

## ON THE STRUCTURE OF COVERS OF SOFIC SHIFTS

RUNE JOHANSEN

Received: June 6, 2010

Revised: March 1, 2011

Communicated by Joachim Cuntz

ABSTRACT. A canonical cover generalizing the left Fischer cover to arbitrary sofic shifts is introduced and used to prove that the left Krieger cover and the past set cover of a sofic shift can be divided into natural layers. These results are used to find the range of a flow-invariant and to investigate the ideal structure of the universal  $C^*$ -algebra associated to a sofic shift space.

2010 Mathematics Subject Classification: Primary: 37B10; Secondary: 46L05

Keywords and Phrases: shift space, sofic shift, Krieger cover, Fischer cover

## 1 INTRODUCTION

Shifts of finite type have been completely classified up to flow equivalence by Boyle and Huang [4, 6, 15], but very little is known about the classification of the class of sofic shift spaces introduced by Weiss [38], even though they are a natural first generalization of shifts of finite type. The purpose of this paper is to investigate the structure of - and relationships between - various standard presentations (the Fischer cover, the Krieger cover, and the past set cover) of sofic shift spaces. These results are used to find the range of the flow-invariant introduced in [1], and to investigate the ideal structure of the  $C^*$ -algebras associated to sofic shifts. In this way, the present paper can be seen as a continuation of the strategy applied in [10, 11, 30], where invariants for shift spaces are extracted from the associated  $C^*$ -algebras.

Section 2 recalls the definitions of shift spaces, labelled graphs, and covers to make the paper self contained. Section 3 introduces a canonical and flow-invariant cover generalizing the left Fischer cover to arbitrary sofic shifts.

Section 4 introduces the concept of a foundation of a cover, which is used to prove that the left Krieger cover and the past set cover can be divided into natural layers and to show that the left Krieger cover of an arbitrary sofic shift can be identified with a subgraph of the past set cover.

In Section 5, the structure of the layers of the left Krieger cover of an irreducible sofic shift is used to find the range of the flow-invariant introduced in [1]. Section 6 uses the results about the structure of covers of sofic shifts to investigate ideal lattices of the associated  $C^*$ -algebras. Additionally, it is proved that Condition (\*) introduced by Carlsen and Matsumoto [12] holds if and only if the left Krieger cover is the maximal essential subgraph of the past set cover.

ACKNOWLEDGEMENTS. This work was supported by the Danish National Research Foundation (DNRF) through the Centre for Symmetry and Deformation. The author would like to thank David Pask, Toke Meier Carlsen, and Søren Eilers for interesting discussions and helpful comments. The author would also like to thank the anonymous referee for useful comments improving the exposition and to thank the University of Wollongong, Australia and the University of Tokyo, Japan where parts of the research for this paper were carried out during visits funded by *Rejselegat for Matematikere*.

## 2 BACKGROUND

SHIFT SPACES. Here, a short introduction to the definition and properties of shift spaces is given to make the present paper self-contained; for a thorough treatment of shift spaces see [21]. Let  $\mathcal{A}$  be a finite set with the discrete topology. The *full shift* over  $\mathcal{A}$  consists of the space  $\mathcal{A}^{\mathbb{Z}}$  endowed with the product topology and the *shift map*  $\sigma: \mathcal{A}^{\mathbb{Z}} \rightarrow \mathcal{A}^{\mathbb{Z}}$  defined by  $\sigma(x)_i = x_{i+1}$  for all  $i \in \mathbb{Z}$ . Let  $\mathcal{A}^*$  be the collection of finite words (also known as blocks) over  $\mathcal{A}$ . A subset  $X \subseteq \mathcal{A}^{\mathbb{Z}}$  is called a *shift space* if it is invariant under the shift map and closed. For each  $\mathcal{F} \subseteq \mathcal{A}^*$ , define  $X_{\mathcal{F}}$  to be the set of bi-infinite sequences in  $\mathcal{A}^{\mathbb{Z}}$  which do not contain any of the *forbidden words* from  $\mathcal{F}$ . A subset  $X \subseteq \mathcal{A}^{\mathbb{Z}}$  is a shift space if and only if there exists  $\mathcal{F} \subseteq \mathcal{A}^*$  such that  $X = X_{\mathcal{F}}$  (cf. [21, Proposition 1.3.4]).  $X$  is said to be a *shift of finite type* (SFT) if this is possible for a finite set  $\mathcal{F}$ .

The *language* of a shift space  $X$  is defined to be the set of all words which occur in at least one  $x \in X$ , and it is denoted  $\mathcal{B}(X)$ .  $X$  is said to be *irreducible* if there for every  $u, w \in \mathcal{B}(X)$  exists  $v \in \mathcal{B}(X)$  such that  $uvw \in \mathcal{B}(X)$ . For each  $x \in X$ , define the *left-ray* of  $x$  to be  $x^- = \cdots x_{-2}x_{-1}$  and define the *right-ray* of  $x$  to be  $x^+ = x_0x_1x_2 \cdots$ . The sets of all left-rays and all right-rays are, respectively, denoted  $X^-$  and  $X^+$ .

A bijective, continuous, and shift commuting map between two shift spaces is called a *conjugacy*, and when such a map exists, the two shift spaces are said to be *conjugate*. Shift spaces  $(X, \sigma_X)$  and  $(Y, \sigma_Y)$  are said to be *flow equivalent* if the corresponding suspension flows  $SX$  and  $SY$  are topologically equivalent. Flow equivalence is generated by conjugacy and *symbol expansion* [35].

GRAPHS. For countable sets  $E^0$  and  $E^1$ , and maps  $r, s: E^1 \rightarrow E^0$  the quadruple  $E = (E^0, E^1, r, s)$  is called a *directed graph*. The elements of  $E^0$  and  $E^1$  are, respectively, the vertices and the edges of the graph. For each edge  $e \in E^1$ ,  $s(e)$  is the vertex where  $e$  starts, and  $r(e)$  is the vertex where  $e$  ends. A *path*  $\lambda = e_1 \cdots e_n$  is a sequence of edges such that  $r(e_i) = s(e_{i+1})$  for all  $i \in \{1, \dots, n-1\}$ . For each  $n \in \mathbb{N}_0$ , the set of paths of length  $n$  is denoted  $E^n$ , and the set of all finite paths is denoted  $E^*$ . Extend the maps  $r$  and  $s$  to  $E^*$  by defining  $s(e_1 \cdots e_n) = s(e_1)$  and  $r(e_1 \cdots e_n) = r(e_n)$ . A *circuit* is a path  $\lambda$  with  $r(\lambda) = s(\lambda)$  and  $|\lambda| > 0$ . For  $u, v \in E^0$ ,  $u$  is said to be *connected* to  $v$  if there is a path  $\lambda \in E^*$  such that  $s(\lambda) = u$  and  $r(\lambda) = v$ , and this is denoted by  $u \geq v$  [21, Section 4.4]. A vertex is said to be *maximal*, if it is connected to all other vertices.  $E$  is said to be *irreducible* if all vertices are maximal. If  $E$  has a unique maximal vertex, this vertex is said to be the *root* of  $E$ .  $E$  is said to be *essential* if every vertex emits and receives an edge. For a finite essential directed graph  $E$ , the *edge shift*  $(X_E, \sigma_E)$  is defined by

$$X_E = \{x \in (E^1)^{\mathbb{Z}} \mid r(x_i) = s(x_{i+1}) \text{ for all } i \in \mathbb{Z}\}.$$

A *labelled graph*  $(E, \mathcal{L})$  over an alphabet  $\mathcal{A}$  consists of a directed graph  $E$  and a surjective labelling map  $\mathcal{L}: E^1 \rightarrow \mathcal{A}$ . Extend the labelling map to  $\mathcal{L}: E^* \rightarrow \mathcal{A}^*$  by defining  $\mathcal{L}(e_1 \cdots e_n) = \mathcal{L}(e_1) \cdots \mathcal{L}(e_n) \in \mathcal{A}^*$ . For a finite essential labelled graph  $(E, \mathcal{L})$ , define the shift space  $(X_{(E, \mathcal{L})}, \sigma)$  by

$$X_{(E, \mathcal{L})} = \{(\mathcal{L}(x_i))_i \in \mathcal{A}^{\mathbb{Z}} \mid x \in X_E\}.$$

The labelled graph  $(E, \mathcal{L})$  is said to be a *presentation* of the shift space  $X_{(E, \mathcal{L})}$ , and a *representative* of a word  $w \in \mathcal{B}(X_{(E, \mathcal{L})})$  is a path  $\lambda \in E^*$  such that  $\mathcal{L}(\lambda) = w$ . Representatives of rays are defined analogously. If  $H \subseteq E^0$  then the *subgraph of  $(E, \mathcal{L})$  induced by  $H$*  is the labelled subgraph of  $(E, \mathcal{L})$  with vertices  $H$  and edges  $\{e \in E^1 \mid s(e), r(e) \in H\}$ .

SOFIC SHIFTS. A function  $\pi: X_1 \rightarrow X_2$  between shift spaces  $X_1$  and  $X_2$  is said to be a *factor map* if it is continuous, surjective, and shift commuting. A shift space is called *sofic* [38] if it is the image of an SFT under a factor map. A shift space is sofic if and only if it can be presented by a finite labelled graph [14]. A sofic shift space is irreducible if and only if it can be presented by an irreducible labelled graph (see [21, Section 3.1]). Let  $(E, \mathcal{L})$  be a finite labelled graph and let  $\pi_{\mathcal{L}}: X_E \rightarrow X_{(E, \mathcal{L})}$  be the factor map induced by the labelling map  $\mathcal{L}: E^1 \rightarrow \mathcal{A}$  then the SFT  $X_E$  is called a *cover* of the sofic shift  $X_{(E, \mathcal{L})}$ , and  $\pi_{\mathcal{L}}$  is called the covering map.

A presentation  $(E, \mathcal{L})$  of a sofic shift space  $X$  is said to be *left-resolving* if no vertex in  $E^0$  receives two edges with the same label. Fischer proved [14] that, up to labelled graph isomorphism, every irreducible sofic shift has a unique left-resolving presentation with fewer vertices than any other left-resolving presentation. This is called the *left Fischer cover* of  $X$ , and it is denoted  $(F, \mathcal{L}_F)$ . An irreducible sofic shift is said to have *almost finite type* (AFT) [22, 33] if the left Fischer cover is right-closing (see e.g. [21, Definition 5.1.4]).

For  $x \in \mathcal{B}(X) \cup X^+$ , define the *predecessor set* of  $x$  to be the set of left-rays which may precede  $x$  in  $X$  (see [17, Sections I and III] and [21, Exercise 3.2.8]). The *follower set* of a left-ray or word is defined analogously. Let  $(E, \mathcal{L})$  be a labelled graph presenting  $X$  and let  $v \in E^0$ . Define the *predecessor set* of  $v$  to be the set of left-rays in  $X$  which have a presentation terminating at  $v$ . This is denoted  $P_\infty^E(v)$ , or just  $P_\infty(v)$  when  $(E, \mathcal{L})$  is understood from the context. The presentation  $(E, \mathcal{L})$  is said to be *predecessor-separated* if  $P_\infty^E(u) \neq P_\infty^E(v)$  when  $u, v \in E^0$  and  $u \neq v$ .

The *left Krieger cover* of the shift space  $X$  is the labelled graph  $(K, \mathcal{L}_K)$  where  $K^0 = \{P_\infty(x^+) \mid x^+ \in X^+\}$ , and where there is an edge labelled  $a \in \mathcal{A}$  from  $P \in K^0$  to  $P' \in K^0$  if and only if there exists  $x^+ \in X^+$  such that  $P = P_\infty(ax^+)$  and  $P' = P_\infty(x^+)$ . The *past set cover* of the shift space  $X$  is the labelled graph  $(W, \mathcal{L}_W)$  where  $W^0 = \{P_\infty(w) \mid w \in \mathcal{B}(X)\}$  and where the edges and labels are constructed as in the Krieger cover. A shift space is sofic if and only if the number of predecessor sets is finite [19, §2], so the left Krieger cover is finite exactly when the shift space is sofic. The left Fischer cover, the left Krieger cover, and the past set cover are left-resolving and predecessor-separated presentations of  $X$ .

The right Krieger cover and the future set cover are right-resolving and follower-separated covers defined analogously to the left Krieger cover and the past set cover, respectively. Every result developed for left-resolving covers in the following has an analogue for the corresponding right-resolving cover. These results can easily be obtained by considering the transposed shift space  $X^T$  (see e.g. [21, p. 39]).

### 3 GENERALIZING THE FISCHER COVER

Jonoska [16] proved that a reducible sofic shift does not necessarily have a unique minimal left-resolving presentation. The aim of this section is to define a generalization of the left Fischer cover as the subgraph of the left Krieger cover induced by a certain subset of vertices. Let  $X$  be a sofic shift space, and let  $(K, \mathcal{L}_K)$  be the left Krieger cover of  $X$ . A predecessor set  $P \in K^0$  is said to be *non-decomposable* if  $V \subseteq K^0$  and  $P = \bigcup_{Q \in V} Q$  implies that  $P \in V$ .

**LEMMA 3.1.** *If  $P \in K^0$  is non-decomposable then the subgraph of  $(K, \mathcal{L}_K)$  induced by  $K^0 \setminus \{P\}$  is not a presentation of  $X$ .*

*Proof.* Let  $E$  be the subgraph of  $K$  induced by  $K^0 \setminus \{P\}$ . Choose  $x^+ \in X^+$  such that  $P = P_\infty(x^+)$ . Let  $V \subseteq K^0 \setminus \{P\}$  be the set of vertices where a presentation of  $x^+$  can start. Then  $Q \subseteq P_\infty(x^+) = P$  for each  $Q \in V$ , and by assumption, there exists  $y^- \in P \setminus \bigcup_{Q \in V} Q$ . Hence, there is no presentation of  $y^-x^+$  in  $(E, \mathcal{L}_K|_E)$ .  $\square$

Lemma 3.1 shows that a subgraph of the left Krieger cover which presents the same shift must contain all the non-decomposable vertices. The next example shows that this subgraph is not always large enough.




Figure 1: Left Krieger cover of the shift considered in Example 3.2. Note that the labelled graph is no longer a presentation of the same shift if the decomposable predecessor set  $P = P_1 \cup P_2$  is removed.

EXAMPLE 3.2. It is easy to check that the labelled graph in Figure 1 is the left Krieger cover of a reducible sofic shift  $X$ . Note that the predecessor set  $P$  is decomposable since  $P = P_1 \cup P_2$ , and that the graph obtained by removing the vertex  $P$  and all edges starting at or terminating at  $P$  is not a presentation of the same sofic shift since there is no presentation of  $f^\infty dbjk^\infty$  in this graph. Note that there is a path from  $P$  to the vertex  $P'$  which is non-decomposable. Together with Lemma 3.1, this example motivates the following definition.

DEFINITION 3.3. The generalized left Fischer cover  $(G, \mathcal{L}_G)$  of a sofic shift  $X$  is defined to be the subgraph of the left Krieger cover induced by  $G^0 = \{P \in K^0 \mid P \geq P', P' \text{ non-decomposable}\}$ .

The following proposition justifies the term generalized left Fischer cover.

PROPOSITION 3.4.

- (i) The generalized left Fischer cover of a sofic shift  $X$  is a left-resolving and predecessor-separated presentation of  $X$ .
- (ii) If  $X$  is an irreducible sofic shift then the generalized left Fischer cover is isomorphic to the left Fischer cover.
- (iii) If  $X_1, X_2$  are sofic shifts with disjoint alphabets then the generalized left Fischer cover of  $X_1 \cup X_2$  is the disjoint union of the generalized left Fischer covers of  $X_1$  and  $X_2$ .

*Proof.* Given  $y^- \in X^-$ , choose  $x^+ \in X^+$  such that  $y^- \in P_\infty(x^+) = P$ . By definition of the generalized left Fischer cover, there exist vertices  $P_1, \dots, P_n \in G^0$  such that  $P = \bigcup_{i=1}^n P_i$ . Choose  $i$  such that  $y^- \in P_i$ . By construction, the left Krieger cover contains a path labelled  $y^-$  terminating at  $P_i$ . Since

$P_i \in G^0$ , this is also a path in the generalized left Fischer cover. This proves that the generalized left Fischer cover is a presentation of  $X^-$ , and hence also a presentation of  $X$ . Since the left Krieger cover is left-resolving and predecessor-separated, so is the generalized left Fischer cover.

Let  $X$  be an irreducible sofic shift, and identify the left Fischer cover  $(F, \mathcal{L}_F)$  with the top irreducible component of the left Krieger cover  $(K, \mathcal{L}_K)$  [19, Lemma 2.7]. By the construction of the generalized left Fischer cover, it follows that the left Fischer cover is a subgraph of the generalized left Fischer cover. Let  $x^+ \in X^+$  such that  $P = P_\infty(x^+)$  is non-decomposable. Let  $S \subseteq F^0$  be the set of vertices where a presentation of  $x^+$  in  $(F, \mathcal{L}_F)$  can start. Then  $P = \bigcup_{v \in S} P_\infty(v)$ , so  $P \in S \subseteq F^0$  by assumption.

Since  $X_1$  and  $X_2$  have no letters in common, the left Krieger cover of  $X_1 \cup X_2$  is just the disjoint union of the left Krieger covers of  $X_1$  and  $X_2$ . The generalized left Fischer cover inherits this property from the left Krieger cover.  $\square$

The shift consisting of two non-interacting copies of the even shift is a simple example where the generalized left Fischer cover is a proper subgraph of the left Krieger cover.

LEMMA 3.5. *Let  $X$  be a sofic shift with left Krieger cover  $(K, \mathcal{L}_K)$ . If there is an edge labelled  $a$  from a non-decomposable  $P \in K^0$  to a decomposable  $Q \in K^0$  then there exists a non-decomposable  $Q' \in K^0$  and an edge labelled  $a$  from  $P$  to  $Q'$ .*

*Proof.* Choose  $x^+ \in X^+$  such that  $P = P_\infty(ax^+)$  and  $Q = P_\infty(x^+)$ . Since  $Q$  is decomposable, there exist  $n > 1$  and non-decomposable  $Q_1, \dots, Q_n \in K^0 \setminus \{Q\}$  such that  $Q = Q_1 \cup \dots \cup Q_n$ . Let  $S$  be the set of predecessor sets  $P' \in K^0$  for which there is an edge labelled  $a$  from  $P'$  to  $Q_j$  for some  $1 \leq j \leq n$ . Given  $y^- \in P$ ,  $y^-ax^+ \in X$ , so  $y^-a \in Q$ . Choose  $1 \leq i \leq n$  such that  $y^-a \in Q_i$ . By construction, there exists  $P' \in S$  such that  $y^- \in P'$ . Reversely, if  $y^- \in P' \in S$  then there is an edge labelled  $a$  from  $P'$  to  $Q_i$  for some  $1 \leq i \leq n$ , so  $y^-a \in Q_i \subseteq Q$ . This implies that  $y^-ax^+ \in X$ , so  $y^- \in P$ . Thus  $P = \bigcup_{P' \in S} P'$ , but  $P$  is non-decomposable, so this means that  $P \in S$ . Hence, there is an edge labelled  $a$  from  $P$  to  $Q_i$  for some  $i$ , and  $Q_i$  is non-decomposable.  $\square$

The following proposition is an immediate consequence of this result and the definition of the generalized left Fischer cover.

PROPOSITION 3.6. *The generalized left Fischer cover is essential.*

The left Fischer cover of an irreducible sofic shift  $X$  is minimal in the sense that no other left-resolving presentation of  $X$  has fewer vertices. This is not always the case for the generalized left Fischer cover.

CANONICAL. Krieger proved that a conjugacy  $\Phi: X_1 \rightarrow X_2$  between sofic shifts with left Krieger covers  $(K_1, \mathcal{L}_1)$  and  $(K_2, \mathcal{L}_2)$ , respectively, induces a conjugacy  $\varphi: \mathcal{X}_{K_1} \rightarrow \mathcal{X}_{K_2}$  such that  $\Phi \circ \pi_1 = \pi_2 \circ \varphi$  when  $\pi_i: \mathcal{X}_{K_i} \rightarrow X_i$  is the

covering map of the left Krieger cover of  $X_i$  [19]. A cover with this property is said to be *canonical*. The next goal is to prove that the generalized left Fischer cover is canonical. This will be done by using results and methods used by Nasu [34] to prove that the left Krieger cover is canonical.

**DEFINITION 3.7** (Bipartite code). *When  $\mathcal{A}, \mathcal{C}, \mathcal{D}$  are alphabets, an injective map  $f: \mathcal{A} \rightarrow \mathcal{CD}$  is called a bipartite expression. If  $X_1, X_2$  are shift spaces with alphabets  $\mathcal{A}_1$  and  $\mathcal{A}_2$ , respectively, and if  $f_1: \mathcal{A}_1 \rightarrow \mathcal{CD}$  is a bipartite expression then a map  $\Phi: X_1 \rightarrow X_2$  is said to be a bipartite code induced by  $f_1$  if there exists a bipartite expression  $f_2: \mathcal{A}_2 \rightarrow \mathcal{DC}$  such that one of the following two conditions is satisfied:*

- (i) *If  $x \in X_1$ ,  $y = \Phi(x)$ , and  $f_1(x_i) = c_i d_i$  with  $c_i \in \mathcal{C}$  and  $d_i \in \mathcal{D}$  for all  $i \in \mathbb{Z}$  then  $f_2(y_i) = d_i c_{i+1}$  for all  $i \in \mathbb{Z}$ .*
- (ii) *If  $x \in X_1$ ,  $y = \Phi(x)$ , and  $f_1(x_i) = c_i d_i$  with  $c_i \in \mathcal{C}$  and  $d_i \in \mathcal{D}$  for all  $i \in \mathbb{Z}$  then  $f_2(y_i) = d_{i-1} c_i$  for all  $i \in \mathbb{Z}$ .*

*A mapping  $\Phi: X_1 \rightarrow X_2$  is called a bipartite code, if it is the bipartite code induced by some bipartite expression.*

It is clear that a bipartite code is a conjugacy and that the inverse of a bipartite code is a bipartite code.

**THEOREM 3.8** (Nasu [34, Thm. 2.4]). *Any conjugacy between shift spaces can be decomposed into a product of bipartite codes.*

Let  $\Phi: X_1 \rightarrow X_2$  be a bipartite code corresponding to bipartite expressions  $f_1: \mathcal{A}_1 \rightarrow \mathcal{CD}$  and  $f_2: \mathcal{A}_2 \rightarrow \mathcal{DC}$ , and use the bipartite expressions to recode  $X_1$  and  $X_2$  to

$$\begin{aligned}\hat{X}_1 &= \{(f_1(x_i))_i \mid x \in X_1\} \subseteq (\mathcal{CD})^{\mathbb{Z}} \\ \hat{X}_2 &= \{(f_2(x_i))_i \mid x \in X_2\} \subseteq (\mathcal{DC})^{\mathbb{Z}}.\end{aligned}$$

For  $i \in \{1, 2\}$ ,  $f_i$  induces a one-block conjugacy from  $X_i$  to  $\hat{X}_i$ , and  $\Phi$  induces a bipartite code  $\hat{\Phi}: \hat{X}_1 \rightarrow \hat{X}_2$  which commutes with these conjugacies. If  $\Phi$  satisfies condition (i) in the definition of a bipartite code then  $(\hat{\Phi}(\hat{x}))_i = d_i c_{i+1}$  when  $\hat{x} = (c_i d_i)_{i \in \mathbb{Z}} \in \hat{X}_1$ . If it satisfies condition (ii) then  $(\hat{\Phi}(\hat{x}))_i = d_{i-1} c_i$  when  $\hat{x} = (c_i d_i)_{i \in \mathbb{Z}} \in \hat{X}_1$ . The shifts  $\hat{X}_1$  and  $\hat{X}_2$  will be called the *recoded shifts* of the bipartite code, and  $\hat{\Phi}$  will be called the *recoded bipartite code*.

A labelled graph  $(G, \mathcal{L})$  is said to be *bipartite* if  $G$  is a bipartite graph (i.e. the vertex set can be partitioned into two sets  $(G^0)_1$  and  $(G^0)_2$  such that no edge has its range and source in the same set). When  $(G, \mathcal{L})$  is a bipartite labelled graph over an alphabet  $\mathcal{A}$ , define two graphs  $G_1$  and  $G_2$  as follows: For  $i \in \{1, 2\}$ , the vertex set of  $G_i$  is  $(G^0)_i$ , the edge set is the set of paths of length 2 in  $(G, \mathcal{L})$  for which both range and source are in  $(G^0)_i$ , and the range and source maps are inherited from  $G$ . For  $i \in \{1, 2\}$ , define  $\mathcal{L}_i: G_i^1 \rightarrow \mathcal{A}^2$

by  $\mathcal{L}_i(e_f) = \mathcal{L}(e)\mathcal{L}(f)$ . The pair  $(G_1, \mathcal{L}_1), (G_2, \mathcal{L}_2)$  is called the *induced pair of labelled graphs* of  $(G, \mathcal{L})$ . This decomposition is not necessarily unique, but whenever a bipartite labelled graph is mentioned, it will be assumed that the induced graphs are specified.

REMARK 3.9 (Nasu [34, Remark 4.2]). Let  $(G, \mathcal{L})$  be a bipartite labelled graph for which the induced pair of labelled graphs is  $(G_1, \mathcal{L}_1), (G_2, \mathcal{L}_2)$ . Let  $X_1$  and  $X_2$  be the sofic shifts presented by these graphs, and let  $X_{G_1}, X_{G_2}$  be the edge shifts generated by  $G_1, G_2$ . The natural embedding  $f: G_1^1 \rightarrow (G^1)^2$  is a bipartite expression which induces two bipartite codes  $\varphi_{\pm}: X_{G_1} \rightarrow X_{G_2}$  such that  $(\varphi_+(x))_i = f_i e_{i+1}$  and  $(\varphi_-(x))_i = f_{i-1} e_i$  when  $x = (e_i f_i)_{i \in \mathbb{Z}} \in X_{G_1}$ . Similarly, the embedding  $F: \mathcal{L}_1(G_1^1) \rightarrow (\mathcal{L}(G^1))^2$  is a bipartite expression which induces bipartite codes  $\Phi_{\pm}: X_1 \rightarrow X_2$  such that  $(\Phi_+(x))_i = b_i a_{i+1}$  and  $(\Phi_-(x))_i = b_{i-1} a_i$  when  $x = (a_i b_i)_{i \in \mathbb{Z}} \in X_1$ . By definition,  $\Phi_{\pm} \circ \pi_1 = \pi_2 \circ \varphi_{\pm}$  when  $\pi_1: X_{G_1} \rightarrow X_1, \pi_2: X_{G_2} \rightarrow X_2$  are the covering maps. The bipartite codes  $\varphi_{\pm}$  and  $\Phi_{\pm}$  are called the *standard bipartite codes induced by  $(G, \mathcal{L})$* .

LEMMA 3.10 (Nasu [34, Cor. 4.6 (1)]). Let  $\Phi: X_1 \rightarrow X_2$  be a bipartite code between sofic shifts  $X_1$  and  $X_2$ . Let  $\hat{X}_1$  and  $\hat{X}_2$  be the recoded shifts of  $X_1$  and  $X_2$  respectively, and let  $(K_1, \mathcal{L}_1)$  and  $(K_2, \mathcal{L}_2)$  be the left Krieger covers of  $\hat{X}_1$  and  $\hat{X}_2$  respectively. Then there exists a sofic shift  $\hat{X}$  for which the left Krieger cover is a bipartite labelled graph such that the induced pair of labelled graphs is  $(K_1, \mathcal{L}_1), (K_2, \mathcal{L}_2)$  and such that the recoded bipartite code  $\hat{\Phi}: \hat{X}_1 \rightarrow \hat{X}_2$  of  $\Phi$  is one of the standard bipartite codes  $\Phi_{\pm}$  induced by the left Krieger cover of  $\hat{X}$  as defined in Remark 3.9.

The proof of the following theorem is very similar to the proof of the corresponding result by Nasu [34, Thm. 3.3] for the left Krieger cover.

THEOREM 3.11. *The generalized left Fischer cover is canonical.*

*Proof.* Let  $\Phi: X_1 \rightarrow X_2$  be a bipartite code. Let  $\hat{X}_1, \hat{X}_2$  be the recoded shifts, let  $(K_1, \mathcal{L}_1), (K_2, \mathcal{L}_2)$  be the corresponding left Krieger covers, and let  $\hat{\Phi}: \hat{X}_1 \rightarrow \hat{X}_2$  be the recoded bipartite code. Use Lemma 3.10 to find a sofic shift  $\hat{X}$  such that the left Krieger cover  $(K, \mathcal{L})$  of  $\hat{X}$  is a bipartite labelled graph for which the induced pair of labelled graphs is  $(K_1, \mathcal{L}_1), (K_2, \mathcal{L}_2)$ . Let  $(G_1, \mathcal{L}_1), (G_2, \mathcal{L}_2)$ , and  $(G, \mathcal{L})$  be the generalized left Fischer covers of respectively  $\hat{X}_1, \hat{X}_2$ , and  $\hat{X}$ .

The labelled graph  $(G, \mathcal{L})$  is bipartite since  $G$  is a subgraph of  $K$ . Note that a predecessor set  $P$  in  $K_1^0$  or  $K_2^0$  is decomposable if and only if the corresponding predecessor set in  $K^0$  is decomposable. If  $i \in \{1, 2\}$  and  $Q \in G_i^0 \subseteq K_i^0$  then there is a path in  $K_i$  from  $Q$  to a non-decomposable  $P \in K_i^0$ . By considering the corresponding path in  $K$ , it is clear that the vertex in  $K^0$  corresponding to  $Q$  is in  $G^0$ . Conversely, if  $Q \in G^0$  then there is a path in  $K$  from  $Q$  to a non-decomposable  $P \in K^0$ . If  $P$  and  $Q$  belong to the same partition  $K_i^0$  then the vertex in  $K_i$  corresponding to  $Q$  is in  $G_i^0$  by definition. On the other hand, if  $Q$  corresponds to a vertex in  $K_i$  and if  $P$  belongs to the other partition

then Lemma 3.5 shows that there exists a non-decomposable  $P'$  in the same partition as  $Q$  and an edge from  $P$  to  $P'$  in  $K$ . Hence, there is also a path in  $K_i$  from the vertex corresponding to  $Q$  to the vertex corresponding to  $P'$ , so  $Q \in G_i^0$ . This proves that the pair of induced labelled graphs of  $(G, \mathcal{L})$  is  $(G_1, \mathcal{L}_1), (G_2, \mathcal{L}_2)$ .

Let  $\hat{\Psi}_{\pm}: \hat{X}_1 \rightarrow \hat{X}_2$  be the standard bipartite codes induced by  $(G, \mathcal{L})$ . Remark 3.9 shows that there exist bipartite codes  $\hat{\psi}_{\pm}: X_{G_1} \rightarrow X_{G_2}$  such that  $\hat{\Psi}_{\pm} \circ \hat{\pi}_1|_{X_{G_1}} = \hat{\pi}_2|_{X_{G_2}} \circ \hat{\psi}_{\pm}$ . The labelled graph  $(G, \mathcal{L})$  presents the same sofic shift as  $(K, \mathcal{L})$ , so they both induce the same standard bipartite codes from  $\hat{X}_1$  to  $\hat{X}_2$ , and by Lemma 3.10,  $\hat{\Phi}$  is one of these standard bipartite codes, so  $\hat{\Phi} = \hat{\Psi}_+$  or  $\hat{\Phi} = \hat{\Psi}_-$ . In particular, there exists a bipartite code  $\hat{\psi}: X_{G_1} \rightarrow X_{G_2}$  such that  $\hat{\Phi} \circ \hat{\pi}_1|_{X_{G_1}} = \hat{\pi}_2|_{X_{G_2}} \circ \hat{\psi}$ .

By recoding  $\hat{X}_1$  to  $X_1$  and  $\hat{X}_2$  to  $X_2$  via the bipartite expressions inducing  $\Phi$ , this gives a bipartite code  $\psi$  such that  $\Phi \circ \pi_1 = \pi_2 \circ \psi$  when  $\pi_1, \pi_2$  are the covering maps of the generalized left Fischer covers of  $X_1$  and  $X_2$  respectively. By Theorem 3.8, any conjugacy can be decomposed as a product of bipartite codes, so this proves that the generalized left Fischer cover is canonical.  $\square$

**THEOREM 3.12.** *For  $i \in \{1, 2\}$ , let  $X_i$  be a sofic shift with generalized left Fischer cover  $(G_i, \mathcal{L}_i)$  and covering map  $\pi_i: X_{G_i} \rightarrow X_i$ . If  $\Phi: SX_1 \rightarrow SX_2$  is a flow equivalence then there exists a unique flow equivalence  $\varphi: SX_{G_1} \rightarrow SX_{G_2}$  such that  $\Phi \circ S\pi_1 = S\pi_2 \circ \varphi$ .*

*Proof.* In [5] it is proved that the left Krieger cover respects symbol expansion: If  $X$  is a sofic shift with alphabet  $\mathcal{A}$ ,  $a \in \mathcal{A}$ ,  $\bullet$  is some symbol not in  $\mathcal{A}$ , and if  $\hat{X}$  is obtained from  $X$  via a symbol expansion which inserts a  $\bullet$  after each  $a$  then the left Krieger cover of  $\hat{X}$  is obtained by replacing each edge labelled  $a$  in the left Krieger cover of  $X$  by two edges in sequence labelled  $a$  and  $\bullet$  respectively. Clearly, the generalized left Fischer cover inherits this property. By [5], any canonical cover which respects flow equivalence has the desired property, so the result follows from Theorem 3.11.  $\square$

#### 4 FOUNDATIONS AND LAYERS OF COVERS

Let  $\mathcal{E} = (E, \mathcal{L})$  be a finite left-resolving and predecessor-separated labelled graph. For each  $V \subseteq E^0$  and each word  $w$  over the alphabet  $\mathcal{A}$  of  $\mathcal{L}$  define

$$wV = \{u \in E^0 \mid u \text{ is the source of a path labelled } w \text{ terminating in } V\}.$$

**DEFINITION 4.1.** *Let  $S$  be a subset of the power set  $\mathcal{P}(E^0)$ , and let  $\sim$  be an equivalence relation on  $S$ . The pair  $(S, \sim)$  is said to be past closed if*

- $\{v\} \in S$ ,
- $\{u\} \sim \{v\}$  implies  $u = v$ ,
- $aV \neq \emptyset$  implies  $aV \in S$ , and

- $U \sim V$  and  $aU \neq \emptyset$  implies  $aV \neq \emptyset$  and  $aU \sim aV$

for all  $u, v \in E^0$ ,  $U, V \in S$ , and  $a \in \mathcal{A}$ .

Let  $(S, \sim)$  be past closed. For each  $V \in S$ , let  $[V]$  denote the equivalence class of  $V$  with respect to  $\sim$ . When  $a \in \mathcal{A}$  and  $V \in S$ ,  $[V]$  is said to receive  $a$  if  $aV \neq \emptyset$ . For each  $[V] \in S/\sim$ , define  $|[V]| = \min_{V \in [V]} |V|$ .

DEFINITION 4.2. Define  $\mathcal{G}(\mathcal{E}, S, \sim)$  to be the labelled graph with vertex set  $S/\sim$  for which there is an edge labelled  $a$  from  $[aV]$  to  $[V]$  whenever  $[V]$  receives  $a$ . For each  $n \in \mathbb{N}$ , the  $n$ th layer of  $\mathcal{G}(\mathcal{E}, S, \sim)$  is the labelled subgraph induced by  $S_n = \{[V] \in S/\sim \mid n = |[V]|\}$ .  $\mathcal{E}$  is said to be a foundation of any labelled graph isomorphic to  $\mathcal{G}(\mathcal{E}, S, \sim)$ .

If a labelled graph  $\mathcal{H}$  is isomorphic to  $\mathcal{G}(\mathcal{E}, S, \sim)$  then the subgraph of  $\mathcal{H}$  corresponding to the  $n$ th layer of  $\mathcal{G}(\mathcal{E}, S, \sim)$  is said to be the  $n$ th layer of  $\mathcal{H}$  with respect to  $\mathcal{E}$ , or simply the  $n$ th layer if  $\mathcal{E}$  is understood from the context.

PROPOSITION 4.3.  $\mathcal{E}$  and  $\mathcal{G}(\mathcal{E}, S, \sim)$  present the same sofic shift, and  $\mathcal{E}$  is labelled graph isomorphic to the first layer of  $\mathcal{G}(\mathcal{E}, S, \sim)$ .

*Proof.* By assumption, there is a bijection between  $E^0$  and the set of vertices in the first layer of  $\mathcal{G}(\mathcal{E}, S, \sim)$ . By construction, there is an edge labelled  $a$  from  $u$  to  $v$  in  $\mathcal{E}$  if and only if there is an edge labelled  $a$  from  $[\{u\}]$  to  $[\{v\}]$  in  $\mathcal{G}(\mathcal{E}, S, \sim)$ . Every finite word presented by  $\mathcal{G}(\mathcal{E}, S, \sim)$  is also presented by  $\mathcal{E}$ , so they present the same sofic shift.  $\square$

The following proposition motivates the use of the term layer by showing that edges can never go from higher to lower layers.

PROPOSITION 4.4. If  $[V] \in S/\sim$  receives  $a \in \mathcal{A}$  then  $|[aV]| \leq |[V]|$ . If  $\mathcal{G}(\mathcal{E}, S, \sim)$  has an edge from a vertex in the  $m$ th layer to a vertex in the  $n$ th layer then  $m \leq n$ .

*Proof.* Choose  $V \in [V]$  such that  $|V| = |[V]|$ . Each  $u \in aV$  emits at least one edge labelled  $a$  terminating in  $V$ , and  $\mathcal{E}$  is left-resolving, so  $|[aV]| \leq |aV| \leq |V| = |[V]|$ . The second statement follows from the definition of  $\mathcal{G}(\mathcal{E}, S, \sim)$ .  $\square$

EXAMPLE 4.5. Let  $(F, \mathcal{L}_F)$  be the left Fischer cover of an irreducible sofic shift  $X$ . For each  $x^+ \in X^+$ , define  $s(x^+) \subseteq F^0$  to be the set of vertices where a presentation of  $x^+$  can start.  $S = \{s(x^+) \mid x^+ \in X^+\} \subseteq \mathcal{P}(F^0)$  is past closed since each vertex in the left Fischer cover is the predecessor set of an intrinsically synchronizing right-ray, so the multiplicity set cover of  $X$  can be defined to be  $\mathcal{G}((F, \mathcal{L}_F), S, =)$ . An analogous cover can be defined by considering the vertices where presentations of finite words can start. Thomsen [37] constructs the derived shift space  $\partial X$  of  $X$  using right-resolving graphs, but an analogous construction works for left-resolving graphs. The procedure from [37, Example 6.10] shows that this  $\partial X$  is presented by the labelled graph obtained by removing the left Fischer cover from the multiplicity set cover.

Let  $X$  be a sofic shift, and let  $(K, \mathcal{L}_K)$  be the left Krieger cover of  $X$ . In order to use the preceding results to investigate the structure of the left Krieger cover and the past set cover, define an equivalence relation on  $\mathcal{P}(K^0)$  by  $U \sim_{\cup} V$  if and only if  $\bigcup_{P \in U} P = \bigcup_{Q \in V} Q$ . Clearly,  $\{P\} \sim_{\cup} \{Q\}$  if and only if  $P = Q$ . If  $U, V \subseteq K^0$ ,  $a \in \mathcal{A}$ ,  $aV \neq \emptyset$ , and  $U \sim_{\cup} V$  then  $aU \sim_{\cup} aV$  by the definition of the left Krieger cover.

**THEOREM 4.6.** *For a sofic shift  $X$ , the generalized left Fischer cover  $(G, \mathcal{L}_G)$  is a foundation of the left Krieger cover  $(K, \mathcal{L}_K)$ , and no smaller subgraph is a foundation.*

*Proof.* Define  $S = \{V \subseteq G^0 \mid \exists x^+ \in X^+ \text{ such that } P_{\infty}(x^+) = \bigcup_{P \in V} P\}$ . Note that  $\{P\} \in S$  for every  $P \in G^0$ . If  $x^+ \in X^+$  with  $P_{\infty}(x^+) = \bigcup_{P \in V} P$  and if  $aV \neq \emptyset$  for some  $a \in \mathcal{A}$  then  $ax^+ \in X^+$  and  $P_{\infty}(ax^+) = \bigcup_{P \in aV} P$ . This proves that the pair  $(S, \sim_{\cup})$  is past closed, so  $\mathcal{G}((G, \mathcal{L}_G), S, \sim_{\cup})$  is well defined. Since  $(G, \mathcal{L}_G)$  is a presentation of  $X$ , there is a bijection  $\varphi: S/\sim_{\cup} \rightarrow K^0$  defined by  $\varphi([V]) = \bigcup_{P \in V} P$ . By construction, there is an edge labelled  $a$  from  $[U]$  to  $[V]$  in  $\mathcal{G}((G, \mathcal{L}_G), S, \sim_{\cup})$  if and only if there exists  $x^+ \in X^+$  such that  $P_{\infty}(ax^+) = \bigcup_{P \in U} P$  and  $P_{\infty}(x^+) = \bigcup_{Q \in V} Q$ , so  $\mathcal{G}((G, \mathcal{L}_G), S, \sim_{\cup})$  is isomorphic to  $(K, \mathcal{L}_K)$ . It follows from Lemma 3.1 that no proper subgraph of  $(G, \mathcal{L}_G)$  can be a foundation of the left Krieger cover.  $\square$

The example from [12, Section 4] shows that the left Krieger cover can be a proper subgraph of the past set cover. The following lemma will be used to further investigate this relationship.

**LEMMA 4.7.** *Let  $X$  be a sofic shift. For every right-ray  $x^+ = x_1x_2x_3 \dots \in X^+$  there exists  $n \in \mathbb{N}$  such that  $P_{\infty}(x^+) = P_{\infty}(x_1x_2 \dots x_k)$  for all  $k \geq n$ .*

*Proof.* It is clear that  $P_{\infty}(x_1) \supseteq P_{\infty}(x_1x_2) \supseteq \dots \supseteq P_{\infty}(x^+)$ . Since  $X$  is sofic, there are only finitely many different predecessor sets of words, so there must exist  $n \in \mathbb{N}$  such that  $P_{\infty}(x_1x_2 \dots x_k) = P_{\infty}(x_1x_2 \dots x_n)$  for all  $k \geq n$ . If  $y^- \in P_{\infty}(x_1x_2 \dots x_n)$  is given, then  $y^-x_1x_2 \dots x_k \in X$  for all  $k \geq n$ , so  $y^-x^+$  contains no forbidden words, and therefore  $y^- \in P_{\infty}(x^+)$ . Since  $y^-$  was arbitrary,  $P_{\infty}(x^+) = P_{\infty}(x_1x_2 \dots x_n)$ .  $\square$

**THEOREM 4.8.** *For a sofic shift  $X$ , the generalized left Fischer cover  $(G, \mathcal{L}_G)$  and the left Krieger cover  $(K, \mathcal{L}_K)$  are both foundations of the past set cover  $(W, \mathcal{L}_W)$ .*

*Proof.* Define  $S = \{V \subseteq G^0 \mid \exists w \in \mathcal{B}(X) \text{ such that } P_{\infty}(w) = \bigcup_{P \in V} P\}$ , and use Lemma 4.7 to conclude that  $S$  contains  $\{P\}$  for every  $P \in G^0$ . By arguments analogous to the ones used in the proof of Theorem 4.6, it follows that  $\mathcal{G}((G, \mathcal{L}_G), S, \sim_{\cup})$  is isomorphic to  $(W, \mathcal{L}_W)$ . To see that  $(K, \mathcal{L}_K)$  is also a foundation, define  $T = \{V \subseteq K^0 \mid \exists w \in \mathcal{B}(X) \text{ such that } P_{\infty}(w) = \bigcup_{P \in V} P\}$ , and apply arguments analogous to the ones used above to prove that  $(W, \mathcal{L}_W)$  is isomorphic to  $\mathcal{G}((K, \mathcal{L}_K), T, \sim_{\cup})$ .  $\square$




Figure 2: Left Fischer cover of the 3-charge constrained shift.




Figure 3: Left Krieger cover of the 3-charge constrained shift.

In the following, the  $n$ th layer of the left Krieger cover (past set cover) will always refer to the  $n$ th layer with respect to the generalized left Fischer cover  $(G, \mathcal{L}_G)$ . For a right-ray (word)  $x$ ,  $P_\infty(x)$  is a vertex in the  $n$ th layer of the left Krieger cover (past set cover) for some  $n \in \mathbb{N}$ , and such an  $x$  is said to be  $1/n$ -synchronizing. Note that  $x$  is  $1/n$ -synchronizing if and only if  $n$  is the smallest number such that there exist  $P_1, \dots, P_n \in G^0$  with  $\bigcup_{i=1}^n P_i = P_\infty(x)$ . In an irreducible sofic shift with left Fischer cover  $(F, \mathcal{L}_F)$ , this happens if and only if  $n$  is the smallest number such that there exist  $u_1, \dots, u_n \in F^0$  with  $\bigcup_{i=1}^n P_\infty(u_i) = P_\infty(x)$ .

EXAMPLE 4.9. Figures 2 and 3 show, respectively, the left Fischer and the left Krieger cover of the 3-charge constrained shift (see e.g. [21, Example 1.2.7] for the definition of charge constrained shifts). There are 3 vertices in the second layer of the left Krieger cover and two in the third. Note how the left Fischer cover can be identified with the first layer of the left Krieger cover.

COROLLARY 4.10. *If the left Krieger cover of a sofic shift is reducible then so is the past set cover.*

*Proof.* This follows from Proposition 4.4 and Theorem 4.8.  $\square$

## 5 THE RANGE OF A FLOW INVARIANT

Let  $E$  be a directed graph. Vertices  $u, v \in E^0$  *properly communicate* [1] if there are paths  $\mu, \lambda \in E^*$  of length greater than or equal to 1 such that  $s(\mu) = u$ ,  $r(\mu) = v$ ,  $s(\lambda) = v$ , and  $r(\lambda) = u$ . This relation is used to construct maximal disjoint subsets of  $E^0$ , called *proper communication sets of vertices*, such that  $u, v \in E^0$  properly communicate if and only if they belong to the same subset. The *proper communication graph*  $PC(E)$  is defined to be the directed graph for which the vertices are the proper communication sets of vertices of  $E$  and for which there is an edge from one proper communication set to another if and only if there is a path from a vertex in the first set to a vertex in the second. The proper communication graph of the left Krieger cover of a sofic shift space is a flow-invariant [1].

Let  $X$  be an irreducible sofic shift with left Fischer cover  $(F, \mathcal{L}_F)$  and left Krieger cover  $(K, \mathcal{L}_K)$ , and let  $E$  be the proper communication graph of  $K$ . By construction,  $E$  is finite and contains no circuit. The left Fischer cover is isomorphic to an irreducible subgraph of  $(K, \mathcal{L}_K)$  corresponding to a root  $r \in E^0$  [19, Lemma 2.7], and by definition, there is an edge from  $u \in E^0$  to  $v \in E^0$  whenever  $u > v$ . The following proposition gives the range of the flow-invariant by proving that all such graphs can occur.

**PROPOSITION 5.1.** *Let  $E$  be a finite directed graph with a root and without circuits.  $E$  is the proper communication graph of the left Krieger cover of an AFT shift if there is an edge from  $u \in E^0$  to  $v \in E^0$  whenever  $u > v$ .*

*Proof.* Let  $E$  be an arbitrary finite directed graph which contains no circuit and which has a root  $r$ , and let  $\tilde{E}$  be the directed graph obtained from  $E$  by adding an edge from  $u \in E^0$  to  $v \in E^0$  whenever  $u > v$ . The goal is to construct a labelled graph  $(F, \mathcal{L}_F)$  which is the left Fischer cover of an irreducible sofic shift with the desired properties. For each  $v \in E^0$ , let  $l(v)$  be the length of the longest path from  $r$  to  $v$ . This is well-defined since  $E$  does not contain any circuits. For each  $v \in E^0$ , define  $n(v) = 2^{l(v)}$  vertices  $v_1, \dots, v_{n(v)} \in F^0$ . The single vertex corresponding to the root  $r \in E^0$  is denoted  $r_1$ . For each  $v \in E^0$ , draw a loop of length 1 labelled  $a_v$  at each of the vertices  $v_1, \dots, v_{n(v)} \in F^0$ . If there is an edge from  $u \in E^0$  to  $v \in E^0$  then  $l(v) > l(u)$ . From each vertex  $u_1, \dots, u_{n(u)}$  draw  $n(u, v) = n(v)/n(u) = 2^{l(v)-l(u)} \geq 2$  edges labelled  $a_{u,v}^1, \dots, a_{u,v}^{n(u,v)}$  such that every vertex  $v_1, \dots, v_{n(v)}$  receives exactly one of these edges. For each sink  $v \in E^0$  draw a uniquely labelled edge from each vertex  $v_1, \dots, v_{n(v)}$  to  $r_1$ . This finishes the construction of  $(F, \mathcal{L}_F)$ .

By construction,  $F$  is irreducible, right-resolving, and left-resolving. Additionally, it is predecessor-separated because there is a uniquely labelled path to every vertex in  $F^0$  from  $r_1$ . Thus,  $(F, \mathcal{L}_F)$  is the left Fischer cover of an AFT shift  $X$ . Let  $(K, \mathcal{L}_K)$  be the left Krieger cover of  $X$ .

For every  $v \in E^0$ ,  $P_\infty(a_v^\infty) = \bigcup_{i=1}^{n(v)} P_\infty(v_i)$  and no smaller set of vertices has this property, so  $P_\infty(a_v^\infty)$  is a vertex in the  $n(v)$ th layer of the left Krieger cover. There is clearly a loop labelled  $a_v$  at the vertex  $P_\infty(a_v^\infty)$ , so it belongs




Figure 4: A directed graph with root  $r$  and without circuits.




Figure 5: Left Fischer cover of the sofic shift  $X$  considered in Example 5.2.

to a proper communication set of vertices. Furthermore,  $ba_v^\infty \in X^+$  if and only if  $b = a_v$  or  $b = a_{u,v}^i$  for some  $u \in E^0$  and  $1 \leq i \leq n(u, v)$ . By construction,  $P_\infty(a_{u,v}^i a_v^\infty) = \bigcup_{i=1}^{n(u,v)} P_\infty(u_i) = P_\infty(a_u^\infty)$ , so there is an edge from  $P_\infty(a_u^\infty)$  to  $P_\infty(a_v^\infty)$  if and only if there is an edge from  $u$  to  $v$  in  $E$ . This proves that  $E$ , and hence also  $\tilde{E}$ , are subgraphs of the proper communication graph of  $K$ . Since the edges which terminate at  $r_1$  are uniquely labelled, any  $x^+ \in X^+$  which contains one of these letters must be intrinsically synchronizing. If  $x^+ \in X^+$  does not contain any of these letters then  $x^+$  must be eventually periodic with  $x^+ = wa_v^\infty$  for some  $v \in E^0$  and  $w \in \mathcal{B}(X)$ . Thus,  $K$  only has the vertices described above, and therefore the proper communication graph of  $K$  is  $\tilde{E}$ .  $\square$

EXAMPLE 5.2. To illustrate the construction used in the proof of Proposition 5.1, let  $E$  be the directed graph drawn in Figure 4.  $E$  has a unique maximal vertex  $r$  and contains no circuit, so it is the proper communication graph of




Figure 6: Left Krieger cover of the shift space  $X$  considered in Example 5.2. The structure of the irreducible component corresponding to the left Fischer cover has been suppressed.

the left Krieger cover of an irreducible sofic shift. Note that  $l(x) = l(y) = 1$  and that  $l(z) = 2$ . Figure 5 shows the left Fischer cover of a sofic shift  $X$  constructed using the method from the proof of Proposition 5.1. Note that the top and bottom vertices should be identified, and that the labelling of the edges terminating at  $r_1$  has been suppressed. Figure 6 shows the left Krieger cover of  $X$ , but the structure of the irreducible component corresponding to the left Fischer cover has been suppressed to emphasize the structure of the higher layers.

In [1], it was also remarked that an invariant analogous to the one discussed in Proposition 5.1 is obtained by considering the proper communication graph of the right Krieger cover. The following example shows that the two invariants may carry different information.

EXAMPLE 5.3. The labelled graph in Figure 7 is left-resolving, irreducible, and predecessor-separated, so it is the left Fischer cover of an irreducible sofic shift. Similarly, the labelled graph in Figure 8 is irreducible, right-resolving and follower-separated, so it is the right Fischer cover of an irreducible sofic shift. By considering the edges labelled  $d$ , it is easy to see that the two graphs present the same sofic shift space  $X$ .

Every right-ray which contains a letter different from  $a$  or  $a'$  is intrinsically synchronizing, so consider a right-ray  $x^+ \in X^+$  such that  $(x^+)_i \in \{a, a'\}$  for all  $i \in \mathbb{N}$ . By considering Figure 7, it is clear that  $P_\infty(x^+) = P_\infty(u) \cup P_\infty(v) \cup P_\infty(y) = P_\infty(y)$ , so  $P(x^+)$  is also in the first layer of the left Krieger cover. Hence, the proper communication graph has only one vertex and no edges.

Every left-ray containing a letter different from  $a$  or  $a'$  is intrinsically synchronizing, so consider the left-ray  $a^\infty \in X^-$ . Figure 8 shows that  $F_\infty(a^\infty) =$




Figure 7: Left Fischer cover of the irreducible sofic shift  $X$  discussed in Example 5.3.




Figure 8: Right Fischer cover of the irreducible sofic shift  $X$  discussed in Example 5.3.

$F_\infty(u') \cup F_\infty(v')$  and that no single vertex  $y'$  in the right Fischer cover has  $F_\infty(y') = F_\infty(a^\infty)$ , so there is a vertex in the second layer of the right Krieger cover. In particular, the corresponding proper communication graph is non-trivial.

## 6 $C^*$ -ALGEBRAS ASSOCIATED TO SOFIC SHIFT SPACES

Cuntz and Krieger [13] introduced a class of  $C^*$ -algebras which can naturally be viewed as the universal  $C^*$ -algebras associated to shifts of finite type. This was generalized by Matsumoto [23] who associated two  $C^*$ -algebras  $\mathcal{O}_X$  and  $\mathcal{O}_{X^*}$  to every shift space  $X$ , and these Matsumoto algebras have been studied intensely [8, 18, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32]. The two Matsumoto algebras  $\mathcal{O}_X$  and  $\mathcal{O}_{X^*}$  are generated by elements satisfying the same relations,

but they are not isomorphic in general [12]. This paper will follow the approach of Carlsen in [9] where a universal  $C^*$ -algebra  $\mathcal{O}_{\tilde{X}}$  is associated to every one-sided shift space  $\tilde{X}$ . This also gives a way to associate  $C^*$ -algebras to every two-sided shift since a two-sided shift  $X$  corresponds to two one-sided shifts  $X^+$  and  $X^-$ .

**IDEAL LATTICES.** Let  $X$  be a sofic shift space and let  $\mathcal{O}_{X^+}$  be the universal  $C^*$ -algebra associated to the one-sided shift  $X^+$  as defined in [9]. Carlsen proved that  $\mathcal{O}_{X^+}$  is isomorphic to the Cuntz-Krieger algebra of the left Krieger cover of  $X$  [8], so the lattice of gauge invariant ideals in  $\mathcal{O}_{X^+}$  is given by the proper communication graph of the left Krieger cover of  $X$  [3, 20], and all ideals are given in this way if the left Krieger cover satisfies Condition (K) [36, Theorem 4.9]. Hence, Proposition 4.4 and Theorem 4.6 can be used to investigate the ideal lattice of  $\mathcal{O}_{X^+}$ . For a reducible sofic shift, a part of the ideal lattice is given by the structure of the generalized left Fischer cover, which is reducible, but if  $X$  is an irreducible sofic shift, and the left Krieger cover of  $X$  satisfies Condition (K) then the fact that the left Krieger cover has a unique top component implies that  $\mathcal{O}_{X^+}$  will always have a unique maximal ideal. The following proposition shows that all these lattices can be realized.

**PROPOSITION 6.1.** *Any finite lattice of ideals with a unique maximal ideal is the ideal lattice of the universal  $C^*$ -algebra  $\mathcal{O}_{X^+}$  associated to an AFT shift  $X$ .*

*Proof.* Let  $E$  be a finite directed graph without circuits and with a unique maximal vertex. Consider the following slight modification of the algorithm from the proof of Proposition 5.1. For each  $v \in E$ , draw two loops of length 1 at each vertex  $v_1, \dots, v_{n(v)}$  associated to  $v$ : One labelled  $a_v$  and one labelled  $a'_v$ . The rest of the construction is as before. Let  $(K, \mathcal{L}_K)$  be the left Krieger cover of the corresponding sofic shift. As before, the proper communication graph of  $K$  is given by  $E$ , and now  $(K, \mathcal{L}_K)$  satisfies Condition (K), so there is a bijective correspondence between the hereditary subsets of  $E^0$  and the ideals of  $C^*(K) \cong \mathcal{O}_{X^+}$ . Since  $E$  was arbitrary, any finite ideal lattice with a unique maximal ideal can be obtained in this way.  $\square$

**THE  $C^*$ -ALGEBRAS  $\mathcal{O}_{X^+}$  AND  $\mathcal{O}_{X^-}$ .** Every two-sided shift space  $X$  corresponds to two one-sided shift spaces  $X^+$  and  $X^-$ , and this gives two natural ways to associate a universal  $C^*$ -algebra to  $X$ . The next goal is to show that these two  $C^*$ -algebras may carry different information about the shift space. Let  $\mathcal{O}_{X^-}$  be the universal  $C^*$ -algebra associated to the one-sided shift space  $(X^T)^+$  as defined in [9]. The left Krieger cover of  $X^T$  is the transpose of the right Krieger cover of  $X$ , so by [8],  $\mathcal{O}_{X^-}$  is isomorphic to the Cuntz-Krieger algebra of the transpose of the right Krieger cover of  $X$ .

**EXAMPLE 6.2.** Let  $X$  be the sofic shift from Example 5.3. Note that the left and right Krieger covers of  $X$  both satisfy Condition (K) from [36], so the corresponding proper communication graphs completely determine the ideal lattices of  $\mathcal{O}_{X^+}$  and  $\mathcal{O}_{X^-}$ . The proper communication graph of the left Krieger cover  $(K, \mathcal{L}_K)$  of  $X$  is trivial, so  $\mathcal{O}_{X^+}$  is simple, while there are precisely two

vertices in the proper communication graph of the right Krieger cover of  $X$ , so there is exactly one non-trivial ideal in  $\mathcal{O}_{X^-}$ . In particular,  $\mathcal{O}_{X^+}$  and  $\mathcal{O}_{X^-}$  are not isomorphic.

Consider the edge shift  $Y = X_K$ . This is an SFT, and the left and right Krieger covers of  $Y$  are both  $(K, \mathcal{L}_{\text{Id}})$ , where  $\mathcal{L}_{\text{Id}}$  is the identity map on the edge set  $K^1$ . By [8],  $\mathcal{O}_{X^+}$  and  $\mathcal{O}_{Y^+}$  are isomorphic to  $C^*(K)$ . Similarly,  $\mathcal{O}_{Y^-}$  is isomorphic to  $C^*(K^T)$  and  $K^T$  is an irreducible graph satisfying Condition (K), so  $\mathcal{O}_{Y^-}$  is simple. In particular,  $\mathcal{O}_{Y^-}$  is not isomorphic to  $\mathcal{O}_{X^-}$ . This shows that the  $C^*$ -algebras associated to  $X^+$  and  $X^-$  are not always isomorphic, and that there can exist a shift space  $Y$  such that  $\mathcal{O}_{Y^+}$  is isomorphic to  $\mathcal{O}_{X^+}$  while  $\mathcal{O}_{Y^-}$  is not isomorphic to  $\mathcal{O}_{X^-}$ .

AN INVESTIGATION OF CONDITION (\*). In [12], two  $C^*$ -algebras  $\mathcal{O}_X$  and  $\mathcal{O}_{X^*}$  are associated to every two-sided shift space  $X$ . The  $C^*$ -algebras  $\mathcal{O}_X$ ,  $\mathcal{O}_{X^*}$ , and  $\mathcal{O}_{X^+}$  are generated by partial isometries satisfying the same relations, but  $\mathcal{O}_{X^+}$  is always universal unlike  $\mathcal{O}_X$  [9]. In [12], it is proved that  $\mathcal{O}_X$  and  $\mathcal{O}_{X^*}$  are isomorphic when  $X$  satisfies a condition called Condition (\*). The example from [12, Section 4] shows that not all sofic shift spaces satisfy this condition by constructing a sofic shift where the left Krieger cover and the past set cover are not isomorphic. The relationship between Condition (\*) and the structure of the left Krieger cover and the past set cover is further clarified by the final main result. For each  $l \in \mathbb{N}$  and  $w \in \mathcal{B}(X)$  define  $P_l(w) = \{v \in \mathcal{B}(X) \mid vw \in \mathcal{B}(X), |v| \leq l\}$ . Two words  $v, w \in \mathcal{B}(X)$  are said to be  $l$ -past equivalent if  $P_l(v) = P_l(w)$ . For  $x^+ \in X^+$ ,  $P_l(x^+)$  and  $l$ -past equivalence are defined analogously.

CONDITION (\*). For every  $l \in \mathbb{N}$  and every infinite  $F \subseteq \mathcal{B}(X)$  such that  $P_l(u) = P_l(v)$  for all  $u, v \in F$  there exists  $x^+ \in X^+$  such that  $P_l(w) = P_l(x^+)$  for all  $w \in F$ .

LEMMA 6.3. *A vertex  $P$  in the past set cover of a sofic shift  $X$  is in an essential subgraph if and only if there exist infinitely many  $w \in \mathcal{B}(X)$  such that  $P_\infty(w) = P$ .*

*Proof.* Let  $P$  be a vertex in an essential subgraph of the past set cover of  $X$ , and let  $x^+ \in X^+$  be a right ray with a presentation starting at  $P$ . Given  $n \in \mathbb{N}$ , there exists  $w_n \in \mathcal{B}(X)$  such that  $P = P_\infty(x_1 x_2 \dots x_n w_n)$ . To prove the converse, let  $P$  be a vertex in the past set cover for which there exist infinitely many  $w \in \mathcal{B}(X)$  such that  $P = P_\infty(w)$ . For each  $w$ , there is a path labelled  $w_{[1, |w|-1]}$  starting at  $P$ . There are no sources in the past set cover, so this implies that  $P$  is not stranded.  $\square$

PROPOSITION 6.4. *A sofic shift  $X$  satisfies Condition (\*) if and only if the left Krieger cover is the maximal essential subgraph of the past set cover.*

*Proof.* Assume that  $X$  satisfies Condition (\*). Let  $P$  be a vertex in an essential subgraph of the past set cover and define  $F = \{w \in \mathcal{B}(X) \mid P_\infty(w) = P\}$ . Choose  $m \in \mathbb{N}$  such that for all  $x, y \in \mathcal{B}(X) \cup X^+$ ,  $P_\infty(x) = P_\infty(y)$  if and only

if  $P_m(x) = P_m(y)$ . By Lemma 6.3,  $F$  is an infinite set, so Condition (\*) can be used to choose  $x^+ \in X^+$  such that  $P_m(x^+) = P_m(w)$  for all  $w \in F$ . By the choice of  $m$ , this means that  $P_\infty(x^+) = P_\infty(w) = P$  for all  $w \in F$ , so  $P$  is a vertex in the left Krieger cover.

To prove the other implication, assume that the left Krieger cover is the maximal essential subgraph of the past set cover. Let  $l \in \mathbb{N}$  be given, and consider an infinite set  $F \subseteq \mathcal{B}(X)$  for which  $P_l(u) = P_l(v)$  for all  $u, v \in F$ . Since  $X$  is sofic, there are only finitely many different predecessor sets, so there must exist  $w \in F$  such that  $P_\infty(w) = P_\infty(v)$  for infinitely many  $v \in F$ . By Lemma 6.3, this proves that  $P = P_\infty(w)$  is a vertex in the maximal essential subgraph of the past set cover. By assumption, this means that it is a vertex in the left Krieger cover, so there exists  $x^+ \in X^+$  such that  $P_\infty(w) = P_\infty(x^+)$ . In particular,  $P_l(x^+) = P_l(w) = P_l(v)$  for all  $v \in F$ , so Condition (\*) is satisfied.  $\square$

In [2], it was proved that  $\mathcal{O}_{X^*}$  is isomorphic to the Cuntz-Krieger algebra of the past set cover of  $X$  when  $X$  satisfies a condition called Condition (I). According to Carlsen [7], a proof similar to the proof which shows that  $\mathcal{O}_{X^+}$  is isomorphic to the Cuntz-Krieger algebra of the left Krieger cover of  $X$  should prove that  $\mathcal{O}_{X^*}$  is isomorphic to the Cuntz-Krieger algebra of the subgraph of the past set cover of  $X$  induced by the vertices  $P$  for which there exist infinitely many words  $w$  such that  $P_\infty(w) = P$ . Using Lemma 6.3, this shows that  $\mathcal{O}_{X^*}$  is always isomorphic to the Cuntz-Krieger algebra of the maximal essential subgraph of the past set cover of  $X$ .

## REFERENCES

- [1] T. Bates, S. Eilers, and D. Pask. Reducibility of covers of AFT shifts. *Israel J. Math.*, 2011 (in press).
- [2] T. Bates and D. Pask.  $C^*$ -algebras of labelled graphs. *J. Operator Theory*, 57(1):207–226, 2007.
- [3] T. Bates, D. Pask, I. Raeburn, and W. Szymański. The  $C^*$ -algebras of row-finite graphs. *New York J. Math.*, 6:307–324 (electronic), 2000.
- [4] M. Boyle. Flow equivalence of shifts of finite type via positive factorizations. *Pacific J. Math.*, 204(2):273–317, 2002.
- [5] M. Boyle, T. M. Carlsen, and S. Eilers. Flow equivalence of sofic shifts. *In preparation*.
- [6] M. Boyle and D. Huang. Poset block equivalence of integral matrices. *Trans. Amer. Math. Soc.*, 355(10):3861–3886 (electronic), 2003.
- [7] T. M. Carlsen. Personal communication.
- [8] T. M. Carlsen. On  $C^*$ -algebras associated with sofic shifts. *J. Operator Theory*, 49(1):203–212, 2003.

- [9] T. M. Carlsen. Cuntz-Pimsner  $C^*$ -algebras associated with subshifts. *Internat. J. Math.*, 19(1):47–70, 2008.
- [10] T. M. Carlsen and S. Eilers. Augmenting dimension group invariants for substitution dynamics. *Ergodic Theory Dynam. Systems*, 24(4):1015–1039, 2004.
- [11] T. M. Carlsen and S. Eilers. Matsumoto  $K$ -groups associated to certain shift spaces. *Doc. Math.*, 9:639–671 (electronic), 2004.
- [12] T. M. Carlsen and K. Matsumoto. Some remarks on the  $C^*$ -algebras associated with subshifts. *Math. Scand.*, 95(1):145–160, 2004.
- [13] J. Cuntz and W. Krieger. A class of  $C^*$ -algebras and topological Markov chains. *Invent. Math.*, 56(3):251–268, 1980.
- [14] R. Fischer. Sofic systems and graphs. *Monatsh. Math.*, 80(3):179–186, 1975.
- [15] D. Huang. Flow equivalence of reducible shifts of finite type. *Ergodic Theory Dynam. Systems*, 14(4):695–720, 1994.
- [16] N. Jonoska. Sofic shifts with synchronizing presentations. *Theoret. Comput. Sci.*, 158(1-2):81–115, 1996.
- [17] N. Jonoska and B. Marcus. Minimal presentations for irreducible sofic shifts. *IEEE Trans. Inform. Theory*, 40(6):1818–1825, 1994.
- [18] Y. Katayama, K. Matsumoto, and Y. Watatani. Simple  $C^*$ -algebras arising from  $\beta$ -expansion of real numbers. *Ergodic Theory Dynam. Systems*, 18(4):937–962, 1998.
- [19] W. Krieger. On sofic systems. I. *Israel J. Math.*, 48(4):305–330, 1984.
- [20] A. Kumjian, D. Pask, I. Raeburn, and J. Renault. Graphs, groupoids, and Cuntz-Krieger algebras. *J. Funct. Anal.*, 144(2):505–541, 1997.
- [21] D. Lind and B. Marcus. *An introduction to symbolic dynamics and coding*. Cambridge University Press, Cambridge, 1995.
- [22] B. Marcus. Sofic systems and encoding data. *IEEE Trans. Inform. Theory*, 31(3):366–377, 1985.
- [23] K. Matsumoto. On  $C^*$ -algebras associated with subshifts. *Internat. J. Math.*, 8(3):357–374, 1997.
- [24] K. Matsumoto.  $K$ -theory for  $C^*$ -algebras associated with subshifts. *Math. Scand.*, 82(2):237–255, 1998.
- [25] K. Matsumoto. Dimension groups for subshifts and simplicity of the associated  $C^*$ -algebras. *J. Math. Soc. Japan*, 51(3):679–698, 1999.

- [26] K. Matsumoto. Relations among generators of  $C^*$ -algebras associated with subshifts. *Internat. J. Math.*, 10(3):385–405, 1999.
- [27] K. Matsumoto. A simple  $C^*$ -algebra arising from a certain subshift. *J. Operator Theory*, 42(2):351–370, 1999.
- [28] K. Matsumoto. On automorphisms of  $C^*$ -algebras associated with subshifts. *J. Operator Theory*, 44(1):91–112, 2000.
- [29] K. Matsumoto. Stabilized  $C^*$ -algebras constructed from symbolic dynamical systems. *Ergodic Theory Dynam. Systems*, 20(3):821–841, 2000.
- [30] K. Matsumoto. Bowen-Franks groups for subshifts and Ext-groups for  $C^*$ -algebras. *K-Theory*, 23(1):67–104, 2001.
- [31] K. Matsumoto.  $C^*$ -algebras associated with presentations of subshifts. *Doc. Math.*, 7:1–30 (electronic), 2002.
- [32] K. Matsumoto, Y. Watatani, and M. Yoshida. KMS states for gauge actions on  $C^*$ -algebras associated with subshifts. *Math. Z.*, 228(3):489–509, 1998.
- [33] M. Nasu. An invariant for bounded-to-one factor maps between transitive sofic subshifts. *Ergodic Theory Dynam. Systems*, 5(1):89–105, 1985.
- [34] M. Nasu. Topological conjugacy for sofic systems. *Ergodic Theory Dynam. Systems*, 6(2):265–280, 1986.
- [35] B. Parry and D. Sullivan. A topological invariant of flows on 1-dimensional spaces. *Topology*, 14(4):297–299, 1975.
- [36] I. Raeburn. *Graph algebras*, volume 103 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC, 2005.
- [37] K. Thomsen. On the structure of a sofic shift space. *Trans. Amer. Math. Soc.*, 356(9):3557–3619 (electronic), 2004.
- [38] B. Weiss. Subshifts of finite type and sofic systems. *Monatsh. Math.*, 77:462–474, 1973.

Rune Johansen  
Department of  
Mathematical Sciences  
University of Copenhagen  
Universitetsparken 5  
2100 København Ø  
Denmark

