

SOME ENDOSCOPIC PROPERTIES OF THE
ESSENTIALLY TAME JACQUET-LANGLANDS
CORRESPONDENCE

KAM-FAI TAM

Received: September 28, 2015

Communicated by Takeshi Saito

ABSTRACT. Let F be a non-Archimedean local field of characteristic 0 and G be an inner form of the general linear group $G^* = \mathrm{GL}_n$ over F . We show that the rectifying character appearing in the essentially tame Jacquet-Langlands correspondence of Bushnell and Henniart for G and G^* can be factorized into a product of some special characters, called zeta-data in this paper, in the theory of endoscopy of Langlands and Shelstad. As a consequence, the essentially tame local Langlands correspondence for G can be described using admissible embeddings of L-tori.

2010 Mathematics Subject Classification: Primary 22E50; Secondary 11S37, 11F70.

Keywords and Phrases: essentially tame Jacquet-Langlands correspondence, inner forms, admissible pairs, zeta-data, endoscopy, admissible embeddings

CONTENTS

1	INTRODUCTION	346
1.1	Background	347
1.2	Main results	348
1.3	Relation with the previous results	350
1.4	Notations	352

2	SOME BASIC SETUPS	352
2.1	Root system	352
2.2	Galois groups	354
2.3	Division algebra	355
2.4	Hereditary orders in central simple algebra	356
2.5	Embedding conditions	357
2.6	Characters	358
3	THE ESSENTIALLY TAME JACQUET-LANGLANDS CORRESPONDENCE	358
3.1	Admissible characters	358
3.2	The correspondences	359
3.3	Some subgroups	361
3.4	Simple characters	362
3.5	Local constructions of attached pairs	363
3.6	Finite symplectic modules	364
3.7	Invariants of finite symplectic modules	366
3.8	Values of rectifiers	367
4	FINITE SYMPLECTIC MODULES	368
4.1	Standard modules of central simple algebra	368
4.2	Complete decomposition of finite symplectic modules	371
4.3	Some properties of parities of jumps	373
4.4	Symmetric submodules	373
4.4.1	Case when f is odd	373
4.4.2	Case when f is even	374
4.4.3	Case when both f and m are even	374
4.4.4	Case when f is even and m is odd	374
4.5	t -factors of isotypic components	375
5	ZETA-DATA	377
5.1	Admissible embeddings of L-tori	377
5.2	Symmetric unramified zeta-data	379
5.3	Zeta-data associated to admissible characters	380
5.3.1	Case when f is odd	381
5.3.2	Case when f is even	382
5.4	The main theorem	383
5.5	Functorial property	384

1 INTRODUCTION

Let G^* be a general linear group over a non-Archimedean local field of characteristic 0, and G be an inner form of G^* . In this paper, we refine the results of the rectifying characters in the context of the essentially tame Jacquet-Langlands correspondence [BH11] by proving that each rectifying character

admits a factorization into a product of characters called ζ -data, defined similarly to χ -data in [LS87], which are significant in describing the essentially tame local Langlands correspondence for G .

We know from [BH11] that the rectifying characters are quadratic characters that measure the difference between two correspondences for essentially tame supercuspidal representations of G and G^* : the representation theoretic one by matching the maximal simple types of the two groups, and the functorial one from the Jacquet-Langlands correspondence. On the representation theoretic side, the maximal simple types of G can be constructed using certain characters of its elliptic maximal tori, while on the functorial side, the Langlands parameters for G can be functorially lifted from the parameters of the same collection of characters.

Combining these results with our result on rectifying characters, we show that the essentially tame local Langlands correspondence for G can be described completely by admissible embeddings, defined in [LS87], of the L-groups of elliptic maximal tori into the L-group of G , generalizing an analogous description proved by the author [Tam] in the split case (when $G = G^*$).

1.1 BACKGROUND

Let F be a non-Archimedean local field, G^* be the group GL_n defined over F , and G be an inner form of G^* defined over F . The set of F -points $G(F)$ of G is therefore isomorphic to $\mathrm{GL}_m(D)$ as a group, where D is a central division algebra over F of dimension d^2 and $m = n/d$.

Let $\mathcal{A}_m^2(D)$ (resp. $\mathcal{A}_n^2(F)$) be the discrete series of $G(F)$ (resp. $G^*(F)$), i.e., the set of equivalence classes of irreducible admissible representations that are essentially square integrable mod-center. The Jacquet-Langlands correspondence asserts a bijection

$$JL : \mathcal{A}_n^2(F) \rightarrow \mathcal{A}_m^2(D)$$

determined by a character relation (see (3.5)) between a representation in $\mathcal{A}_n^2(F)$ and its image in $\mathcal{A}_m^2(D)$. The existence of this bijection is known, starting from the case $n = 2$ [JL70], when $G(F)$ is the multiplicative group of the quaternion algebra over F . For arbitrary n , when $G(F)$ is the multiplicative group of a division algebra, the existence is proved by [Rog83]. The general situations are treated by [DKV84] in the characteristic zero case and by [Bad02], [BHL10] in the positive characteristic case.

Bushnell and Henniart describe in [BH11, (2.1)] the image of JL when it is restricted to the subset $\mathcal{A}_n^0(F)$ of supercuspidal representations. The image is the subset of representations in $\mathcal{A}_m^2(D)$, each of whose *parametric degree* is equal to n . We do not need the full definition of the parametric degree of a representation, so we only refer to [BH11, Section 2.7] for details. We only need to know that

- the parametric degree of a representation in $\mathcal{A}_m^2(D)$ is a positive integer divisor of n ,

- a representation in $\mathcal{A}_m^2(D)$ is supercuspidal if its parametric degree is n ; the converse is true in the split case (when $G = G^*$) but not in general, and
- the parametric degree is preserved under JL .

Furthermore, we can describe the image of JL of *each* representation $\pi \in \mathcal{A}_n^0(F)$ when π is *essentially tame*, a notion we will explain in Section 3.2. More precisely, if we let $\mathcal{A}_m^{\text{et}}(D)$ (resp. $\mathcal{A}_n^{\text{et}}(F)$) be the set of essentially tame representations in $\mathcal{A}_m^2(D)$ (resp. $\mathcal{A}_n^2(F)$) whose parametric degree is n , then we can describe completely the *essentially tame Jacquet-Langlands correspondence*:

$$JL : \mathcal{A}_n^{\text{et}}(F) \rightarrow \mathcal{A}_m^{\text{et}}(D),$$

as in [BH11, Theorem A].

To explain the theorem and describe JL completely, we require the notion of admissible characters from [How77]. In Section 3.1, we define the set $P_n(F)$ of (equivalence classes of) admissible pairs $(E/F, \xi) \in P_n(F)$, where E/F is a tamely ramified extension of degree n and ξ is a character of E^\times admissible over F . This set bijectively parametrizes both $\mathcal{A}_n^{\text{et}}(F)$ and $\mathcal{A}_m^{\text{et}}(D)$ explicitly [BH11], using the theory of simple types of $G(F)$ developed in [BF85], [Gra07], [Séc04], [Séc05a], [Séc05b], [SS08], [BSS12] which generalizes the corresponding theory in the split case [BK93], [BH96] and the division algebra case [Zin92], [Bro96].

If we denote by

$${}_F\Pi : P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F), (E/F, \xi) \mapsto {}_F\Pi_\xi \quad (1.1)$$

and

$${}_D\Pi : P_n(F) \rightarrow \mathcal{A}_m^{\text{et}}(D), (E/F, \xi) \mapsto {}_D\Pi_\xi \quad (1.2)$$

the above bijections, then Bushnell and Henniart proved in [BH11] that the composition

$$\nu : P_n(F) \xrightarrow{{}_F\Pi} \mathcal{A}_n^{\text{et}}(F) \xrightarrow{JL} \mathcal{A}_m^{\text{et}}(D) \xrightarrow{{}_D\Pi^{-1}} P_n(F)$$

maps an admissible pair $(E/F, \xi) \in P_n(F)$ to another pair of the form $(E/F, \xi \cdot {}_D\nu_\xi)$, where ${}_D\nu_\xi$ is a tamely ramified character ${}_D\nu_\xi$ of E^\times depending on ξ . We borrow the terminology from [BH10] and call the character ${}_D\nu_\xi$ the *rectifier* of ξ for the essentially tame Jacquet-Langlands correspondence.

1.2 MAIN RESULTS

The main result of this paper is to relate the rectifier ${}_D\nu_\xi$ with a special set of characters, called ζ -data in this paper, introduced in the theory of endoscopy of Langlands and Shelstad [LS87]. The significance of ζ -data will be explained in the next section, together with a brief summary of the previous results of the author [Tam].

To describe the main result, we first assume that $\text{char}(F) = 0$ (see Remark 1.3 about this assumption). Let T be the F -torus such that $T(F) = E^\times$. We regard T as a maximal torus embedded in G . In contrast to the split case, we have to carefully choose the embedding $T \rightarrow G$ relative to an hereditary \mathfrak{o}_F -order in $G(F)$ associated to ξ . This will be explained in Section 2.5. Given this embedding, let $\Phi = \Phi(G, T)$ be the root system, which is invariant under the action of the absolute Galois group Γ_F of F if we view Φ as a subset of the character group of T . For each root $\lambda \in \Phi$, we denote by E_λ the fixed field of the stabilizer of λ in Γ_F , so that E_λ is a field extension of a Γ_F -conjugate of E . We recall from [LS87, Corollary 2.5.B] that ζ -data is a set of characters

$$\{\zeta_\lambda\} = \{\zeta_\lambda\}_{\lambda \in \Gamma_F \backslash \Phi},$$

where each ζ_λ is a character of E_λ^\times satisfying the conditions in *loc. cit.* (and will be recalled in Section 5.1). Here λ ranges over a suitable subset of roots in Φ , denoted by $\Gamma_F \backslash \Phi$ for the moment, representing the Γ_F -orbits of Φ and such that E_λ is a field extension of E (but not just its conjugate).

The following theorem restates the main result, Theorem 5.5, in a simpler way.

THEOREM 1.1. *Given a character ξ of E^\times admissible over F .*

(i) *There exists a set of ζ -data $\{\zeta_{\lambda, \xi}\}_{\lambda \in \Gamma_F \backslash \Phi}$ such that*

$${}_D\nu_\xi = \prod_{\lambda \in \Gamma_F \backslash \Phi} \zeta_{\lambda, \xi}|_{E^\times}.$$

(ii) *The values of each $\zeta_{\lambda, \xi}$ can be expressed in terms of certain invariants, called t -factors in this paper, of the corresponding component in the complete symmetric decomposition of the finite symplectic modules associated to ξ (see the notations and definitions in Sections 4.2 and 3.7).*

□

We explain statement (ii) of the above theorem. The finite symplectic modules appear in the respective constructions of the extended maximal simple types inducing ${}_F\Pi_\xi$ and ${}_D\Pi_\xi$ in (1.1) and (1.2) (see [BH11, (2.5.4)], or the summary in Section 3.5). Each of these modules admits an orthogonal decomposition, called a complete symmetric decomposition in this paper (Proposition 4.4), whose components are parametrized by the same set $\Gamma_F \backslash \Phi$ parameterizing the factors in (i) of the Theorem. The t -factors are, roughly speaking, defined by the symplectic signs attached to these components.

When proving Theorem 1.1, we pick a choice of characters $\{\zeta_{\lambda, \xi}\}_{\lambda \in \Gamma_F \backslash \Phi}$, where each character $\zeta_{\lambda, \xi}$ has values in terms of the t -factors of the corresponding component, the one indexed by λ . We then show that these characters constitute a set of ζ -data. Moreover, using the multiplicativity of t -factors, the product of these ζ -data, when restricted to E^\times , is equal to the rectifier ${}_D\nu_\xi$, whose values

are given in the First and Second Comparison Theorems of [BH11]. Hence our result refines the one in *loc. cit.*

While the finite symplectic modules and their decompositions are also studied by the author in the split case [Tam], there are extra conditions on the components of these modules in the general case. These conditions come from the extra ramifications of the related compact subgroups in constructing the extended maximal simple types. The degrees of these ramifications depend on the residue degree $f(E/F)$ and, with other conditions similar to those in the split case, determine whether each component is trivial or not. This new phenomenon will be fully studied in Section 4. In particular, when E/F is totally ramified and ξ is fixed, the finite symplectic modules are isomorphic to each other for all inner forms of G^* , a fact already known in [BH11, Proposition 5.6].

1.3 RELATION WITH THE PREVIOUS RESULTS

The significance of the factorization of ${}_{D\nu\xi}$ in Theorem 1.1(i) comes from [Tam], which proves an analogous factorization of the rectifier ${}_{F\mu\xi}$ for the essentially tame local Langlands correspondence [BH05a].

We first recall from *loc. cit.* that the rectifier ${}_{F\mu\xi}$ measures the difference between the “naïve correspondence” and the essentially tame Langlands correspondence for G^* ; more precisely, the Langlands parameter of ${}_{F\Pi\xi}$ defined in (1.1) is the induced representation

$$\text{Ind}_{\mathcal{W}_E}^{\mathcal{W}_F}(\xi \cdot {}_{F\mu\xi}) \tag{1.3}$$

of the Weil group \mathcal{W}_F of F , where $\xi \cdot {}_{F\mu\xi}$ is regarded as a character of \mathcal{W}_E by class field theory [Tat79]. In [Tam, Theorem 1.1], the author proved that the rectifier ${}_{F\mu\xi}$ admits a factorization

$${}_{F\mu\xi} = \prod_{\lambda \in \Gamma_F \setminus \Phi} \chi_{\lambda, \xi}|_{E^\times},$$

where $\{\chi_{\lambda, \xi}\}_{\lambda \in \Gamma_F \setminus \Phi}$ is a set of χ -data, consisting of characters of E_λ^\times satisfying the conditions similar to those of ζ -data (see Section 5.1).

With a collection of χ -data, we follow [LS87, Section 2.6] to construct an admissible embedding

$$I_{\{\chi_{\lambda, \xi}\}} : {}^L T \rightarrow {}^L G$$

of the L-group ${}^L T$ of the maximal torus T into the L-group ${}^L G$ of G^* . (For convenience, we call ${}^L T$ an L-torus in this paper.) Let $\tilde{\xi} : \mathcal{W}_F \rightarrow {}^L T$ be an L-homomorphism whose class is the parameter of the character ξ of $E^\times = T(F)$ under the local Langlands correspondence of the torus T , i.e., the Artin reciprocity for E^\times [Tat79]. In a previous result of the author [Tam, Corollary 1.2], the Langlands parameter (1.3) of ${}_{F\Pi\xi}$ is isomorphic to the composition

$$I_{\{\chi_{\lambda, \xi}\}} \circ \tilde{\xi} : \mathcal{W}_F \rightarrow {}^L T \rightarrow {}^L G \xrightarrow{\text{natural proj.}} \text{GL}_n(\mathbb{C})$$

as a representation of \mathcal{W}_F . In other words, the essentially tame local Langlands correspondence for G^* can be described by admissible embeddings of L-tori. A set of ζ -data is the ‘difference’ of two sets of χ -data, in the sense that, given a set $\{\chi_\lambda\}$ of χ -data, we have

$$\{\chi'_\lambda\} \text{ is another set of } \chi\text{-data} \iff \{\chi_\lambda(\chi'_\lambda)^{-1}\} \text{ is a set of } \zeta\text{-data.}$$

If we define the local Langlands correspondence for G as the composition of the local Langlands correspondence for G^* and the Jacquet-Langlands correspondence JL , then we can express the Langlands parameter of ${}_D\Pi_\xi$ using an admissible embedding of L-tori, as follows.

COROLLARY 1.2. *Let $\{\chi_{\lambda,\xi}\}$ and $\{\zeta_{\lambda,\xi}\}$ be respectively the χ -data and the ζ -data associated to an admissible character ξ . The Langlands parameter*

$$\text{Ind}_{\mathcal{W}_E}^{\mathcal{W}_F}(\xi \cdot {}_F\mu_\xi \cdot {}_D\nu_\xi)$$

of ${}_D\Pi_\xi$ is isomorphic to

$$I_{\{\chi_{\lambda,\xi}, \zeta_{\lambda,\xi}\}} \circ \tilde{\xi} : \mathcal{W}_F \rightarrow {}^L T \rightarrow {}^L G \xrightarrow{\text{natural proj.}} \text{GL}_n(\mathbb{C})$$

as a representation of \mathcal{W}_F . □

Hence analogously we can describe the essentially tame local Langlands correspondence for G by admissible embeddings of L-tori.

As a consequence, we show in Proposition 5.7 that the factorization of ${}_D\nu_\xi$ in Theorem 1.1(i) is functorial, in the following sense. Let K/F be an intermediate extension of E/F , so that if the pair $(E/F, \xi)$ is admissible over F , then $(E/K, \xi)$ is admissible over K by definition. We denote the centralizer of K^\times in $G(F)$ by $GL_{m_K}(D_K)$, where D_K is a K -division algebra and m_K a positive integer. If $\{\zeta_{\lambda,\xi}\}$ is the set of ζ -data associated to ξ , then the partial product

$$\prod_{\lambda \in \Gamma_F \setminus \Phi, \lambda|_{K^\times} \neq 1} \zeta_{\lambda,\xi}|_{E^\times}.$$

(a product similar to Theorem 1.1(i), with factors ranging over the characters being non-trivial on K^\times) is the rectifier ${}_{D_K}\nu_\xi$ of ξ over K .

REMARK 1.3. We would like to remark on the condition of the characteristic $\text{char}(F) = 0$, as we also did in [Tam, Remark 1.3]. The readers should be aware that the works of [JL70], [DKV84], [Bad02], [BH11] make the Jacquet-Langlands correspondence valid for local fields of arbitrary characteristic. In our paper, we apply the condition $\text{char}(F) = 0$ only because we refer to the theory of endoscopy from [LS87], [KS99]. However, we do not actually need this condition for the part of the theory that we allude to, which is about the admissible embeddings of L-tori. In [Tam, Section 6] (or rather [LS87, Section 2.5]), we see that these kind of embeddings can be defined without any condition of $\text{char}(F)$. Therefore, the condition $\text{char}(F) = 0$ in this paper should be treated as a mild condition. □

1.4 NOTATIONS

Throughout the paper, F denotes a non-Archimedean local field of characteristic 0. Its ring of integers is \mathfrak{o}_F with the maximal ideal \mathfrak{p}_F . The residue field $\mathbf{k}_F = \mathfrak{o}_F/\mathfrak{p}_F$ has q elements and is of characteristic p . We denote by $v_F : F^\times \rightarrow \mathbb{Z}$ the discrete valuation on F . We denote by Γ_F the absolute Galois group of F , and by \mathcal{W}_F the Weil group of F .

The multiplicative group F^\times decomposes into a product of subgroups

$$\langle \varpi_F \rangle \times \boldsymbol{\mu}_F \times U_F^1.$$

They are namely the group generated by a prime element ϖ_F , the group $\boldsymbol{\mu}_F$ of roots of unity of order prime to p , and the 1-unit group $U_F^1 := 1 + \mathfrak{p}_F$. We will identify $\boldsymbol{\mu}_F$ with \mathbf{k}_F^\times in the natural way. We then write $U_F = U_F^0 := \boldsymbol{\mu}_F \times U_F^1$ and $U_F^i := 1 + \mathfrak{p}_F^i$ for each positive integer i . Let $\boldsymbol{\mu}_n$ be the group of n th roots of unity in the algebraic closure \bar{F} of F , and z_n be a choice of primitive n th root in $\boldsymbol{\mu}_n$.

The F -level of a character ξ of F^\times is the smallest integer $a \geq -1$ such that $\xi|_{U_F^{a+1}}$ is trivial. A character ξ of F^\times is called unramified if $\xi|_{U_F}$ is trivial, or equivalently, if its F -level is -1 . It is called tamely ramified if $\xi|_{U_F^1}$ is trivial, or equivalently, if its F -level is 0.

Given a field extension E/F , we denote its ramification index by $e = e(E/F)$ and its residue degree by $f = f(E/F)$. We also denote by $\text{tr}_{E/F}$ and norm $N_{E/F}$ the trace and norm respectively.

We fix an additive character ψ_F of F of level 0, which means that ψ_F is trivial on \mathfrak{p}_F but is non-trivial on \mathfrak{o}_F . Hence $\psi_F|_{\mathfrak{o}_F}$ induces a non-trivial character of \mathbf{k}_F . We write $\psi_E = \psi_F \circ \text{tr}_{E/F}$.

Suppose that A is a central simple algebra over F . We denote the reduced trace by $\text{trd}_{A/F}$ and the reduced norm by $\text{Nrd}_{A/F}$.

Given a set X , we denote its cardinality by $\#X$. If H is a group and X is a H -set, then we denote the action of $h \in H$ on $x \in X$ by $x \mapsto {}^h x$. The set of H -orbits is denoted by $H \backslash X$. If π is a representation of H (over a given field), we denote its equivalence class by (H, π) .

2 SOME BASIC SETUPS

2.1 ROOT SYSTEM

Given a field extension E/F of degree n , we let T be the induced torus $\text{Res}_{E/F} \mathbb{G}_m$. We embed T into G as an elliptic maximal torus, and denote the image still by T . The choice of this embedding will be specific in Section 2.5, but at this moment this choice is irrelevant. Let $\Phi = \Phi(G, T)$ be the root system of T in G . Following [Tam, Section 3.1], we can denote each root in Φ by $[\frac{g}{h}]$ where $g = g\Gamma_E$ and $h = h\Gamma_E$ are distinct cosets in Γ_F/Γ_E . (We use the same notation g for an element in Γ_F and its Γ_E -cosets, for notation convenience.) The Γ_F -action on Φ is given by $x \cdot [\frac{g}{h}] = [\frac{xg}{xh}]$. For each root

$\lambda \in \Phi$, we denote by $[\lambda]$ its Γ_F -orbit $\mathcal{W}_F\lambda$. Each Γ_F -orbit contains a root of the form $\left[\frac{1}{g}\right]$ for some non-trivial coset $g \in \Gamma_F/\Gamma_E$.

For each root $\lambda \in \Phi$, we denote the stabilizers $\{g \in \Gamma_F \mid {}^g\lambda = \lambda\}$ and $\{g \in \Gamma_F \mid {}^g\lambda = \pm\lambda\}$ by Γ_λ and $\Gamma_{\pm\lambda}$ respectively and their fixed fields by E_λ and $E_{\pm\lambda}$ respectively. We call a root λ *symmetric* if $[E_\lambda : E_{\pm\lambda}] = 2$, and *asymmetric* otherwise. Equivalently, λ is symmetric if and only if λ and $-\lambda$ are in the same Γ_F -orbit. Note that the symmetry of Φ is preserved by the Γ_F -action. Let

- (i) $\Gamma_F \backslash \Phi_{\text{sym}}$ be the set of Γ_F -orbits of symmetric roots,
- (ii) $\Gamma_F \backslash \Phi_{\text{asym}}$ be the set of Γ_F -orbits of asymmetric roots, and
- (iii) $\Gamma_F \backslash \Phi_{\text{asym}/\pm}$ be the set of equivalence classes of asymmetric Γ_F -orbits by identifying $[\lambda]$ and $[-\lambda]$.

We denote by $(\Gamma_E \backslash \Gamma_F / \Gamma_E)'$ the collection of non-trivial double cosets, and by $[g]$ the double coset $\Gamma_E g \Gamma_E$. We can deduce the following proposition easily.

PROPOSITION 2.1. *The map*

$$\Gamma_F \backslash \Phi \rightarrow (\Gamma_E \backslash \Gamma_F / \Gamma_E)', [\lambda] = \mathcal{W}_F \left[\frac{1}{g}\right] \mapsto [g],$$

is a bijection between the set $\Gamma_F \backslash \Phi$ of Γ_F -orbits of the root system Φ and the set $(\Gamma_E \backslash \Gamma_F / \Gamma_E)'$ of non-trivial double cosets.

We can therefore call $g \in \Gamma_F$ *symmetric* if $[g] = [g^{-1}]$, and *asymmetric* otherwise, so that the bijection in Proposition 2.1 preserves symmetries on both sides. Let

- (i) $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}}$ be the set of symmetric non-trivial double cosets,
- (ii) $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}}$ be the set of asymmetric non-trivial double cosets, and
- (iii) $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}/\pm}$ be the set of equivalence classes of $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}}$ by identifying $[g]$ with $[g^{-1}]$.

We choose subsets \mathcal{D}_{sym} and $\mathcal{D}_{\text{asym}/\pm}$ of representatives in Γ_F/Γ_E of $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}}$ and $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}/\pm}$ respectively, and write

$$\mathcal{D}_{\text{asym}} = \mathcal{D}_{\text{asym}/\pm} \sqcup \{g^{-1} \mid g \in \mathcal{D}_{\text{asym}/\pm}\}.$$

We also choose subsets \mathcal{R}_{sym} and $\mathcal{R}_{\text{asym}/\pm}$ of representatives in Φ of orbits in $\Gamma_F \backslash \Phi_{\text{sym}}$ and $\Gamma_F \backslash \Phi_{\text{asym}/\pm}$ respectively such that every root $\lambda \in \mathcal{R}_\pm := \mathcal{R}_{\text{sym}} \sqcup \mathcal{R}_{\text{asym}/\pm}$ is of the form $\left[\frac{1}{g}\right]$ for some $g \in \mathcal{D}_\pm := \mathcal{D}_{\text{sym}} \sqcup \mathcal{D}_{\text{asym}/\pm}$, and write

$$\mathcal{R}_{\text{asym}} = \mathcal{R}_{\text{asym}/\pm} \sqcup (-\mathcal{R}_{\text{asym}/\pm}).$$

Hence \mathcal{R}_{sym} , $\mathcal{R}_{\text{asym}}$, and $\mathcal{R}_{\text{asym}/\pm}$ correspond bijectively to \mathcal{D}_{sym} , $\mathcal{D}_{\text{asym}}$, and $\mathcal{D}_{\text{asym}/\pm}$ respectively by the identification in Proposition 2.1. Denote $E_g := E_\lambda$ and $E_{\pm g} := E_{\pm\lambda}$. Notice that $E_g = E({}^gE)$, composite field of E and gE .

2.2 GALOIS GROUPS

Let E/F be a field extension of degree n . In most of the paper, we assume that E/F is tamely ramified, which means that p is coprime to e . By [Lan94, II.§5], we can choose ϖ_E and ϖ_F such that

$$\varpi_E^e = z_{E/F}\varpi_F, \text{ for some } z_{E/F} \in \mu_E. \tag{2.1}$$

Choose in \bar{F}^\times a primitive e th root of unity z_e and an e th root $z_{E/F,e}$ of $z_{E/F}$. (We do not require that $z_{E/F,e}^a = z_e$, if a is the multiplicative order of $z_{E/F}$.) Denote $L = E[z_e, z_{E/F,e}]$ and $l = [L : E]$. With the choices of ϖ_F and ϖ_E as in (2.1), we define the following F -automorphisms on L .

- (i) $\phi : z \mapsto z^q$, for all $z \in \mu_L$, and $\phi : \varpi_E \mapsto z_\phi \varpi_E$.
- (ii) $\sigma : z \mapsto z$, for all $z \in \mu_L$, and $\sigma : \varpi_E \mapsto z_e \varpi_E$.

Here z_ϕ lies in μ_E satisfying $(z_\phi \varpi_E)^e = z_{E/F}^q \varpi_F$. More generally, we write $\phi^i \varpi_E = z_{\phi^i} \varpi_E$ where $z_{\phi^i} = z_\phi^{1+q+\dots+q^{i-1}}$ is an e th root of $z_{E/F}^{q^i-1}$. Therefore, $\Gamma_{L/F} = \langle \sigma \rangle \rtimes \langle \phi \rangle$ with relation $\phi \circ \sigma \circ \phi^{-1} = \sigma^q$. Suppose that $\Gamma_{L/E} = \langle \sigma^c \phi^f \rangle$ for some integer c satisfying the condition:

$$e \text{ divides } c \left(\frac{q^{fl} - 1}{q^f - 1} \right).$$

We can choose

$$\{\sigma^i \phi^j \mid i = 0, \dots, e - 1, j = 0, \dots, f - 1\}$$

as coset representatives for the quotient $\Gamma_{E/F} = \Gamma_F/\Gamma_E$. Moreover, elements in a fixed double coset are of the form $[\sigma^i \phi^j]$ with a fixed $j \pmod f$.

PROPOSITION 2.2 ([Tam, Proposition 3.3]). *The double coset $[g] = [\sigma^i \phi^j]$ is symmetric only if $j = 0$ or, when f is even, $j = f/2$. \square*

We call those symmetric $[\sigma^i]$ ramified and those symmetric $[\sigma^i \phi^{f/2}]$ unramified, and denote by $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym-ram}}$ and $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym-unram}}$ respectively the collections of symmetric ramified and symmetric unramified double cosets. We provide several useful results concerning the parity of certain subsets in $\Gamma_E \backslash \Gamma_F / \Gamma_E$.

PROPOSITION 2.3 ([Tam, Propositions 3.4 and 3.5]). *(i) If $[g]$ is symmetric unramified, then the degree $[E_g : E]$ is odd.*

(ii) The parity of $\#(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym-unram}}$ is equal to that of $e(f - 1)$. \square

LEMMA 2.4. *Suppose that f is even. The following are equivalent.*

- (i) *There exists $\sigma^i \phi^{f/2} \in W_{F[\varpi_E]}$ for some i .*

- (ii) $z_{\phi^{f/2}}$ is an e th root of unity.
- (iii) $z_{E/F} \in K_+$, where K_+/F is unramified of degree $f/2$.
- (iv) $f_{\varpi} := [E : F[\varpi_E]]$ is even.

Proof. (i) is equivalent to (ii) since $\sigma^i \phi^{f/2} \varpi_E = z_e^i z_{\phi^{f/2}} \varpi_E$. To show that (iii) implies (ii), we recall that $z_{\phi^{f/2}}$ is an e th root of $z_{E/F}^{q^{f/2}-1}$. If $z_{E/F} \in K_+$, then $z_{E/F}^{q^{f/2}-1} = 1$ and $z_{\phi^{f/2}}$ is an e th root of unity. The converse is similar. To show the equivalence of (iii) and (iv), we notice that $f(F[\varpi_E]/F) = f(F[z_{E/F}]/F) = f/f_{\varpi}$. Hence that $F[z_{E/F}] \subseteq K_+$ is equivalent to that f_{ϖ} is even. \square

LEMMA 2.5. *Suppose that $g = \sigma^i \phi^{f/2}$ satisfies the conditions in Lemma 2.4.*

- (i) The double coset $[\sigma^i \phi^{f/2}]$ is automatically symmetric.
- (ii) The set $(\Gamma_E \backslash \Gamma_{F[\varpi_E]} / \Gamma_E)_{\text{sym-unram}} = (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym-unram}} \cap (\Gamma_E \backslash \Gamma_{F[\varpi_E]} / \Gamma_E)$ contains a single element $[\sigma^i \phi^{f/2}]$.

Proof. For (i), we consider the actions of $\sigma^i \phi^{f/2}$ and its inverse $(\sigma^i \phi^{f/2})^{-1}$ on E . We certainly have $\sigma^i \phi^{f/2} \varpi_E = (\sigma^i \phi^{f/2})^{-1} \varpi_E = \varpi_E$ by definition. We also have $\sigma^i \phi^{f/2} z = (\sigma^i \phi^{f/2})^{-1} z = z^{q^{f/2}}$ for all $z \in \mu_E$. Therefore $\sigma^i \phi^{f/2} \Gamma_E = (\sigma^i \phi^{f/2})^{-1} \Gamma_E$ and in particular $[\sigma^i \phi^{f/2}] = [(\sigma^i \phi^{f/2})^{-1}]$. For (ii), we know by Lemma 2.4.(ii) that the double coset is the one containing $\sigma^i \phi^{f/2}$ where $z_{\phi^{f/2}} = z_e^{-i}$. \square

PROPOSITION 2.6. *When f is even, the parity of the cardinality of $(\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym-unram}} - \Gamma_{F[\varpi_E]}$ is equal to $e + f_{\varpi} - 1$.*

Proof. Recall

- (i) by Lemma 2.4 that there exists $\sigma^i \phi^{f/2} \in W_{F[\varpi_E]}$ if and only if f_{ϖ} is even, and
- (ii) by Proposition 2.3 that the parity of the number of symmetric $[\sigma^i \phi^{f/2}]$ is the same as that of e .

By combining these facts, we have the assertion. \square

2.3 DIVISION ALGEBRA

Let D be a division algebra over F of dimension n^2 . Denote its unique maximal order by \mathfrak{o}_D and the maximal ideal of \mathfrak{o}_D by \mathfrak{p}_D . Suppose that the Hasse-invariant of D is $h = h(D)$, so that $\gcd(n, h) = 1$. By [Rei03, (14.5) Theorem], we can choose a primitive $(q^n - 1)$ th root z of unity in D and a uniformizer ϖ_D such that

$$\varpi_D^n = \varpi_F \text{ and } \varpi_D z \varpi_D^{-1} = z^{q^h}. \tag{2.2}$$

We write $K_n = F[z]$ and $\mu_D = \langle z \rangle$, then K_n is a maximal unramified extension (of degree n) in D , and μ_D is a group of roots of unity of order $q^n - 1$, both defined up to conjugacy by D^\times . Therefore, the conjugation of ϖ_D acts on K_n as the h th power of the Frobenius automorphism, i.e.,

$$\varpi_D u \varpi_D^{-1} = \phi^h u \text{ for all } u \in K_n.$$

We write $U_D^i := 1 + \mathfrak{p}_D^i$ for all positive integer i . The multiplicative subgroup D^\times hence decomposes into a semi-direct product

$$(\langle \varpi_D \rangle \rtimes \mu_D) \rtimes U_D^1.$$

Let E/F be a tamely ramified field extension of degree n , and let K be the maximal unramified sub-extension in E/F . We assume that $K \subseteq K_n$ and that the uniformizers ϖ_E and ϖ_F satisfy $\varpi_E^e = z_{E/F} \varpi_F$ as in (2.1) for some $z_{E/F} \in \mu_E$. If we define $z_{D/E} \in \mu_D = \mu_{K_n}$ to be a solution of

$$N_{K_n/K}(z_{D/E}) = z_{E/F}, \tag{2.3}$$

then we may take $\varpi_E = \varpi_D^f z_{D/E}$ and this defines an embedding of E into D over F . Note that from (2.2)

$$z \varpi_D^i z^{-1} = z^{1-q^{hi}} \varpi_D^i \tag{2.4}$$

for all $z \in \mu_E = \mu_K$ and all $i \in \mathbb{Z}$.

2.4 HEREDITARY ORDERS IN CENTRAL SIMPLE ALGEBRA

If G is an F -inner form of $G^* = \text{GL}_n$, then $G(F) = A^\times$, where A be a central simple algebra over F . By Wedderburn Theorem [Rei03, (7.4)Theorem], A is isomorphic to $\text{Mat}_m(D)$, where D is a division algebra of F -dimension d^2 and $md = n$. Therefore, $G(F) \cong \text{GL}_m(D)$. Any field extension of degree n can be embedded into A as a maximal subfield in A , and any two such embeddings are conjugate under $G(F)$.

Let \mathfrak{A} be an \mathfrak{o}_F -hereditary order in A , $\mathfrak{P}_{\mathfrak{A}}$ be its Jacobson radical, and $\mathfrak{K}_{\mathfrak{A}}$ be the $G(F)$ -normalizer of \mathfrak{A}^\times . If \mathfrak{A} is principal, in the sense that there exists $\varpi_{\mathfrak{A}} \in \mathfrak{K}_{\mathfrak{A}}$ such that $\varpi_{\mathfrak{A}} \mathfrak{A} = \mathfrak{A} \varpi_{\mathfrak{A}} = \mathfrak{P}_{\mathfrak{A}}$, then the valuation $v_{\mathfrak{A}} : \mathfrak{K}_{\mathfrak{A}} \rightarrow \mathbb{Z}$ is defined by $x \mathfrak{A} = \mathfrak{A} x = \mathfrak{P}_{\mathfrak{A}}^{v_{\mathfrak{A}}(x)}$ for all $x \in \mathfrak{K}_{\mathfrak{A}}$. We also write $U_{\mathfrak{A}} = U_{\mathfrak{A}}^0 = \mathfrak{A}^\times$, $U_{\mathfrak{A}}^i = 1 + \mathfrak{P}_{\mathfrak{A}}^i$ for each positive integer i , $U_{\mathfrak{A}}^x = U_{\mathfrak{A}}^{\lceil x \rceil}$ for all $x \in \mathbb{R}_{\geq 0}$, and

$$U_{\mathfrak{A}}^{x+} = \bigcup_{x \in \mathbb{R}_{\geq 0}, y > x} U_{\mathfrak{A}}^y.$$

Suppose that E_0 is a subfield in A and \mathfrak{A} is E_0 -pure, i.e., $E_0^\times \subseteq \mathfrak{K}_{\mathfrak{A}}$, then we define the ramification index $e(\mathfrak{A}/\mathfrak{o}_{E_0})$ to be the integer e satisfying $v_{\mathfrak{A}}|_{E_0^\times} = e v_{E_0}$. We therefore have

$$\mathfrak{p}_{E_0}^i \mathfrak{P}_{\mathfrak{A}}^j = \mathfrak{P}_{\mathfrak{A}}^{ie+j}$$

and

$$\mathfrak{p}_{E_0}^i \cap \mathfrak{P}_{\mathfrak{A}}^j = \mathfrak{p}_{E_0}^{\max\{i,j/e\}}$$

for all $i, j \in \mathbb{Z}$.

In the split case $G = G^*$, i.e., when $A = A^* = \text{Mat}_n(F)$, we denote the hereditary order \mathfrak{A} by \mathfrak{A}^* .

2.5 EMBEDDING CONDITIONS

Suppose we fix an F -embedding $E_0 \hookrightarrow A$, let E_0 and \mathfrak{A} be as in the previous section, and write A_0 the centralizer of E_0 in A . We can restrict the embedding to $E_0^\times \hookrightarrow G(F)$, and denote the centralizer by $A_0^\times = Z_{G(F)}(E_0^\times)$. Under the above setup, there are many choices of \mathfrak{A} among its $Z_{G(F)}(E_0^\times)$ -conjugacy class. In this paper, we assume the conditions (i)-(iii) below, all adopted from [BH11], to fix a unique \mathfrak{A} .

- (i) [BH11, Section 3.2] If E/E_0 is an unramified extension in A_0 such that $[E : F] = n$, then we require that \mathfrak{A} is E -pure, i.e., $E^\times \subseteq \mathfrak{K}_{\mathfrak{A}}$.

Let \mathfrak{A}_0 be the centralizer of E_0 in \mathfrak{A} , i.e., $\mathfrak{A}_0 = \mathfrak{A} \cap A_0$, which is a hereditary \mathfrak{o}_{E_0} -order in A_0 , with Jacobson radical $\mathfrak{P}_{\mathfrak{A}_0} = \mathfrak{P}_{\mathfrak{A}} \cap A_0$.

- (ii) [BH11, (2.3.1)(2)] There exists a fixed integer $e(\mathfrak{A}/\mathfrak{A}_0) \geq 1$ such that

$$\mathfrak{P}_{\mathfrak{A}}^k \cap A_0 = \mathfrak{P}_{\mathfrak{A}_0}^{k/e(\mathfrak{A}/\mathfrak{A}_0)} \text{ for every } k \in \mathbb{Z}. \tag{2.5}$$

We say that \mathfrak{A} is the *canonical continuation* of \mathfrak{A}_0 in A . Under (ii), we have moreover $\mathfrak{K}_{\mathfrak{A}} \cap A_0 = \mathfrak{K}_{\mathfrak{A}_0}$.

- (iii) [BH11, (2.3.2)] \mathfrak{A}_0 is a maximal hereditary \mathfrak{o}_{E_0} -order in A_0 , i.e., $e(\mathfrak{A}_0/\mathfrak{o}_{E_0}) = 1$.

Under these conditions, \mathfrak{A} is the unique E_0 -pure hereditary order in A such that $\mathfrak{A} \cap A_0 = \mathfrak{A}_0$. Moreover, \mathfrak{A} is maximal among all E_0 -pure hereditary orders in A , and both \mathfrak{A} and \mathfrak{A}_0 are principal (by [BH11, the remark after (2.3.2)]).

By [Zin99, 0. Theorem], if the \mathfrak{o}_D -period of \mathfrak{A} is denoted by $r = r(\mathfrak{A}) = e(\mathfrak{A}/\mathfrak{o}_D)$, i.e. $\varpi_D \mathfrak{A} = \mathfrak{P}_{\mathfrak{A}}^r$, then we have an isomorphism

$$\mathfrak{A}/\mathfrak{P}_{\mathfrak{A}} \cong \text{Mat}_s(\mathbf{k}_D)^r, \tag{2.6}$$

where $s = s(\mathfrak{A}) = f/e(\mathfrak{A}/\mathfrak{o}_E)$. Once A (and hence D) is fixed, the integers r and s depend only on E ; indeed $s = s(E/F) = \gcd(f, m)$ and $r = r(E/F) = e/\gcd(d, e) = m/s$.

If K is an intermediate subfield in E/F , we write $f_K = f(E/K)$ and $e_K = e(E/K)$. By [Zin99, 1. Prop.] the centralizer $A_K = Z_A(K)$ is isomorphic to $\text{Mat}_{m_K}(D_K)$, where D_K is a division algebra over K of degree d_K^2 , with

$$d_K = \frac{d}{\gcd(d, n(K/F))} \text{ and } m_K = \gcd(m, n(E/K)).$$

Let \mathfrak{A}_K be the centralizer of K in \mathfrak{A} . It is routine to check that (E_0K, \mathfrak{A}_K) is a canonical continuation of \mathfrak{A}_{E_0K} and is also E -pure, i.e., (i)-(iii) are satisfied when we change our base field from F to K . We therefore have an isomorphism

$$\mathfrak{A}_K/\mathfrak{P}_{\mathfrak{A}_K} \cong \text{Mat}_{s_K}(\mathbf{k}_{D_K})^{r_K}, \tag{2.7}$$

where $r_K = e_K/\text{gcd}(d_K, e_K)$ and $s_K = \text{gcd}(f_K, m_K)$.

2.6 CHARACTERS

Let $\text{Nrd}_{A/F} : A^\times \rightarrow F^\times$ be the reduced norm of $G(F) = A^\times$. By [Rei03, (33.4)Theorem] and [NM43, Satz 2], $\text{Nrd}_{A/F}$ is surjective and its kernel is the commutator subgroup of $G(F)$. Therefore, any character of $G(F)$ factors through $\text{Nrd}_{A/F}$.

We define the \mathfrak{A} -level of a character ξ of A^\times as the smallest integer $a \geq -1$ such that $\xi|_{U_{\mathfrak{A}}^{a+1}}$ is trivial. This is analogous to the F -level of a character of F^\times defined in Section 1.4.

PROPOSITION 2.7. *If the F -level of a character ξ of F^\times is r , then the \mathfrak{A} -level of the character $\xi \circ \text{Nrd}_{A/F}$ is $r \cdot e(\mathfrak{A}/\mathfrak{o}_F)$.*

Proof. It suffices to show that

$$\text{Nrd}_{A/F}(U_{\mathfrak{A}}^{r \cdot e(\mathfrak{A}/\mathfrak{o}_F)}) = U_F^r \text{ for all } r \in \mathbb{R}_{\geq 0}.$$

This follows from [BF85, (2.8.3)]. □

3 THE ESSENTIALLY TAME JACQUET-LANGLANDS CORRESPONDENCE

3.1 ADMISSIBLE CHARACTERS

We recall the definition of admissible characters in [How77], [Moy86]. Given a tamely ramified finite extension E/F , let ξ be a character of E^\times . We call a pair $(E/F, \xi)$ *admissible* over F if ξ is an admissible character over F , i.e., for every intermediate subfield K between E/F ,

- (i) if ξ factors through the norm $N_{E/K}$, then $E = K$;
- (ii) if $\xi|_{U_E^1}$ factors through $N_{E/K}$ then E/K is unramified.

Two admissible pairs $(E/F, \xi)$ and $(E'/F, \xi')$ are called F -equivalent if there is $g \in \Gamma_F$ such that ${}^gE = E'$ and ${}^g\xi = \xi'$. Let $P_n(F)$ be the set of F -equivalence classes of the admissible pair $(E/F, \xi)$, with each class in $P_n(F)$ still denoted by $(E/F, \xi)$ for convenience.

Every admissible character ξ admits a factorization (see [How77, Corollary of Lemma 11] or [Moy86, Lemma 2.2.4])

$$\xi = \xi_{-1}(\xi_0 \circ N_{E/E_0}) \cdots (\xi_t \circ N_{E/E_t})(\xi_{t+1} \circ N_{E/F}), \tag{3.1}$$

where, in the notations above, the decreasing sequence of fields

$$E = E_{-1} \supseteq E_0 \supsetneq E_1 \supsetneq \cdots \supsetneq E_t \supsetneq E_{t+1} = F.$$

and the increasing E -levels $a_{-1} = 0 < a_0 < a_1 < \cdots < a_t \leq a_{t+1}$ of the characters $\xi_k \circ N_{E/E_k}$, $k = 0, \dots, t + 1$, are uniquely determined. We call the E -levels a_k the *jumps* of ξ and call the collection $\{E_k, a_k | k = 0, \dots, t\}$ the *jump data* of ξ . By convention, when $E_0 = E$, we replace $(\xi_0 \circ N_{E/E_0})\xi_{-1}$ by ξ_0 and assume that ξ_{-1} is trivial; otherwise, ξ_{-1} is tamely ramified and E/E_0 is unramified [Moy86, Definition 2.2.3]. There are certain generic conditions imposed on the jump data of the character by its admissibility, but we do not need them fully in this paper. We refer the interested reader to [Moy86, Definition 2.2.3] and [BH10, Section 8.2] for these conditions (and when E/F is totally ramified, see also [BH05b, Section 1]), and only use one of their consequences in (3.11).

We fix a (non-canonical) choice of ξ_{-1} in the factorization (3.1) as follows. We fix a choice of the wild component ξ_w of ξ to be the product

$$(\xi_0 \circ N_{E/E_0}) \cdots (\xi_t \circ N_{E/E_t})(\xi_{t+1} \circ N_{E/F})$$

which satisfies

$$\xi_w(\varpi_F) = 1 \text{ and } \xi_w \text{ has a } p\text{-power order} \tag{3.2}$$

(see [BH11, Lemma 1 of Section 4.3]), and define the tame component of ξ to be $\xi_{-1} = \xi \xi_w^{-1}$. We write

$$\Xi = \Xi(\xi) = \xi_0(\xi_1 \circ N_{E_0/E_1}) \cdots (\xi_{t+1} \circ N_{E_0/F}), \tag{3.3}$$

such that $\xi_w = \Xi \circ N_{E/E_0}$.

Suppose that E_0/F is a tamely ramified extension of degree dividing n and E/E_0 is unramified. Let Ξ be a character of $U_{E_0}^1$. Following [BH05a, Section 1.3], we call $(E_0/F, \Xi)$ an *admissible 1-pair* over F if Ξ does not factor through any norm $N_{E_0/K}$ with $F \subseteq K \subsetneq E_0$. We denote by $P_n^1(F)$ the set of F -equivalence classes of these pairs. Therefore, the map

$$P_n(F) \rightarrow P_n^1(F), (E/F, \xi) \mapsto (E_0/F, \Xi(\xi)|_{U_{E_0}^1}),$$

is well-defined and surjective. Notice that we can define the jump-data of a 1-pair, such that the jump-data of an admissible pair is the same as that of its associated 1-pair.

3.2 THE CORRESPONDENCES

Let G^* be GL_n defined over F , and G be an inner form of G^* whose F -point is isomorphic to $GL_m(D)$ for some central F -division algebra D of dimension d^2 and $n = md$. Let $\mathcal{A}_n^2(F)$ (resp. $\mathcal{A}_m^2(D)$) be the collection of equivalence classes

of irreducible smooth representations of $G^*(F)$ (resp. $G(F)$) which are essentially square-integrable modulo center. The Jacquet-Langlands correspondence is a canonical bijection

$$JL : \mathcal{A}_n^2(F) \rightarrow \mathcal{A}_m^2(D) \quad (3.4)$$

between the two collections determined by a character relation between $\pi \in \mathcal{A}_n^2(F)$ and its image $JL(\pi) \in \mathcal{A}_m^2(D)$: for every pair of semi-simple elliptic regular elements (g, g^*) , where $g \in G(F)$ and $g^* \in G^*(F)$, with the same reduced characteristic polynomial, we have [BH11, Section 1.4]

$$(-1)^{n-m} \Theta_\pi(g^*) = \Theta_{JL(\pi)}(g), \quad (3.5)$$

where Θ_π (resp. $\Theta_{JL(\pi)}$) is the character of π (resp. $JL(\pi)$).

For each representation $\pi \in \mathcal{A}_m^2(D)$, let

- (i) $f(\pi)$ be the number of unramified characters χ of F^\times that $\chi \otimes \pi \cong \pi$ (here χ is regarded as a representation of $G(F)$ by composing with the reduced norm map $\text{Nrd} : G(F) \rightarrow F^\times$), and
- (ii) $\delta(\pi)$ be the parametric degree of π (we do not require its full definition, so we only refer to [BH11, Section 2.7] for details).

It is known that $\delta(\pi)$ is a positive integer and is a multiple of $f(\pi)$. Moreover, π is supercuspidal if $\delta(\pi) = n$, while the converse is only true in the split case (when $G = G^*$).

Recall that we denote by $\mathcal{A}_n^0(F)$ the set of supercuspidal representations of $G^*(F)$. The correspondence (3.4) restricts to a bijection

$$JL : \mathcal{A}_n^0(F) \rightarrow \{\pi \in \mathcal{A}_m^2(D) \mid \delta(\pi) = n\}.$$

We call π *essentially tame* if p does not divide $\delta(\pi)/f(\pi)$. Let $\mathcal{A}_m^{\text{et}}(D)$ be the set of isomorphism classes of irreducible representations in $\mathcal{A}_m^2(D)$ which are essentially tame and satisfy $\delta(\pi) = n$. Therefore $\mathcal{A}_n^{\text{et}}(F)$ is the same collection defined in [BH05a]. Since the Jacquet-Langlands correspondence in (3.4) preserves the invariants $\delta(\pi)$ and $f(\pi)$, we have the following theorem [BH11, 2.8. Corollary 2].

THEOREM 3.1 (Essentially tame Jacquet-Langlands correspondence). *The restriction of the Jacquet-Langlands correspondence induces a bijection*

$$JL : \mathcal{A}_n^{\text{et}}(F) \rightarrow \mathcal{A}_m^{\text{et}}(D).$$

This bijection preserves the central characters on both sides. □

Bushnell and Henniart described this bijection explicitly in a way parallel to [BH05a], [BH05b], [BH10]. We recall the results briefly as follows. On the one hand, we have the bijection

$${}_F\Pi : P_n(F) \rightarrow \mathcal{A}_n^{\text{et}}(F), (E/F, \xi) \mapsto {}_F\Pi_\xi, \quad (3.6)$$

generalizing the construction of Howe [How77]. On the other hand, there is an analogous bijection

$${}_D\Pi : P_n(F) \rightarrow \mathcal{A}_m^{\text{et}}(D), (E/F, \xi) \mapsto {}_D\Pi_\xi, \tag{3.7}$$

using the constructions in [Séc04], [Séc05a], [Séc05b], [SS08]. In fact, what is constructed in [BH11] is the inverse of (3.7), using the method called ‘attached-pairs’, since the construction parallel to (3.6) exhibits some ‘novel technical difficulties’ as mentioned in [BH11, Introduction 4.]. In the split case, the attached-pair method yields the inverse of (3.6) (see 4.4 of [BH11]). The composition of the bijection in (3.6), the correspondence in Theorem 3.1, and the inverse of (3.7),

$${}_D\nu : P_n(F) \xrightarrow{{}_F\Pi} \mathcal{A}_n^{\text{et}}(F) \xrightarrow{JL} \mathcal{A}_m^{\text{et}}(D) \xrightarrow{{}_D\Pi^{-1}} P_n(F), \tag{3.8}$$

determines a tamely ramified quadratic character ${}_D\nu_\xi$ of E^\times for each admissible character ξ of E^\times , depending only on the wild part of ξ , such that for each admissible pair $(E/F, \xi)$, the pair $(E/F, {}_D\nu_\xi \cdot \xi)$ is also admissible and

$${}_D\nu(E/F, \xi) = (E/F, {}_D\nu_\xi \cdot \xi). \tag{3.9}$$

We call this character ${}_D\nu_\xi$ the *rectifier* of ξ (for the Jacquet-Langlands correspondence). Using the First and Second Comparison Theorems of [BH11], we can compute the values of ${}_D\nu_\xi$. To express these values, we need the knowledge of certain invariants of finite symplectic modules, which will be described in Section 3.8. Finally, with the expression of ${}_D\nu_\xi$, we see that we can describe the correspondence in Theorem 3.1 explicitly, using (3.6), (3.7), and (3.8).

3.3 SOME SUBGROUPS

We recall certain subgroups of $G(F)$. Suppose that the jump data $\{E_k, a_k | k = 0, \dots, t\}$ are defined by the factorization (3.1) of an admissible pair $(E/F, \xi)$, or equivalently, of its associated 1-pair $(E_0/F, \Xi)$. We require that (E_0, \mathfrak{A}) satisfy the conditions in Section 2.5. We write A_k the centralizer of E_k in A and $\mathfrak{A}_k = A_k \cap \mathfrak{A}$. We can then define $\mathfrak{P}_{\mathfrak{A}_k}$, $U_{\mathfrak{A}_k}$, $U_{\mathfrak{A}_k}^x$, and $U_{\mathfrak{A}_k}^{x+}$ for $x \in \mathbb{R}_{\geq 0}$ analogously as in Section 2.4. Following [Gra07, Definition 4.1], we construct the pro- p subgroups

$$\begin{aligned} H^1(\Xi, \mathfrak{A}) &= U_{\mathfrak{A}_0}^1 U_{\mathfrak{A}_1}^{(a_0 e(\mathfrak{A}_1/\mathfrak{o}_E)/2)^+} \dots U_{\mathfrak{A}_t}^{(a_{t-1} e(\mathfrak{A}_t/\mathfrak{o}_E)/2)^+} U_{\mathfrak{A}}^{(a_t e(\mathfrak{A}/\mathfrak{o}_E)/2)^+} \text{ and} \\ J^1(\Xi, \mathfrak{A}) &= U_{\mathfrak{A}_0}^1 U_{\mathfrak{A}_1}^{a_0 e(\mathfrak{A}_1/\mathfrak{o}_E)/2} \dots U_{\mathfrak{A}_t}^{a_{t-1} e(\mathfrak{A}_t/\mathfrak{o}_E)/2} U_{\mathfrak{A}}^{a_t e(\mathfrak{A}/\mathfrak{o}_E)/2}. \end{aligned} \tag{3.10}$$

We also construct the subgroups

$$\begin{aligned} J(\Xi, \mathfrak{A}) &= U_{\mathfrak{A}_0} U_{\mathfrak{A}_1}^{a_0 e(\mathfrak{A}_1/\mathfrak{o}_E)/2} \dots U_{\mathfrak{A}_t}^{a_{t-1} e(\mathfrak{A}_t/\mathfrak{o}_E)/2} U_{\mathfrak{A}}^{a_t e(\mathfrak{A}/\mathfrak{o}_E)/2} \text{ and} \\ \mathbf{J}(\Xi, \mathfrak{A}) &= E^\times J(\Xi, \mathfrak{A}) = E_0^\times J(\Xi, \mathfrak{A}). \end{aligned}$$

We abbreviate these groups by H^1, J^1, J and \mathbf{J} if the admissible character ξ is fixed. Notice that H^1, J^1, J are compact subgroups of $G(F)$ and \mathbf{J} is a compact-mod-center subgroup of $G(F)$.

In [Séc04, Section 3.1], these subgroups are defined based on a simple stratum $[\mathfrak{A}, -v_{\mathfrak{A}}(\beta), 0, \beta]$, where β is a suitable element in E_0 , depending on Ξ and such that

$$-v_{\mathfrak{A}}(\beta) = \text{the } E\text{-level of } \Xi \circ N_{E/E_0}.$$

The group $H^1(\Xi, \mathfrak{A})$ is denoted by $H^1(\beta, \mathfrak{A})$ in *loc. cit.* (and similarly for the other subgroups). This construction is an obvious generalization of [BK93, Section 3.1] (see also [BH11, Section 2.5] and the Comment therewithin).

3.4 SIMPLE CHARACTERS

Given an admissible 1-pair $(E_0/F, \Xi)$ and a finite unramified extension E/E_0 , we define $H^1(\Xi, \mathfrak{A})$ as in (3.10). Using the idea of [Moy86, Section 3.2] (see also [Séc04, Definitions 3.22, 3.45, Proposition 3.47]), we attach to $(E_0/F, \Xi)$ a simple character $(H^1(\Xi, \mathfrak{A}), \theta_{\Xi, E})$ as follows. Suppose that Ξ admits a factorization of the form (3.3), with each $\xi_k \circ N_{E_0/E_k}$, where $k = 0, \dots, t + 1$, a character of $U_{E_0}^1$. The generic conditions on the factorization imply that for each ξ_k there is $c_k \in E_k \cap \mathfrak{p}_E^{-a_k}$ such that $E_{k+1}[c_k] = E_k$ and

$$\xi_k \circ N_{E/E_k}(1 + x) = \psi_F(\text{tr}_{E/F}(c_k x)) \text{ for all } x \in \mathfrak{p}_E^{(a_k/2)^+}. \tag{3.11}$$

Note that the element c_k can be chosen mod $\mathfrak{p}_E^{-a_k/2}$. We denote the character on the right side of (3.11) by ψ_{c_k} .

We define a character $\theta_{\Xi, E}$ of the subgroup $H^1(\Xi, \mathfrak{A})$ in (3.10) by the following inductive procedure. We first define a character θ_{t+1} on the subgroup $U_{\mathfrak{A}_{t+1}}^{(a_t e(\mathfrak{A}_{t+1}/\mathfrak{o}_E)/2)^+}$ (note that indeed $E_{t+1} = F$ and $\mathfrak{A}_{t+1} = \mathfrak{A}$) by

$$\xi_{t+1} \circ \text{Nrd}_{A/F} \text{ on } U_{\mathfrak{A}_{t+1}}^{(a_t e(\mathfrak{A}_{t+1}/\mathfrak{o}_E)/2)^+}.$$

Inductively, suppose θ_{k+1} is defined, we construct θ_k on the subgroup

$$U_{\mathfrak{A}_k}^{(a_{k-1} e(\mathfrak{A}_k/\mathfrak{o}_E)/2)^+} U_{\mathfrak{A}_{k+1}}^{(a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2)^+} \dots U_{\mathfrak{A}}^{(a_t e(\mathfrak{A}/\mathfrak{o}_E)/2)^+}$$

by

$$\begin{aligned} &(\xi_k \circ \text{Nrd}_{A_k/E_k}) \cdots (\xi_{t+1} \circ \text{Nrd}_{A/F}) \text{ on } U_{\mathfrak{A}_k}^{(a_{k-1} e(\mathfrak{A}_k/\mathfrak{o}_E)/2)^+} \text{ and} \\ &(\psi_{c_k} \circ \text{trd}_{\mathfrak{A}_k/E_k}) \theta_{k+1} \text{ on } U_{\mathfrak{A}_{k+1}}^{(a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2)^+} \dots U_{\mathfrak{A}}^{(a_t e(\mathfrak{A}/\mathfrak{o}_E)/2)^+}. \end{aligned} \tag{3.12}$$

On the intersection $U_{\mathfrak{A}_k}^{(a_k e(\mathfrak{A}_k/\mathfrak{o}_E)/2)^+}$, we have

$$\theta_{k+1} = (\xi_{k+1} \circ \text{Nrd}_{A_{k+1}/E_{k+1}}) \cdots (\xi_{t+1} \circ \text{Nrd}_{A/F})$$

and

$$\psi_{c_k} \circ \text{trd}_{\mathfrak{A}_k/E_k} = \xi_k \circ \text{Nrd}_{A_k/E_k}.$$

By Proposition 2.7 and (3.11), the two characters in (3.12) agree on the intersection, and so θ_k is defined. Finally, we define $\theta_{\Xi,E} = \theta_0$ on $H^1(\Xi, \mathfrak{A})$. By construction, this character $\theta_{\Xi,E}$ is normalized by $\mathfrak{K}_{\mathfrak{A}_0}$, hence it is a simple character by [Gra07, 5.3 Definition.(i)].

To get back our character Ξ from $\theta_{\Xi,E}$, just notice that $\theta_{\Xi,E}|_{U_E^1}$ factors through N_{E/E_0} , hence there is a unique character Ξ of $U_{E_0}^1$ such that $\Xi \circ N_{E/E_0} = \theta_{\Xi,E}|_{U_E^1}$.

3.5 LOCAL CONSTRUCTIONS OF ATTACHED PAIRS

We briefly summarize the construction in [BH11, Sec. 4] of the bijection ${}_D\Pi$ in (3.7). We will distinguish between the ‘level-zero’ case and the ‘positive-level’ case (we refer to [BH11, Sections 2.4 and 2.6] the definition of these cases). As we remarked in the Introduction, it is indeed the inverse of ${}_D\Pi$ that we are going to describe.

In the level-zero case, the construction is similar to the one in the split case. Each level zero $\pi \in \mathcal{A}_m^{\text{et}}(D)$ contains a representation $(\text{GL}_m(\mathfrak{o}_D), \lambda)$, called a maximal simple type of level zero, inflated from an irreducible cuspidal representation $(\text{GL}_m(\mathbf{k}_D), \bar{\lambda})$. This representation $\bar{\lambda}$ corresponds, via Green’s parametrization [Gre55], to a $\mathbf{k}_E/\mathbf{k}_F$ -regular character $\bar{\xi}$ of \mathbf{k}_E^\times , where E/F is the unramified extension of degree n . We define the character ξ of E^\times such that $\xi|_{\mathfrak{o}_E^\times}$ is the inflation of $\bar{\xi}$ and $\xi|_{F^\times}$ is the central character of π . By [BH11, 4.2. Proposition], the attached pair $(E/F, \xi)$ is admissible and the correspondence

$$\mathcal{A}_m^{\text{et}}(D)_{\text{level}-0} \rightarrow P_n(F)_{\text{level}-0}, \pi \mapsto (E/F, \xi), \tag{3.13}$$

is bijective. We can show that $\pi = \text{cInd}_{\mathbf{J}}^{G(F)} \Lambda$, where the condition $\delta(\pi) = n$ implies that $\mathbf{J} = F^\times \text{GL}_m(\mathfrak{o}_D)$, and (\mathbf{J}, Λ) is defined by the conditions

$$\Lambda|_{\text{GL}_m(\mathfrak{o}_D)} = \lambda \text{ and } \Lambda|_{F^\times} \text{ is a multiple of } \xi|_{F^\times}. \tag{3.14}$$

The representation (\mathbf{J}, Λ) is called an extended maximal simple type of level zero.

In the positive level case, we first recall the construction of a extended maximal simple type in general. Suppose we have a simple character (H^1, θ) . For example, we can construct a simple character $\theta = \theta_{\Xi,E}$ as in Section 3.4 using an admissible pair $(E/F, \xi)$. We notice that the commutator subgroup $[J^1, J^1]$ lies in H^1 [BK93, (3.1.15)]. By [Séc04, Théorème 3.52], the above simple character θ induces an non-degenerate alternating bilinear form

$$h_\theta(x, y) = \theta([1 + x, 1 + y]), \text{ for all } 1 + x, 1 + y \in J^1, \tag{3.15}$$

on the \mathbb{F}_p -vector space

$${}_A\mathfrak{V}_\xi := J^1/H^1.$$

The classical theory of Heisenberg representation implies that there is a unique representation $\bar{\eta}$ of $J^1/\ker \theta$ containing the character θ of $H^1/\ker \theta$ as a central

character. We then define η as the inflation of $\bar{\eta}$ to J^1 . By [Séc05a, Théorème 2.28], there exists a unique irreducible representation (J, κ) , which is called a β -extension (or wide extension in [BH11]) of η and satisfies certain conditions on its intertwining in $G(F)$ (see [BH11, (2.5.5)]). We now choose a maximal simple type $(\mathrm{GL}_{m_0}(\mathfrak{o}_{D_0}), \sigma)$ of $A_0^\times = Z_{G(F)}(E_0^\times)$ of level zero and inflate it to a representation (J, σ) , since we know that $J = \mathrm{GL}_{m_0}(\mathfrak{o}_{D_0})J^1$. We obtain a maximal simple type (J, λ) , where $\lambda = \kappa \otimes \sigma$. By [Séc05b, Théorème 5.2], there exists an irreducible representation Λ of $\mathbf{J} = E_0^\times J^1$ (by the condition $\delta(\pi) = n$), extending λ and whose compact-induction to $G(F)$ is irreducible and supercuspidal. The representation (\mathbf{J}, Λ) is called an extended maximal simple type. By [BH11, Lemma 2 of Section 4.3], we can fix a unique extended type containing (J, λ) and satisfying the (non-canonical) conditions:

$$\varpi_F \in \ker \Lambda \text{ and } \det \Lambda \text{ has a } p\text{-power order.}$$

Following [BH11, Section 3 and 4], we have to approach indirectly to describe the inverse of ${}_D\Pi$. Suppose that we are given a representation $\pi \in \mathcal{A}_m^{\mathrm{et}}(D)$ of positive level. By [SS08, Théorème 5.21], it contains an extended maximal simple type (\mathbf{J}, Λ) of the above form, such that $\Lambda|_{H^1}$ is a multiple of a simple character (H^1, θ) . There is a unique character ξ_w of E^\times , depending on $\theta|_{U_{\mathfrak{a}_0}^1}$, satisfying the conditions in (3.2). In particular, we have

$$\theta|_{U_{\mathfrak{a}_0}^1} = \Xi \circ \mathrm{Nrd}_{A_0/E_0}, \text{ such that } \xi_w|_{U_E^1} = \Xi \circ N_{E/E_0}.$$

By the discussion of the previous paragraph, we can attach to ξ_w an extended maximal simple type (\mathbf{J}, Λ_w) such that $\Lambda \cong \Lambda_{-1} \otimes \Lambda_w$ for a uniquely determined extended maximal simple type $(\mathbf{J}, \Lambda_{-1})$ of level zero. Attached to $(\mathbf{J}, \Lambda_{-1})$ is a level zero character ξ_{-1} of E^\times admissible over E_0 , as mentioned in the level zero case. Finally, by [BH11, 4.3. Proposition], the attached pair $(E/F, \xi)$, where $\xi = \xi_{-1}\xi_w$, is admissible and independent of the various choices above. We call $(E/F, \xi)$ a pair attached to π .

The technical part is to show that the attaching map

$$\mathcal{A}_m^{\mathrm{et}}(D) \rightarrow P_n(F), \pi \mapsto (E/F, \xi)$$

is well-defined and injective. This is done in the Parametrization Theorem of [BH11, Section 6]. The composition (3.8) is then injective (since the maps ${}_F\Pi$ and JL are known to be bijective) and preserves the restriction of each character to the subgroup $F^\times U_E^1$, which is of finite index of E^\times . Therefore, the map in (3.8) and hence ${}_D\Pi$ in (3.7), is bijective.

3.6 FINITE SYMPLECTIC MODULES

Since the group \mathbf{J} normalizes the subgroups H^1, J^1 , and the simple character θ of H^1 , it acts on the finite quotient ${}_A\mathfrak{V}_\xi := J^1/H^1$. This quotient is denoted by ${}_{A^*}\mathfrak{V}_\xi$ in the split case $A = A^*$, which is studied in [BH10] and [Tam]. Notice

that the quotient is clearly a finite dimensional \mathbb{F}_p -vector space. The action of \mathbf{J} induces a symplectic $\mathbb{F}_p\mathbf{J}$ -module structure on this space with respect to the non-degenerate alternating form h_θ in (3.15). We have a decomposition of

$${}_A\mathfrak{A}_\xi = {}_A\mathfrak{A}_{\xi,0} \oplus \cdots \oplus {}_A\mathfrak{A}_{\xi,t}, \tag{3.16}$$

into $\mathbb{F}_p\mathbf{J}$ -submodules, where

$$\begin{aligned} {}_A\mathfrak{A}_{\xi,k} &= \frac{U_{\mathfrak{A}_{k+1}}^{a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2}}{U_{\mathfrak{A}_k}^{a_k e(\mathfrak{A}_k/\mathfrak{o}_E)/2} U_{\mathfrak{A}_{k+1}}^{(a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2)+}} \\ &= \frac{\mathfrak{P}_{\mathfrak{A}_{k+1}}^{a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2}}{\mathfrak{P}_{\mathfrak{A}_k}^{a_k e(\mathfrak{A}_k/\mathfrak{o}_E)/2} + \mathfrak{P}_{\mathfrak{A}_{k+1}}^{(a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2)+}} \end{aligned} \tag{3.17}$$

for $k = 0, \dots, t$. In this paper, We call this decomposition the *coarse decomposition* of ${}_A\mathfrak{A}_\xi$. By [Séc04, Proposition 3.9], the decomposition (3.16) is orthogonal.

In the sequel, we will be interested in the adjoint action of E^\times on ${}_A\mathfrak{A}_\xi$ restricted from that of \mathbf{J} , which factors through the finite group $\Psi_{E/F} := E^\times/F^\times(E^\times \cap J^1)$.

The following Proposition appears in [BH11, Proposition 5.6]. We re-interpret its proof here.

PROPOSITION 3.2. *If E/F is totally ramified, then ${}_A\mathfrak{A}_\xi \cong {}_{A^*}\mathfrak{A}_\xi$.*

Proof. From the proofs of Propositions 4.1 and 4.2 (which are purely algebra and do not require the knowledge of this section), we see that the totally ramified condition implies that

$$\mathfrak{P}_{\mathfrak{A}_k^j}^j / \mathfrak{P}_{\mathfrak{A}_k^{j+1}}^{j+1} \cong \mathfrak{P}_{\mathfrak{A}_k^j}^j / \mathfrak{P}_{\mathfrak{A}_k^j}^{j+1} \cong \text{Ind}_1^{\Psi_{E/E_k}} \mathbf{k}_F \tag{3.18}$$

as a $\mathbf{k}_F \Psi_{E/E_k}$ -module, for all $j \in \mathbb{Z}$ and $k = 1, \dots, t + 1$. Moreover, we know that in the split case the index $e(\mathfrak{A}_k^*/\mathfrak{o}_E)$ (appearing in the powers in (3.17), when $A = A^*$) is always 1, and in general each $e(\mathfrak{A}_k/\mathfrak{o}_E)$ (again appearing in (3.17)) divides $f(E/E_k)$ (remember that $f(E/E_k)/e(\mathfrak{A}_k/\mathfrak{o}_E)$ is the integer s_{E_k} appearing in (2.7), when $K = E_k$), which is equal to 1 in the totally ramified case. Hence from the expression in (3.17) and using (3.18), we see that both ${}_A\mathfrak{A}_{\xi,k}$ and ${}_{A^*}\mathfrak{A}_{\xi,k}$ are isomorphic to

$$\frac{\mathfrak{P}_{\mathfrak{A}_{k+1}}^{a_k/2}}{\mathfrak{P}_{\mathfrak{A}_k}^{a_k/2} + \mathfrak{P}_{\mathfrak{A}_{k+1}}^{(a_k/2)+}}.$$

Hence their sums ${}_A\mathfrak{A}_\xi$ and ${}_{A^*}\mathfrak{A}_\xi$ are also isomorphic. □

3.7 INVARIANTS OF FINITE SYMPLECTIC MODULES

Let Γ be a finite cyclic group whose order is not divisible by p . We call a finite $\mathbb{F}_p\Gamma$ -module V *symplectic* if there is a non-degenerate alternating form $\mathbf{h} : V \times V \rightarrow \mathbb{F}_p$ which is Γ -invariant, in the sense that

$$\mathbf{h}(\gamma v_1, \gamma v_2) = \mathbf{h}(v_1, v_2), \text{ for all } \gamma \in \Gamma, v_1, v_2 \in V.$$

The simple module V_λ corresponding to a character $\lambda \in \text{Hom}(\Gamma, \overline{\mathbb{F}}_p^\times)$ is the field $\mathbb{F}_p[\lambda(\Gamma)]$ generated over \mathbb{F}_p by the image $\lambda(\Gamma)$, with Γ -action

$$\gamma v = \lambda(\gamma)v, \text{ for all } \gamma \in \Gamma, v \in V_\lambda.$$

Its \mathbb{F}_p -linear dual $V_\lambda^* = \text{Hom}(V_\lambda, \mathbb{F}_p)$ is isomorphic to $V_{\lambda^{-1}}$ by the map

$$V_{\lambda^{-1}} \rightarrow V_\lambda^*, v \mapsto (w \mapsto \text{tr}_{\mathbb{F}_p[\lambda(\Gamma)]/\mathbb{F}_p}(wv)),$$

such that the canonical pairing $\langle \cdot, \cdot \rangle : V_\lambda \times V_{\lambda^{-1}} \rightarrow \mathbb{F}_p$ is Γ -invariant.

We recall some basic facts from [BF83, (8.2.3)] and [BH10, Sec. 3, Prop. 4].

PROPOSITION 3.3. (i) *An indecomposable symplectic $\mathbb{F}_p\Gamma$ -module is isomorphic to either one of the following two kinds,*

- (a) *a hyperbolic module of the form $\mathbf{V}_\lambda = V_\lambda \oplus V_{\lambda^{-1}}$ such that either $\lambda^2 = 1$ or $V_\lambda \not\cong V_{\lambda^{-1}}$, with the alternating form*

$$\mathbf{h}_{\mathbf{V}_\lambda}((v_1, v_1^*), (v_2, v_2^*)) = \langle v_1, v_2^* \rangle - \langle v_2, v_1^* \rangle,$$

for all $(v_1, v_1^), (v_2, v_2^*) \in \mathbf{V}_\lambda$;*

- (b) *an anisotropic module of the form V_λ with $\lambda^2 \neq 1$ and $V_\lambda \cong V_{\lambda^{-1}}$. In this case, $[\mathbb{F}_p[\lambda(\Gamma)] : \mathbb{F}_p]$ is even and the alternating form $\mathbf{h}_{\mathbf{V}_\lambda}$ is defined, up to Γ -isometry, by*

$$(v_1, v_2) \mapsto \text{tr}_{\mathbb{F}_p[\lambda(\Gamma)]/\mathbb{F}_p}(\alpha v_1 \bar{v}_2), \text{ for all } v_1, v_2 \in V_\lambda,$$

where $v \mapsto \bar{v}$ is the \mathbb{F}_p -automorphism of $\mathbb{F}_p[\lambda(\Gamma)]$ of order 2 and $\alpha \in \mathbb{F}_p[\lambda(\Gamma)]^\times$ satisfies $\bar{\alpha} = -\alpha$.

- (ii) *If V_λ is anisotropic and $\mathbb{F}_p[\lambda(\Gamma)]_\pm$ denotes the subfield of $\mathbb{F}_p[\lambda(\Gamma)]$ such that $\mathbb{F}_p[\lambda(\Gamma)]/\mathbb{F}_p[\lambda(\Gamma)]_\pm$ is quadratic, then $\lambda(\Gamma)$ is a subgroup of $\ker(N_{\mathbb{F}_p[\lambda(\Gamma)]/\mathbb{F}_p[\lambda(\Gamma)]_\pm})$*

- (iii) *The Γ -isometry class of a symplectic $\mathbb{F}_p\Gamma$ -module (V, \mathbf{h}) is determined by the underlying $\mathbb{F}_p\Gamma$ -module V .*

Part (iii) is particularly useful because, when we talk about invariants of Γ -isometry classes of symplectic $\mathbb{F}_p\Gamma$ -modules, we do not have to write down the alternating forms explicitly.

Given a finite symplectic $\mathbb{F}_p\Gamma$ -module \mathfrak{V} , we attach a sign $t_{\Gamma}^0(\mathfrak{V}) \in \{\pm 1\}$ and a quadratic character $t_{\Gamma}^1(\mathfrak{V})$ of Γ . We also set

$$t_{\Gamma}(\mathfrak{V}) = t_{\Gamma}^0(\mathfrak{V})t_{\Gamma}^1(\mathfrak{V})(\gamma),$$

where γ is any generator of Γ . We call these *t-factors* of \mathfrak{V} . We recall from [BH10, Section 3] the definition the t-factors.

(i) If Γ acts on \mathfrak{V} trivially, then

$$t_{\Gamma}^0(\mathfrak{V}) = 1 \text{ and } t_{\Gamma}^1(\mathfrak{V}) \equiv 1.$$

(ii) Let \mathfrak{V} be an indecomposable symplectic $\mathbb{F}_p\Gamma$ -module.

(a) If $\mathfrak{V} = \mathfrak{V}_{\lambda} \oplus \mathfrak{V}_{\lambda^{-1}}$ is hyperbolic, then

$$t_{\Gamma}^0(\mathfrak{V}) = 1 \text{ and } t_{\Gamma}^1(\mathfrak{V}) = \text{sgn}_{\lambda(\Gamma)}(\mathfrak{V}_{\lambda}).$$

Here $\text{sgn}_{\lambda(\Gamma)}(\mathfrak{V}_{\lambda}) : \Gamma \rightarrow \{\pm 1\}$ is the character whose image $\gamma \mapsto \text{sgn}_{\lambda(\gamma)}(\mathfrak{V}_{\lambda})$ is the signature of the multiplication by $\lambda(\gamma)$ as a permutation of the set \mathfrak{V}_{λ} .

(b) If $\mathfrak{V} = \mathfrak{V}_{\lambda}$ is anisotropic, then

$$t_{\Gamma}^0(\mathfrak{V}) = -1 \text{ and } t_{\Gamma}^1(\mathfrak{V})(\gamma) = \left(\frac{\gamma}{\ker(N_{\mathbb{F}_p[\lambda(\Gamma)]/\mathbb{F}_p[\lambda(\Gamma)]_{\pm}})} \right) \text{ for any } \gamma \in \Gamma.$$

Here (\cdot) is the symbol defined as follows: for every finite cyclic group H ,

$$\left(\frac{x}{H} \right) = \begin{cases} 1 & \text{if } x \in H^2, \\ -1 & \text{otherwise.} \end{cases}$$

(iii) If \mathfrak{V} decomposes into an orthogonal sum $\mathfrak{V}_1 \perp \cdots \perp \mathfrak{V}_t$ of indecomposable symplectic $\mathbb{F}_p\Gamma$ -modules, then

$$t_{\Gamma}^i(\mathfrak{V}) = t_{\Gamma}^i(\mathfrak{V}_1) \cdots t_{\Gamma}^i(\mathfrak{V}_t) \text{ for } i = 0, 1.$$

Notice that when $p = 2$, the order of Γ is odd. In this case, $t_{\Gamma}^1(\mathfrak{V})$ is always trivial, because all signature characters and symbols (\cdot) are trivial.

3.8 VALUES OF RECTIFIERS

Given a tamely ramified extension E/F and an F -admissible character ξ of E^{\times} , let ${}_D\nu_{\xi}$ be the rectifier of ξ defined in (3.9). To describe the values of ${}_D\nu_{\xi}$, we need to impose a condition on ϖ_E defined in (2.1):

$$\varpi_E \in E_0, \tag{3.19}$$

where E_0 be the first field appearing in the factorization (3.1) of ξ . This condition is the same as in the Second Comparison Theorem of [BH11, Section 7], where we further require that $\varpi_E^r \in F$ for some integer r coprime to p . Indeed, from the assumption in (2.1) this extra requirement is automatic in our situation. Under (3.19), the roots of unity $z_{E/F}, z_{\phi^i}$ (defined in Section 2.2), and others related to ϖ_E in later sections all depend on the first field E_0 in the jump data of ξ .

The values of the rectifier $D\nu_\xi$ depends on the t-factors

$$t_\mu^1(A\mathfrak{A}_\xi), t_\mu^1(A^*\mathfrak{A}_\xi), t_{\langle\varpi\rangle}(A\mathfrak{A}_\xi) \text{ and } t_{\langle\varpi\rangle}(A^*\mathfrak{A}_\xi),$$

where $\mu = \mu_{E/F}$ and $\langle\varpi\rangle = \langle\varpi\rangle_{E/F}$ abbreviate the following subgroups of $\Psi_{E/F} = E^\times / F^\times U_E^1$,

$$\begin{aligned} \mu &:= \mu_E / \mu_F \text{ and} \\ \langle\varpi\rangle &:= \text{the subgroup generated by the image of } \varpi_E. \end{aligned} \tag{3.20}$$

By the First and Second Comparison Theorems of [BH11], the rectifier $D\nu_\xi$ has values

$$\begin{aligned} D\nu_\xi|_{\mu_E} &= t_\mu^1(A\mathfrak{A}_\xi)t_\mu^1(A^*\mathfrak{A}_\xi) \text{ and} \\ D\nu_\xi(\varpi_E) &= (-1)^{n-m+f_\varpi-m\varpi} t_{\langle\varpi\rangle}(A\mathfrak{A}_\xi)t_{\langle\varpi\rangle}(A^*\mathfrak{A}_\xi), \end{aligned} \tag{3.21}$$

where $f_\varpi := [E : F[\varpi_E]] = f(E/F[\varpi_E])$ and $m_\varpi = \gcd(m, f_\varpi)$.

In the case when E/F is totally ramified, Proposition 3.2 implies that

$$D\nu_\xi \text{ is unramified and } D\nu_\xi(\varpi_E) = (-1)^{n-m},$$

as stated in [BH11, 5.3.Theorem].

4 FINITE SYMPLECTIC MODULES

4.1 STANDARD MODULES OF CENTRAL SIMPLE ALGEBRA

Let \mathfrak{A} be the hereditary E -pure order in A , as discussed in Sections 2.4 and 2.5. The isomorphism (2.6) implies that $\mathfrak{P}_\mathfrak{A}^j / \mathfrak{P}_\mathfrak{A}^{j+1} \cong \text{Mat}_s(\mathbf{k}_D)^r$ for all $j \in \mathbb{Z}$, where $s = \gcd(f, m)$ and $r = e / \gcd(d, e)$. We denote this quotient by $(\text{Mat}_s(\mathbf{k}_D)^r)_j$ when we want to emphasize the index j . Notice that as $\mathbf{k}_F \Psi_{E/F}$ -modules, all $(\text{Mat}_s(\mathbf{k}_D)^r)_j$, for j ranges over all \mathbb{Z} , are isomorphic to each other. When $\mathfrak{A} = \mathfrak{A}^*$ and $j = 0$, we know that $\mathfrak{U}_{\mathfrak{A}^*} := \mathfrak{A}^* / \mathfrak{P}_{\mathfrak{A}^*} \cong \text{Mat}_n(\mathbf{k}_F)$ admits a root-space decomposition

$$\mathfrak{U}_{\mathfrak{A}^*} \cong \mathfrak{U}_0 \bigoplus_{[\lambda] \in \Gamma_F \setminus \Phi} \mathfrak{U}_{[\lambda]},$$

where $\mathfrak{U}_0 \cong \mathfrak{o}_E / \mathfrak{p}_E$ on which $\Psi_{E/F}$ acts trivially, and $\mathfrak{U}_{[\lambda]}$ is the \mathbf{k}_F -subspace on which $\Psi_{E/F}$ acts by the character λ . Note that the equivalence class of the $\mathbf{k}_F \Psi_{E/F}$ -module $\mathfrak{U}_{[\lambda]}$ depends only on the Γ_F -orbit of λ .

For future computation, we rewrite the above decomposition as

$$\mathfrak{U}_{\mathfrak{A}^*} \cong \bigoplus_{[g] \in \Gamma_E \backslash \Gamma_F / \Gamma_E} \mathfrak{U}_{[g]} \tag{4.1}$$

using the identification in Proposition 2.1. Here $\mathfrak{U}_{[g]} \cong \mathbf{k}_{E^{(gE)}}$ as a \mathbf{k}_F -vector space for each $[g] \in \Gamma_E \backslash \Gamma_F / \Gamma_E$, and the $\Psi_{E/F}$ -action on each $v \in \mathfrak{U}_{[g]}$ is given as follows: if $[g] = [\sigma^i \phi^j]$, then

$$z v = (z^{q^j-1})^{-1} v \text{ for all } z \in \mu_E \text{ and } \varpi_E v = (z_e^i z_{\phi^j})^{-1} v, \tag{4.2}$$

where z_e and z_{ϕ^j} are defined in Section 2.2.

For general inner form G , we first consider a simple case when A is a division algebra D . We write $(\mathbf{k}_D)_j := \mathfrak{P}_D^j / \mathfrak{P}_D^{j+1}$ for each $j \in \mathbb{Z}$.

PROPOSITION 4.1. *For each $j \in \mathbb{Z}$, the $\mathbf{k}_F \Psi_{E/F}$ -module $(\mathbf{k}_D)_j$ is isomorphic to*

$$\bigoplus_{[\sigma^i \phi^{hj}] \in \Gamma_E \backslash \Gamma_F / \Gamma_E} \mathfrak{U}_{[\sigma^i \phi^{hj}]}. \tag{4.3}$$

Proof. Recall that, if we denote by K the maximal unramified extension in D (of degree n over F), then

$$\mathfrak{P}_D^i = \cdots \oplus \mathfrak{p}_K \varpi_D^i \oplus \mathfrak{o}_K \varpi_D^{i+1} \oplus \cdots.$$

Hence we can use $x \varpi_D^i$, with $x \in \mu_K \cup \{0\}$, as a representative in \mathfrak{P}_D^i of an element in $(\mathbf{k}_D)_j$. We regard $(\mathbf{k}_D)_j$ as a \mathbf{k}_E -vector space of dimension e such that $z \in \mu_E$ acts on each piece $(\mathbf{k}_E)_j$ by the character $\left[\frac{1}{\phi^{hj}} \right] (z) = z^{1-q^{hj}}$ as in (2.4). Therefore, a Frobenius reciprocity argument (which is still valid when p does not divide $\#\Psi_{E/F}$) implies that

$$(\mathbf{k}_D)_j \cong \text{Ind}_{\mu_E / \mu_F}^{\Psi_{E/F}} (\mathbf{k}_E)_j;$$

more precisely, the action of ϖ_E has eigenvalues $(z_e^i z_{\phi^{hj}})^{-1}$, $i = 0, \dots, e - 1$, where we recall that $z_{\phi^{hj}}$ is an e th root of $z_{E/F}^{q^{hj}-1}$. Hence we have the decomposition (4.3) with the $\Psi_{E/F}$ -action on each component $\mathfrak{U}_{[\sigma^i \phi^{hj}]}$ as in (4.2) for each fixed j . \square

For the general A , if we write $(\text{Mat}_s(\mathbf{k}_D)^r)_{j'} := \mathfrak{P}_{\mathfrak{A}}^{j'} / \mathfrak{P}_{\mathfrak{A}}^{j'+1}$ for each $j' \in \mathbb{Z}$, then we have the following result.

PROPOSITION 4.2. *For each $j' \in \mathbb{Z}$, we have a decomposition*

$$(\text{Mat}_s(\mathbf{k}_D)^r)_{j'} \cong \bigoplus_{\substack{[\sigma^i \phi^j] \in \Gamma_E \backslash \Gamma_F / \Gamma_E \\ j \equiv hj' \pmod{f/s}}} \mathfrak{U}_{[\sigma^i \phi^j]}.$$

as a $\Psi_{E/F}$ -module. (Recall that $f/s = d / \gcd(d, e) = e(\mathfrak{A} / \mathfrak{o}_E)$.)

Proof. Denote by $E_D = E \cap D$ the maximal subfield contained in both E and D , then $[\mathbf{k}_{E_D} : \mathbf{k}_F] = f/s$ and $[\mathbf{k}_D : \mathbf{k}_{E_D}] = \gcd(d, e) = e/r$. By Proposition 4.1, we know that $z \in \mu_{E_D}$ acts on $(\text{Mat}_s(\mathbf{k}_D)^r)_{j'}$ as a sum of $(\mathbf{k}_{E_D})_{hj'}$, i.e., z acts by the character $\left[\begin{smallmatrix} 1 \\ \bar{\phi}^{hj'} \end{smallmatrix} \right] (z) = z^{1-q^{hj'}}$, where $\bar{\phi}$ is the image of ϕ under the natural projection

$$\Gamma_{\mathbf{k}_E/\mathbf{k}_F} \rightarrow \Gamma_{\mathbf{k}_{E_D}/\mathbf{k}_F}. \tag{4.4}$$

(Note that the arguments above concerning Proposition 4.1 still valid even though E_D may not be a maximal subfield of D .) We now consider the \mathbf{k}_{E_D} -embeddings (where all choices are conjugate to each other)

$$\mathbf{k}_{E_D} \rightarrow \mathbf{k}_E \rightarrow \text{Mat}_s(\mathbf{k}_{E_D}) \rightarrow \text{Mat}_s(\mathbf{k}_D),$$

Notice that \mathbf{k}_E is a maximal subfield of $\text{Mat}_s(\mathbf{k}_{E_D})$. By the “twisted group ring decomposition”, we know that $z \in \mu_E$ acts on $\text{Mat}_s(\mathbf{k}_{E_D})$ as a sum of $((\mathbf{k}_E)_{hj'})^s$, i.e., z acts on each of the s summands of \mathbf{k}_E by the character $\left[\begin{smallmatrix} 1 \\ \phi^j \end{smallmatrix} \right] (z) = z^{1-q^j}$, for ϕ^j ranges over the s pre-images of $\bar{\phi}^{hj'}$ under the natural projection (4.4). We denote this μ_E -module by $\text{Mat}_s(\mathbf{k}_{E_D})_{j'}$. Finally, since the relative degree of $(\text{Mat}_s(\mathbf{k}_D))^r$ over $\text{Mat}_s(\mathbf{k}_{E_D})$ is $e = [\Psi_{E/F} : \mu_E/\mu_F]$, a Frobenius reciprocity argument (which is still valid when p does not divide $\#\Psi_{E/F}$) implies that

$$(\text{Mat}_s(\mathbf{k}_D)^r)_{j'} \cong \text{Ind}_{\mu_E/\mu_F}^{\Psi_{E/F}} \text{Mat}_s(\mathbf{k}_{E_D})_{j'};$$

Therefore, we have obtained the desired decomposition and proved the proposition. □

The following Corollary is a direct consequence of Proposition 4.2.

COROLLARY 4.3. *The graded algebra*

$$\mathfrak{U}_{\mathfrak{A}} := \bigoplus_{j'=0}^{f/s-1} (\text{Mat}_s(\mathbf{k}_D)^r)_{j'},$$

is isomorphic to $\mathfrak{U}_{\mathfrak{A}^}$ as a $\Psi_{E/F}$ -module.* □

We provide some notations for later use. We write

$$\mathfrak{U}_{\text{sym}} := \bigoplus_{[g] \in (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}}} \mathfrak{U}_{[g]}$$

and also $\mathfrak{U}_{\text{sym-ram}}$ and $\mathfrak{U}_{\text{sym-unram}}$ analogously. Given intermediate extensions $F \subseteq K \subseteq L \subseteq E$, we write

$$\mathfrak{U}_{K/L} := \bigoplus_{[g] \in \Gamma_E \backslash (\Gamma_L - \Gamma_K) / \Gamma_E} \mathfrak{U}_{[g]}$$

We also define the *symmetric module* associated to $\mathfrak{U}_{[g]}$ (or $\mathfrak{U}_{[g^{-1}]}$) by

$$\mathfrak{U}_{[g]} := \begin{cases} \mathfrak{U}_{[g]} \oplus \mathfrak{U}_{[g^{-1}]} & \text{if } [g] \text{ is asymmetric,} \\ \mathfrak{U}_{[g]} & \text{if } [g] \text{ is symmetric,} \end{cases} \tag{4.5}$$

and call

$$\mathfrak{U}_{\mathfrak{A}} \cong \mathfrak{U}_0 \bigoplus_{[g] \in (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}} \sqcup (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}} / \pm} \mathfrak{U}_{[g]} \tag{4.6}$$

the *complete symmetric decomposition* of $\mathfrak{U}_{\mathfrak{A}}$. If \mathfrak{V} is a submodule of $\mathfrak{U}_{\mathfrak{A}}$, we also use the same convention to denote its submodules, for example, $\mathfrak{V}_{K/L} = \mathfrak{U}_{K/L} \cap \mathfrak{V}$ and $\mathfrak{V}_{[g]} = \mathfrak{U}_{[g]} \cap \mathfrak{V}$, and also call

$$\mathfrak{V} \cong (\mathfrak{U}_0 \cap \mathfrak{V}) \bigoplus_{[g] \in (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}} \sqcup (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}} / \pm} \mathfrak{V}_{[g]}$$

the complete symmetric decomposition of \mathfrak{V}

4.2 COMPLETE DECOMPOSITION OF FINITE SYMPLECTIC MODULES

We are interested in the adjoint action of E^\times on ${}_A\mathfrak{V}_\xi$ restricted from that of \mathbf{J} , which factors through the finite group $E^\times / F^\times (E^\times \cap J^1) \cong \Psi_{E/F}$. We also know that this action preserves the symplectic structure h_θ (3.15) on ${}_A\mathfrak{V}_\xi$. Hence ${}_A\mathfrak{V}_\xi$ is moreover a finite symplectic $\mathbb{F}_p\Gamma$ -module for each cyclic subgroup Γ of $\Psi_{E/F}$. We denote the $\mathfrak{U}_{[g]}$ -isotypic component in ${}_A\mathfrak{V}_\xi$ by ${}_A\mathfrak{V}_{\xi,[g]}$, and obtain the decompositions

$${}_A\mathfrak{V}_\xi = \bigoplus_{[g] \in (\Gamma_E \backslash \Gamma_F / \Gamma_E)'} \bigoplus_{[g] \in (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{sym}} \sqcup (\Gamma_E \backslash \Gamma_F / \Gamma_E)_{\text{asym}} / \pm} {}_A\mathfrak{V}_{\xi,[g]} \tag{4.7}$$

inherited from (4.1) and (4.6) respectively. These decomposition are finer than the one in (3.16). Indeed, it is easy to see that

$${}_A\mathfrak{V}_{\xi,k} := {}_A\mathfrak{V}_{\xi,E_k/E_{k+1}} = \bigoplus_{[g] \in \Gamma_E \backslash (\Gamma_{E_{k+1}} - \Gamma_{E_k}) / \Gamma_E} {}_A\mathfrak{V}_{\xi,[g]}$$

for $k = 0, \dots, t$.

PROPOSITION 4.4. *The complete symmetric decomposition of ${}_A\mathfrak{V}_\xi$ is orthogonal with respect to the alternating form h_θ .*

Proof. Since we know that the $\Psi_{E/F}$ -components of ${}_A\mathfrak{V}_\xi$ consist of those in the standard module $\mathfrak{U}_{\mathfrak{A}}$, which is isomorphic to the standard one $\mathfrak{U}_{\mathfrak{A}^*}$ in the split case, the proof of the assertion is just analogous to the one in the split case [Tam, (5.10)], based on the argument of [BF83, (8.2.3),(8.2.4)]. \square

We would like to describe the isotypic component appearing in the complete decomposition (4.7) of ${}_A\mathfrak{V}_\xi$. We first write $e(\mathfrak{A}/\mathfrak{A}_{k+1}) := e(\mathfrak{A}/\mathfrak{o}_E)/e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)$.

PROPOSITION 4.5. *The quotient $\mathfrak{P}_{\mathfrak{A}}^j \cap A_{k+1} / \mathfrak{P}_{\mathfrak{A}}^{j+1} \cap A_{k+1}$ is non-trivial if and only if $j \in e(\mathfrak{A}/\mathfrak{A}_{k+1})\mathbb{Z}$.*

Proof. Since $\mathfrak{P}_{\mathfrak{A}}^j \cap A_{k+1} = \mathfrak{P}_{\mathfrak{A}_{k+1}}^{j/e(\mathfrak{A}/\mathfrak{A}_{k+1})}$ for all $j \in \mathbb{Z}$, the assertion follows directly. \square

We now specify $j = j_k = e(\mathfrak{A}/\mathfrak{o}_E)a_k/2$ for some integer a_k , and so $\mathfrak{P}_{\mathfrak{A}}^{j_k} \cap A_{k+1} = \mathfrak{P}_{\mathfrak{A}_{k+1}}^{a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2}$, such that the index on the right side is the one appearing in the group J^1 (3.10). The condition in Proposition 4.5 is satisfied if and only if a_k is even or $e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)$ is even, in which case

$$\begin{aligned} {}_A\mathfrak{V}_{\xi,k} &\cong \frac{\mathfrak{P}_{\mathfrak{A}_{k+1}}^{a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2}}{\mathfrak{P}_{\mathfrak{A}_{k+1}}^{a_k e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)/2+} + \mathfrak{P}_{\mathfrak{A}_k}^{a_k e(\mathfrak{A}_k/\mathfrak{o}_E)/2}} \\ &\cong \begin{cases} \text{Mat}_{s_{k+1}}(\mathbf{k}_{D_{k+1}})^{r_{k+1}} / \text{Mat}_{s_k}(\mathbf{k}_{D_k})^{r_k} & \text{when } a_k e(\mathfrak{A}_k/\mathfrak{o}_E)/2 \in \mathbb{Z}, \\ \text{Mat}_{s_{k+1}}(\mathbf{k}_{D_{k+1}})^{r_{k+1}} & \text{otherwise,} \end{cases} \end{aligned}$$

where r_k and s_k are the invariants of \mathfrak{A}_k analogous to r and s of \mathfrak{A} .

To summarize, ${}_A\mathfrak{V}_{\xi,k}$ is isomorphic to

$$\begin{aligned} &0 \quad \text{when } a_k \text{ is odd and } e(\mathfrak{A}_{k+1}/\mathfrak{o}_E) \text{ is odd,} \\ &\text{Mat}_{s_{k+1}}(\mathbf{k}_{D_{k+1}})^{r_{k+1}} \\ &\quad \text{when } a_k \text{ is odd, } e(\mathfrak{A}_k/\mathfrak{o}_E) \text{ is odd, and } e(\mathfrak{A}_{k+1}/\mathfrak{o}_E) \text{ is even,} \\ &\text{Mat}_{s_{k+1}}(\mathbf{k}_{D_{k+1}})^{r_{k+1}} / \text{Mat}_{s_k}(\mathbf{k}_{D_k})^{r_k} \\ &\quad \text{when } a_k \text{ is even or } e(\mathfrak{A}_k/\mathfrak{o}_E) \text{ is even.} \end{aligned}$$

The action of $\Psi_{E/F}$ on $\mathfrak{P}_{\mathfrak{A}}^{j_k} \cap A_{k+1}$ is given by $\sigma^i \phi^{jh} \in \Gamma_{E_{k+1}}$, where j has image hj_k in the natural projection $\mathbb{Z}/f\mathbb{Z} \rightarrow \mathbb{Z}/e(\mathfrak{A}/\mathfrak{o}_E)\mathbb{Z}$. Therefore, directly from the description of ${}_A\mathfrak{V}_{\xi,k}$ above, we have the following decompositions.

PROPOSITION 4.6. *The complete decomposition of the component ${}_A\mathfrak{V}_{\xi,k}$ is given as follows.*

- (i) *When a_k is odd and $e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)$ is odd, then ${}_A\mathfrak{V}_{\xi,k}$ is trivial.*
- (ii) *When a_k is odd, $e(\mathfrak{A}_k/\mathfrak{o}_E)$ is odd, and $e(\mathfrak{A}_{k+1}/\mathfrak{o}_E)$ is even, then*

$${}_A\mathfrak{V}_{\xi,k} \cong \bigoplus_{\substack{[g]=[\sigma^i \phi^j] \in \Gamma_E \setminus \Gamma_{E_{k+1}} / \Gamma_E \\ j \equiv hj_k \pmod{e(\mathfrak{A}/\mathfrak{o}_E)}}} \mathfrak{U}_{[g]}.$$

- (iii) *When a_k is even or $e(\mathfrak{A}_k/\mathfrak{o}_E)$ is even, then*

$${}_A\mathfrak{V}_{\xi,k} \cong \bigoplus_{\substack{[g]=[\sigma^i \phi^j] \in \Gamma_E \setminus (\Gamma_{E_{k+1}} - \Gamma_{E_k}) / \Gamma_E \\ j \equiv hj_k \pmod{e(\mathfrak{A}/\mathfrak{o}_E)}}} \mathfrak{U}_{[g]}.$$

\square

4.3 SOME PROPERTIES OF PARITIES OF JUMPS

Let R be the index when $f(E/E_R)$ is odd and $f(E/E_{R+1})$ is even.

LEMMA 4.7. *We have $(\Gamma_{E_{R+1}}/\Gamma_E)_{\text{sym-unram}} = ((\Gamma_{E_{R+1}} - \Gamma_{E_R})/\Gamma_E)_{\text{sym-unram}}$.*

Proof. Recall Proposition 2.2 that every symmetric unramified $[g]$ are of the form $[\sigma^i \phi^{f/2}]$, so there is no coset of the form $\sigma^i \phi^{f/2}$ belonging to Γ_{E_R} . \square

Let Q be the index when $e(\mathfrak{A}_Q/\mathfrak{o}_E)$ is odd and $e(\mathfrak{A}_{Q+1}/\mathfrak{o}_E)$ is even.

LEMMA 4.8. *Suppose that f is even. We always have $R \leq Q$. If moreover m is odd, then $Q = R$.*

Proof. We know that $e(\mathfrak{A}_k/\mathfrak{o}_E)$ divides $f(E/E_k)$, so that if $Q < R$, then the even number $e(\mathfrak{A}_{Q+1}/\mathfrak{o}_E)$ divides $f(E/E_{Q+1})$, which divides the odd number $f(E/E_R)$. This is a contradiction. Hence $R \leq Q$. When $R \leq Q$, then

$$e(\mathfrak{A}_{R+1}/\mathfrak{o}_E) \text{ is odd and } f(E/E_{R+1}) \text{ is even.} \tag{4.8}$$

Since

$$e(\mathfrak{A}_{R+1}/\mathfrak{o}_E) = f_{E_{R+1}}/s_{E_{R+1}} = \frac{f(E/E_{R+1})}{\gcd(f(E/E_{R+1}), \gcd(m, n(E/E_{R+1})))}, \tag{4.9}$$

the statement (4.8) is equivalent to saying that

$$\begin{aligned} &\text{the 2-powers of the numerator and} \\ &\text{the denominator on the right side of (4.9) are equal.} \end{aligned} \tag{4.10}$$

This power is greater than 0. Hence (4.10) is equivalent to that

$$(\text{the 2-power of } m) \geq (\text{the 2-power of } f(E/E_{R+1})) \geq 0. \tag{4.11}$$

If m is odd, then (4.11) is a contradiction. \square

4.4 SYMMETRIC SUBMODULES

We write $A\mathfrak{A}_{\xi, \text{sym}} = A\mathfrak{A}_{\xi} \cap \mathfrak{A}_{\text{sym}}$ and $A\mathfrak{A}_{\xi, \text{sym-ram}}$ and $A\mathfrak{A}_{\xi, \text{sym-unram}}$ analogously.

4.4.1 CASE WHEN f IS ODD

From Proposition 3.2, we always have

$$A\mathfrak{A}_{\xi, \text{sym}} = A\mathfrak{A}_{\xi, \text{sym-ram}} \cong A^* \mathfrak{A}_{\xi, \text{sym-ram}} = A^* \mathfrak{A}_{\xi, \text{sym}}. \tag{4.12}$$

4.4.2 CASE WHEN f IS EVEN

Notice that the natural projection $\mathbb{Z}/f\mathbb{Z} \rightarrow \mathbb{Z}/e(\mathfrak{A}/\mathfrak{o}_E)\mathbb{Z}$ maps

$$f/2 \mapsto \begin{cases} 0 & \text{if } e(\mathfrak{A}/\mathfrak{o}_E) \text{ divides } f/2, \\ e(\mathfrak{A}/\mathfrak{o}_E)/2 \neq 0 & \text{otherwise} \end{cases} \quad (4.13)$$

The condition that $e(\mathfrak{A}/\mathfrak{o}_E)$ divides $f/2$ is equivalent to that s is even. When f is even, then $s = \gcd(f, m)$ is even if and only if m is even. We hence separate the cases according to the parity of m .

4.4.3 CASE WHEN BOTH f AND m ARE EVEN

In this case, $f/2$ is mapped to 0 by $\mathbb{Z}/f\mathbb{Z} \rightarrow \mathbb{Z}/e(\mathfrak{A}/\mathfrak{o}_E)\mathbb{Z}$. We separate the cases according to the parity of the jump a_k . When a_k is odd, neither 0 or $f/2$ is mapped to $hj_k \neq 0 \in \mathbb{Z}/e(\mathfrak{A}/\mathfrak{o}_E)\mathbb{Z}$, and so ${}^A\mathfrak{V}_{\xi,k}$ is trivial. When a_k is even, both 0 and $f/2$ are mapped to $hj_k = 0$ by (4.13), and so ${}^A\mathfrak{V}_{\xi,k,\text{sym}} \cong \mathfrak{U}_{k,\text{sym}}$. We also recall that

$${}^A\mathfrak{V}_{\xi,k} \cong \begin{cases} 0 & \text{if } a_k \text{ is odd,} \\ \mathfrak{U}_{k,\text{sym}} & \text{if } a_k \text{ is even.} \end{cases}$$

Whatever the parity of a_k is, we always have ${}^A\mathfrak{V}_{\xi,\text{sym}} \cong {}^A\mathfrak{V}_{\xi,\text{sym}}$.

4.4.4 CASE WHEN f IS EVEN AND m IS ODD

In this case, notice that $e(\mathfrak{A}/\mathfrak{o}_E)$ must be even, and

$$j_k = e(\mathfrak{A}/\mathfrak{o}_E)a_k/2 \equiv \begin{cases} 0 & \text{if } a_k \text{ is even,} \\ e(\mathfrak{A}/\mathfrak{o}_E)/2 & \text{if } a_k \text{ is odd.} \end{cases} \pmod{e(\mathfrak{A}/\mathfrak{o}_E)}$$

Therefore,

$${}^A\mathfrak{V}_{\xi,k,\text{sym}} = \begin{cases} {}^A\mathfrak{V}_{\xi,k,\text{sym-ram}} & \text{if } a_k \text{ is even,} \\ {}^A\mathfrak{V}_{\xi,k,\text{sym-unram}} & \text{if } a_k \text{ is odd.} \end{cases}$$

Using Proposition 4.6, we find that when a_k is even,

$${}^A\mathfrak{V}_{\xi,k,\text{sym}} = \mathfrak{U}_{E_k/E_{k+1},\text{sym-ram}} = \bigoplus_{[\sigma^i] \in (\Gamma_E \setminus (\Gamma_{E_{R+1}} - \Gamma_{E_R}) / \Gamma_E)_{\text{sym}}} \mathfrak{U}_{[\sigma^i]},$$

Here the index R is defined in Section 4.3. When a_k is odd, ${}^A\mathfrak{V}_{\xi,k,\text{sym}}$ is trivial when $k < R$, and is isomorphic to

$$\mathfrak{U}_{E/E_{R+1},\text{sym-unram}} = \bigoplus_{[\sigma^i \phi^{f/2}] \in (\Gamma_E \setminus \Gamma_{E_R} / \Gamma_E)_{\text{sym}}} \mathfrak{U}_{[\sigma^i \phi^{f/2}]}$$

when $k = R$, and to

$$\mathfrak{U}_{E_k/E_{k+1}, \text{sym-unram}} = \bigoplus_{[\sigma^i \phi^{f/2}] \in (\Gamma_E \backslash (\Gamma_{E_{k+1}} - \Gamma_{E_k}) / \Gamma_E)_{\text{sym}}} \mathfrak{U}_{[\sigma^i \phi^{f/2}]}$$

when $k > R$.

We observe that, whether a_k is odd or even, the symmetric unramified part of ${}_A \mathfrak{V}_\xi$ and ${}_{A^*} \mathfrak{V}_\xi$ are complementary, in the sense that

$${}_A \mathfrak{V}_{\xi, k, \text{sym-unram}} \oplus {}_{A^*} \mathfrak{V}_{\xi, k, \text{sym-unram}} = \mathfrak{U}_{E_k/E_{k+1}, \text{sym-unram}}$$

for all $k = 0, \dots, t$

We summarize the above in the following.

PROPOSITION 4.9. *We always have ${}_A \mathfrak{V}_{\xi, \text{sym-ram}} \cong {}_{A^*} \mathfrak{V}_{\xi, \text{sym-ram}}$ and*

- (i) *when f is odd, or when both f and m are even, then ${}_A \mathfrak{V}_{\xi, \text{sym-unram}} \cong {}_{A^*} \mathfrak{V}_{\xi, \text{sym-unram}}$;*
- (ii) *when f is even and m is odd, then ${}_A \mathfrak{V}_{\xi, \text{sym-unram}} \oplus {}_{A^*} \mathfrak{V}_{\xi, \text{sym-unram}} = \mathfrak{U}_{\text{sym-unram}}$.*

4.5 T-FACTORS OF ISOTYPIC COMPONENTS

We recall the values of the t-factors $t_\Gamma^i(\mathfrak{V})$, $i = 0, 1$, when Γ is one of the cyclic subgroups μ and $\langle \varpi \rangle$ of $\Psi_{E/F}$ defined in (3.20), and \mathfrak{V} is a symmetric module $\mathfrak{U}_{[g]}$ defined in (4.5). The following Proposition describes all $t_\Gamma^i(\mathfrak{U}_{[g]})$ except when $[g] = [\sigma^{e/2}]$.

PROPOSITION 4.10 ([Tam, Proposition 4.9]). (i) *If $[g] = [\sigma^i \phi^j]$ is asymmetric, then*

$$t_\mu^0(\mathfrak{U}_{[g]}) = 1, \quad t_\mu^1(\mathfrak{U}_{[g]}) : z \mapsto \text{sgn}_{z^{q^i-1}}(\mathfrak{U}_{[g]}),$$

$$t_{\langle \varpi \rangle}^0(\mathfrak{U}_{[g]}) = 1, \quad \text{and } t_{\langle \varpi \rangle}^1(\mathfrak{U}_{[g]}) (\varpi_E) = \text{sgn}_{z_e^i z_{\phi^j}}(\mathfrak{U}_{[g]}).$$

(ii) *If $[g] = [\sigma^i]$ is symmetric and not equal to $[1]$ or $[\sigma^{e/2}]$, then*

$$t_\mu^0(\mathfrak{U}_{[g]}) = 1, \quad t_\mu^1(\mathfrak{U}_{[g]}) \equiv 1,$$

$$t_{\langle \varpi \rangle}^0(\mathfrak{U}_{[g]}) = -1, \quad \text{and } t_{\langle \varpi \rangle}^1(\mathfrak{U}_{[g]}) : \varpi_E \mapsto \left(\frac{z_e^i}{\ker(N_{\mathbb{F}_p[z_e^i]/\mathbb{F}_p[z_e^i]_\pm})} \right).$$

(iii) *If $[g] = [\sigma^i \phi^{f/2}]$ is symmetric, then*

$$t_\mu^0(\mathfrak{U}_{[g]}) = -1, \quad t_\mu^1(\mathfrak{U}_{[g]}) \text{ is quadratic,}$$

and

- (I) if $z_e^i z_{\phi^{f/2}} = 1$, then $t_{\langle \varpi \rangle}^0(\mathfrak{U}_{[g]}) = 1$ and $t_{\langle \varpi \rangle}^1(\mathfrak{U}_{[g]}) \equiv 1$;
- (II) if $z_e^i z_{\phi^{f/2}} = -1$, then $t_{\langle \varpi \rangle}^0(\mathfrak{U}_{[g]}) = 1$ and $t_{\langle \varpi \rangle}^1(\mathfrak{U}_{[g]})(\varpi_E) = (-1)^{\frac{1}{2}(q^{f/2}-1)}$;
- (III) if $z_e^i z_{\phi^{f/2}} \neq \pm 1$, then $t_{\langle \varpi \rangle}^0(\mathfrak{U}_{[g]}) = -1$ and

$$t_{\langle \varpi \rangle}^1(\mathfrak{U}_{[g]}) : \varpi_E \mapsto \left(\frac{z_e^i z_{\phi^{f/2}}}{\ker(N_{\mathbb{F}_p[z_e^i z_{\phi^{f/2}]} / \mathbb{F}_p[z_e^i z_{\phi^{f/2}]_{\pm}}})} \right).$$

□

In the exceptional case, when $[g] = [\sigma^{e/2}]$, we have $\mu_{E_g} = \mu_E$. To unify notation, we define

$$t_{\mu}^1(\mathfrak{U}_{[\sigma^{e/2}]}): \mu_E \rightarrow \{\pm 1\}, x \mapsto \left(\frac{x}{\mu_E} \right). \tag{4.14}$$

The $\mathbb{F}_p\langle \varpi \rangle$ -module structure of $\mathfrak{U}_{[\sigma^{e/2}]}$ does not concern us (see the explanation after Formula (5.5)).

The following properties concerning symmetric double cosets are useful when computing the above t-factors.

PROPOSITION 4.11. *Suppose that $[g]$ is symmetric.*

- (i) *If $[g]$ is ramified (resp. unramified), then $[\mathfrak{U}_{[g]} : \mathbf{k}_E]$ is even (resp. odd).*
- (ii) *Let $\mathbb{F}_p[[\frac{1}{g}]](\varpi_E)$ be the field extension of \mathbb{F}_p generated by the image of $[\frac{1}{g}](\varpi_E)$ in \mathbf{k}_F^\times . If $[g] \neq [\sigma^{e/2}]$, then the degree $[\mathfrak{U}_{[g]} : \mathbb{F}_p[[\frac{1}{g}]](\varpi_E)]$ is odd.*

Proof. The first statement for ramified $[g]$ is a simple calculation, and that for unramified $[g]$ is a consequence of Proposition 2.3. The second statement is proved in [Tam, Lemma 4.8]. □

We would like to extend our definition of the t-factors $t_{\mu}^i(\mathfrak{U}_{[g]})$, with $i = 0, 1$, from μ to $\mu_g = \mu_{E_g} / \mu_F$. We define

$$t_{\mu_g}^0(\mathfrak{U}_{[g]}) := t_{\mu}^0(\mathfrak{U}_{[g]})$$

and for all $z \in \mu_g$,

$$t_{\mu_g}^1(\mathfrak{U}_{[g]}) : z \mapsto \begin{cases} \text{sgn}_{[\frac{1}{g}]}(z)(\mathfrak{U}_{[g]}) & \text{if } [g] \text{ is asymmetric,} \\ \left(\frac{[\sigma^i \phi^{f[\mathfrak{U}_{[g]} : \mathbf{k}_E] / 2}]}{\ker N_{\mathbf{k}_{E_g} / \mathbf{k}_{E_{\pm g}}}} \right)(z) & \text{if } [g] \text{ is symmetric.} \end{cases}$$

PROPOSITION 4.12. *The restriction $t_{\mu_g}^1(\mathfrak{U}_{[g]})$ to μ is $t_{\mu}^1(\mathfrak{U}_{[g]})$.*

Proof. For asymmetric $[g]$, the result is immediate by definition. For symmetric ramified $[g]$, the restriction of the root $\left[\sigma^i \phi^{f[\mathfrak{u}_{[g]}] : \mathfrak{k}_E} \right]$ to $\mu = \mu_E / \mu_F$ is trivial, so the assertion is again true. When $[g]$ is symmetric unramified, we have to show that the restriction of $\left(\frac{\cdot}{\ker N_{\mathfrak{k}_{E_g} / \mathfrak{k}_{E \pm g}}} \right)$ to μ is $\left(\frac{\cdot}{\ker N_{\mathfrak{k}_E / \mathfrak{k}_{E \pm}}} \right)$, or equivalently, to show that the index of the subgroup $\ker N_{\mathfrak{k}_E / \mathfrak{k}_{E \pm}} \cong \mu_{q^{f/2+1}}$ of $\ker N_{\mathfrak{k}_E / \mathfrak{k}_{E \pm}} \cong \mu_{q^{f[\mathfrak{u}_{[g]}] : \mathfrak{k}_E} / 2 + 1}$ is odd, which follows from Proposition 4.11. \square

5 ZETA-DATA

5.1 ADMISSIBLE EMBEDDINGS OF L-TORI

As mentioned in Section 1.3, to understand ζ -data, it is better to first understand χ -data, which is motivated by constructing admissible embeddings of L-tori [LS87, Section 2.6].

We take T to be an elliptic torus of G isomorphic to $\text{Res}_{E/F} \mathbb{G}_m$. Its dual torus \hat{T} is $\text{Ind}_{E/F}(\mathbb{C}^\times)$, which is isomorphic to $(\mathbb{C}^\times)^n$ as a group. It is equipped with the induced action of the Weil group \mathcal{W}_F , which factors through the action of the Galois group Γ_F . We define the L-torus ${}^L T := \hat{T} \rtimes \mathcal{W}_F$ as the L-group of T .

We assume that the dual torus \hat{T} is embedded into the L-group ${}^L G = \hat{G} \rtimes \mathcal{W}_F$ of G , where $\hat{G} = \text{GL}_n(\mathbb{C})$, with image \mathcal{T} . For convenience, we simply denote the image of $t \in \hat{T}$ by the embedding $\hat{T} \rightarrow \mathcal{T} \subset \hat{G}$ also by $t \in \mathcal{T}$. This embedding should be defined using the chosen splittings of G and \hat{G} . As we do not need the full detail of the definition of this embedding, we only refer to [LS87, Section 2.5] for details (or, when $(G, T) = (\text{GL}_n, \text{Res}_{E/F} \mathbb{G}_m)$, see [Tam, Section 6.1]). All we need to know is that we can always assume that the image \mathcal{T} is the diagonal subgroup of \hat{G} .

With the embedding $\hat{T} \rightarrow \mathcal{T}$ chosen, an *admissible embedding* from ${}^L T$ to ${}^L G$ is a morphism of groups $I : {}^L T \rightarrow {}^L G$ of the form

$$I(t \rtimes w) = tI(1 \rtimes w) \text{ for all } t \rtimes w \in {}^L T.$$

Note that an admissible embedding maps \mathcal{W}_F into $N_{\hat{G}}(\mathcal{T})$, i.e., the factor $I(1 \rtimes w)$ above lies in $N_{\hat{G}}(\mathcal{T})$. Two admissible embeddings I_1, I_2 are called $\text{Int}(\mathcal{T})$ -equivalent if there is $t \in \mathcal{T}$ such that

$$I_1(w) = tI_2(w)t^{-1} \text{ for all } w \in \mathcal{W}_F.$$

By [LS87, Section 2.6], admissible embeddings exist, and the collection of these embeddings can be described as follows.

PROPOSITION 5.1. *The set of admissible embeddings from ${}^L T$ to ${}^L G$ is a $Z^1(\mathcal{W}_F, \hat{T})$ -torsor, and the set of the $\text{Int}(\mathcal{T})$ -equivalence classes of these embeddings is an $H^1(\mathcal{W}_F, \hat{T})$ -torsor.*

The idea in [LS87, Section 2.5] of constructing an admissible embedding is to choose a set of characters

$$\{\chi_\lambda\}_{\lambda \in \Phi}, \text{ where } \chi_\lambda : E_\lambda^\times \rightarrow \mathbb{C}^\times,$$

called χ -data, such that the following conditions hold.

DEFINITION 5.2. (i) For each $\lambda \in \Phi$, we have $\chi_{-\lambda} = \chi_\lambda^{-1}$ and $\chi^{w\lambda} = {}^w\chi_\lambda$ for all $w \in \mathcal{W}_F$.

(ii) If λ is symmetric, then $\chi|_{E_{\pm\lambda}^\times}$ equals the quadratic character $\delta_{E_\lambda/E_{\pm\lambda}}$ attached to the extension $E_\lambda/E_{\pm\lambda}$.

Remember that, in Section 2.1, we choose a subset $\mathcal{R}_\pm = \mathcal{R}_{\text{sym}} \sqcup \mathcal{R}_{\text{asym}/\pm}$ of Φ representing the orbits $\mathcal{W}_F \backslash \Phi_{\text{sym}}$ and $\mathcal{W}_F \backslash \Phi_{\text{asym}/\pm}$. Hence, by condition (i), the set of χ -data depends completely on the subset $\{\chi_\lambda\}_{\lambda \in \mathcal{R}_\pm}$. We still call such a subset a set of χ -data. Moreover, using Artin reciprocity [Tat79], we may regard each χ_λ as a character of the Weil group \mathcal{W}_{E_λ} .

Following the recipe in [LS87, Section 2.5], we can define an admissible embedding

$$I_{\{\chi_\lambda\}} : {}^L T \rightarrow {}^L G$$

depending on a given set of χ -data. In our present situation, we can describe the admissible embedding $I_{\{\chi_\lambda\}}$ in Proposition 5.3 below. We first recall the Langlands correspondence for the torus $T = \text{Res}_{E/F} \mathbb{G}_m$, which is a bijection

$$\text{Hom}(T(F), \mathbb{C}^\times) \rightarrow H^1(\mathcal{W}_F, \hat{T}). \tag{5.1}$$

Given a character ξ of $T(F) = E^\times$, we denote by $\tilde{\xi}$ a 1-cocycle in $Z^1(\mathcal{W}_F, \hat{T})$ whose class is the image of ξ under (5.1). Given χ -data $\{\chi_\lambda\}_{\lambda \in \mathcal{R}_\pm}$, we define

$$\mu := \mu_{\{\chi_\lambda\}} = \prod_{\lambda \in \mathcal{R}} \text{Res}_{E_\lambda^\times}^{E^\times} \chi_\lambda,$$

where $\mathcal{R} = \mathcal{R}_\pm \sqcup (-\mathcal{R}_{\text{asym}/\pm})$ is a subset representing $\Gamma_F \backslash \Phi$. It is easy to check that the product of the restricted characters is independent of representatives in \mathcal{R} , so we usually write

$$\mu = \prod_{[\lambda] \in \Gamma_F \backslash \Phi} \text{Res}_{E_\lambda^\times}^{E^\times} \chi_\lambda.$$

PROPOSITION 5.3 ([Tam, Proposition 6.5]). *For every character ξ of E^\times , the composition*

$$I_{\{\chi_\lambda\}} \circ \tilde{\xi} : \mathcal{W}_F \rightarrow {}^L T \rightarrow {}^L G \xrightarrow{\text{proj}} \text{GL}_n(\mathbb{C})$$

is isomorphic to $\text{Ind}_{E/F}(\xi \cdot \mu_{\{\chi_\lambda\}})$ as a representation of \mathcal{W}_F . □

We now define an analogous set of characters

$$\{\zeta_\lambda\}_{\lambda \in \Phi}, \text{ where } \zeta_\lambda : E_\lambda^\times \rightarrow \mathbb{C}^\times,$$

called ζ -data, such that the following conditions hold.

DEFINITION 5.4. (i) For each $\lambda \in \Phi$, we have $\zeta_{-\lambda} = \zeta_\lambda^{-1}$ and $\zeta_{w\lambda} = {}^w\zeta_\lambda$ for all $w \in \mathcal{W}_F$.

(ii) If λ is symmetric, then $\zeta|_{E_{\pm\lambda}^\times}$ is trivial.

We can view a set of ζ -data as the difference of two sets of χ -data. Motivated from Propositions 5.1 and 5.3, the product character

$$\nu := \nu_{\{\zeta_\lambda\}} = \prod_{[\lambda] \in \Gamma_F \backslash \Phi} \text{Res}_{E_\lambda^\times}^{E_\lambda^\times} \zeta_\lambda.$$

can be viewed as measuring the difference of two admissible embeddings. Recall that, similar to choosing \mathcal{R}_\pm , we can also choose $\mathcal{D}_\pm = \mathcal{D}_{\text{sym}} \sqcup \mathcal{D}_{\text{asym}/\pm}$ to be a subset of Γ_F/Γ_E consisting of representatives of $(\Gamma_E \backslash \Gamma_F/\Gamma_E)_{\text{sym}}$ and $(\Gamma_E \backslash \Gamma_F/\Gamma_E)_{\text{asym}/\pm}$ respectively, and obtain a bijection from Proposition 2.1,

$$\mathcal{R}_\pm = \mathcal{R}_{\text{sym}} \bigsqcup \mathcal{R}_{\text{asym}/\pm} \rightarrow \mathcal{D}_\pm = \mathcal{D}_{\text{sym}} \bigsqcup \mathcal{D}_{\text{asym}/\pm}, \lambda = \left[\frac{1}{g} \right] \mapsto g.$$

We usually denote by E_g and $E_{\pm g}$ the fields E_λ and $E_{\pm\lambda}$ respectively, if $g \in \mathcal{D}_\pm$ corresponds to $\lambda \in \mathcal{R}_\pm$. We also denote by ζ_g the character ζ_λ , and write

$$\nu := \nu_{\{\zeta_g\}} = \prod_{[g] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)'} \text{Res}_{E_g^\times}^{E_g^\times} \zeta_g.$$

5.2 SYMMETRIC UNRAMIFIED ZETA-DATA

We choose a specific ζ -data ζ_g for each $[g] = [\sigma^i \phi^{f/2}] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)_{\text{sym-unram}}$, base on the results from the χ -datum χ_g .

Notice that, since $E_g/E_{\pm g}$ is quadratic unramified, the norm group $N_{E_g/E_{\pm g}}(E_g^\times)$ has a decomposition

$$\mu_{E_{\pm g}} \times \langle z_e^i z_{\phi^{f/2}} \varpi_E^2 \rangle \times U_{E_{\pm g}}^1$$

and we take a root of unity $z_0 \in \mu_{E_g}$ such that

$$z_0 \varpi_E \in E_{\pm g}^\times - N_{E_g/E_{\pm g}}(E_g^\times).$$

We only consider tamely ramified χ -data and ζ -data, i.e., we require that

$$\chi_g|_{U_{E_g}^1} \equiv 1 \text{ and } \zeta_g|_{U_{E_g}^1} \equiv 1.$$

Therefore, the Definition 5.2.(ii) of χ -data is explicitly (see [Tam, (7.6)])

$$\chi_g(\mu_{E_{\pm g}}) = 1, \chi_g(z_e^i z_{\phi^{f/2}} \varpi_E^2) = 1, \text{ and } \chi_g(z_0 \varpi_E) = -1. \tag{5.2}$$

Hence, given a χ -datum χ_g , we can obtain a ζ -datum ζ_g easily by requiring

$$\zeta_g|_{\mu_{E_g}} = \chi_g|_{\mu_{E_g}} \text{ and } \zeta_g(\varpi_E) = -\chi_g(\varpi_E).$$

In [Tam, Section 7.4], in the cases when $\mathfrak{Y}_{\xi, [g]} \cong \mathfrak{U}_{[g]}$ is non-trivial, we construct a χ -datum

$$\chi_g|_{\mu_{E_g}} = t_{\mu_g}^1(\mathfrak{U}_{[g]}) \text{ and } \chi_g(\varpi_E) = \begin{cases} -t_{\langle \varpi \rangle}(\mathfrak{U}_{[g]}) & \text{if } \sigma^i \phi^{f/2} \varpi_E = \varpi_E, \\ t_{\langle \varpi \rangle}(\mathfrak{U}_{[g]}) & \text{otherwise.} \end{cases}$$

In other words, the character χ_g satisfies the conditions in (5.2). Hence the character

$$\zeta_g|_{\mu_{E_g}} = t_{\mu_g}^1(\mathfrak{U}_{[g]}) \text{ and } \zeta_g(\varpi_E) = \begin{cases} t_{\langle \varpi \rangle}(\mathfrak{U}_{[g]}) & \text{if } \sigma^i \phi^{f/2} \varpi_E = \varpi_E, \\ -t_{\langle \varpi \rangle}(\mathfrak{U}_{[g]}) & \text{otherwise} \end{cases}$$

is a ζ -datum. This ζ -datum will be used in the next section.

5.3 ZETA-DATA ASSOCIATED TO ADMISSIBLE CHARACTERS

Given an admissible character ξ of E^\times over F , we first assign, for each $[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)_{\text{asym}/\pm}$, the values of the ζ -data to

$$\zeta_{g, \xi}|_{\mu_{E_g}} = \text{sgn}_{\mu_{E_g}}(A\mathfrak{Y}_{[g]}) \text{sgn}_{\mu_{E_g}}(A^*\mathfrak{Y}_{[g]}) = t_{\mu_g}^1(A\mathfrak{Y}_{\xi, [g]}) t_{\mu_g}^1(A^*\mathfrak{Y}_{\xi, [g]}).$$

In this way, the product of the characters

$$\zeta_{g, \xi} \zeta_{g^{-1}, \xi} = \zeta_{g, \xi} \left(\zeta_{g, \xi}^g \right)^{-1} = \zeta_{g, \xi} \circ \left[\frac{1}{g} \right]$$

has values

$$\begin{aligned} (\zeta_{g, \xi} \circ \left[\frac{1}{g} \right])|_{\mu_E}(z) &= \text{sgn}_{\mu_E}(A\mathfrak{Y}_{\xi, [g]})(\left[\frac{1}{g} \right](z)) \text{sgn}_{\mu_E}(A^*\mathfrak{Y}_{\xi, [g]})(\left[\frac{1}{g} \right](z)) \\ &= t_{\mu}^1(A\mathfrak{Y}_{\xi, [g]}) t_{\mu}^1(A^*\mathfrak{Y}_{\xi, [g]}) \end{aligned}$$

and

$$\begin{aligned} (\zeta_{g, \xi} \circ \left[\frac{1}{g} \right])(\varpi_E) &= \zeta_{g, \xi}|_{\mu_E} \left(\left[\frac{1}{g} \right](\varpi_E) \right) \\ &= \text{sgn}_{\left[\frac{1}{g} \right](\varpi_E)}(A\mathfrak{Y}_{\xi, [g]}) \text{sgn}_{\left[\frac{1}{g} \right](\varpi_E)}(A^*\mathfrak{Y}_{\xi, [g]}) \\ &= t_{\langle \varpi \rangle}^1(A\mathfrak{Y}_{\xi, [g]})(\varpi_E) t_{\langle \varpi \rangle}^1(A^*\mathfrak{Y}_{\xi, [g]})(\varpi_E) \\ &= t_{\langle \varpi \rangle}(A\mathfrak{Y}_{\xi, [g]}) t_{\langle \varpi \rangle}(A^*\mathfrak{Y}_{\xi, [g]}). \end{aligned} \tag{5.3}$$

We then assign, for each $[g] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)_{\text{asym}/\pm}$, arbitrary values to $\zeta_{g,\xi}(\varpi_E)$ and $\zeta_{g^{-1},\xi}(\varpi_E)$, as long as the product satisfies (5.3). (This phenomenon is comparable to [LS87, Lemma 3.3.A], as explained in [Tam, Remark 7.2].) It is routine to check that each $\zeta_{g,\xi}$ is a ζ -datum. Indeed, this checking is exactly the same as that in the χ -data case (see [Tam, Section 7.2]), since Definition 5.2.(i) is the same as that of χ -data.

We then assign values to the ζ -data for each $[g] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)_{\text{sym}}$ case by case.

5.3.1 CASE WHEN f IS ODD

Recall from (4.12) that

$$t_{\Gamma}^i(A\mathfrak{V}_{\xi,\text{sym}}) = t_{\Gamma}^i(A^*\mathfrak{V}_{\xi,\text{sym}}), \text{ for } i = 0, 1 \text{ and } \Gamma = \mu, \langle \varpi \rangle. \tag{5.4}$$

We assign the ζ -data to the following values. If e is odd (so that m is odd since m divides e), we assign all $\zeta_{g,\xi}$ to be trivial. If e is even, then we just take all $\zeta_{g,\xi}$, $[g] \neq [\sigma^{e/2}]$, to be trivial and

$$\zeta_{\sigma^{e/2},\xi}|_{\mu_E} \equiv 1 \text{ and } \zeta_{\sigma^{e/2},\xi}(\varpi_E) = (-1)^m.$$

To show that $\zeta_{\sigma^{e/2},\xi}$ is a ζ -datum, notice that since $N_{E/E_{\pm\sigma^{e/2}}}(\varpi_E) = -\varpi_{E_{\pm\sigma^{e/2}}} = -\varpi_E^2$, and since

$$\zeta_{\sigma^{e/2},\xi}(\varpi_E)^2 = \zeta_{\sigma^{e/2},\xi}(-1)\zeta_{\sigma^{e/2},\xi}(-\varpi_E^2) = (1)(1) = 1,$$

we can assign $\chi_{\sigma^{e/2},\xi}(\varpi_E)$ to either 1 or -1 to obtain a ζ -datum. By (5.4), we can rewrite our assigned ζ -data as

$$\begin{aligned} \zeta_{g,\xi}|_{\mu_E} &= t_{\mu}^1(A\mathfrak{V}_{\xi,[g]})t_{\mu}^1(A^*\mathfrak{V}_{\xi,[g]}) \\ \zeta_{g,\xi}(\varpi_E) &= \begin{cases} t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi,[g]}) & \text{if } g \neq \sigma^{e/2}, \\ (-1)^m t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi,[g]}) & \text{if } g = \sigma^{e/2}. \end{cases} \end{aligned} \tag{5.5}$$

Note that $t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[\sigma^{e/2}]})$ is not defined (see the paragraph containing Formula (4.14)). In fact, we just take

$$t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[\sigma^{e/2}]}) = t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi,[\sigma^{e/2}]}) = 1,$$

since $A\mathfrak{V}_{\xi,[\sigma^{e/2}]} \cong A^*\mathfrak{V}_{\xi,[\sigma^{e/2}]}$ by Proposition 3.2, and it is shown in [Tam, Proposition 5.3] that $A^*\mathfrak{V}_{\xi,[\sigma^{e/2}]}$ is always trivial.

The product of ζ -data is equal to

$$\prod_{[g] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}(\varpi_E) \equiv 1$$

and

$$\prod_{[g] \in (\mathcal{W}_E \backslash \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}(\varpi_E) = \begin{cases} (-1)^m & \text{if } e \text{ is even,} \\ 1 & \text{if } e \text{ is odd,} \end{cases}$$

which is rewritten as

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}|_{\mu_E} \equiv t_{\mu}^1(A\mathfrak{A}_{\xi})t_{\mu}^1(A^*\mathfrak{A}_{\xi})$$

and

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}(\varpi_E) = \begin{cases} (-1)^m t_{\langle \varpi \rangle}(A\mathfrak{A}_{\xi})t_{\langle \varpi \rangle}(A^*\mathfrak{A}_{\xi}) & \text{if } e \text{ is even,} \\ t_{\langle \varpi \rangle}(A\mathfrak{A}_{\xi})t_{\langle \varpi \rangle}(A^*\mathfrak{A}_{\xi}) & \text{if } e \text{ is odd.} \end{cases}$$

The product is equal to the rectifier given in (3.21),

$$D\nu_{\xi}|_{\mu_E} \equiv 1 \text{ and } D\nu_{\xi}(\varpi_E) = (-1)^{m(d-1)} = (-1)^{e-m},$$

when E/F is totally ramified.

5.3.2 CASE WHEN f IS EVEN

Let K be the maximal unramified extension of E/F . If we define $D_K\nu_{\xi}$ to be the ramified part of $D\nu_{\xi}$, which is also the rectifier corresponding to the admissible pair $(E/K, \xi)$, then we have

$$D_K\nu_{\xi} = \prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_K / \mathcal{W}_E)'} \zeta_{g,\xi}|_{E^{\times}},$$

as when f is odd, and in particular

$$D_K\nu_{\xi}(\varpi_E) = (-1)^{e-m_K},$$

where $m_K = \gcd(e, m)$. Therefore, our plan is to distribute the sign

$$(-1)^{e-m_K} (-1)^{n-m+f_{\varpi}-m_{\varpi}} = \begin{cases} 1 & \text{if } m \text{ is even,} \\ (-1)^{e+f_{\varpi}+1} & \text{if } m \text{ is odd,} \end{cases} \tag{5.6}$$

to each $\zeta_{g,\xi}(\varpi_E)$, where $[g]$ is symmetric unramified, multiplying the product of t -factors $t_{\langle \varpi \rangle}(A\mathfrak{A}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{A}_{\xi,[g]})$. As before, we separate the cases according to the parity of m .

When m is even, recall from Proposition 4.9 that we have either

$$\text{both } A\mathfrak{A}_{\xi,k,\text{sym}} \text{ and } A^*\mathfrak{A}_{\xi,k,\text{sym}} \text{ are trivial,}$$

or

$$\text{both } A\mathfrak{A}_{\xi,k,\text{sym}} \text{ and } A^*\mathfrak{A}_{\xi,k,\text{sym}} \text{ are isomorphic to } \mathfrak{U}_{k,\text{sym}}.$$

We assign the trivial ζ -data for all $[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)_{\text{sym-unram}}$, so that

$$\zeta_{g,\xi}|_{\mu_E} = t_{\mu}^1(A\mathfrak{A}_{\xi,[g]})t_{\mu}^1(A^*\mathfrak{A}_{\xi,[g]}) \text{ and } \zeta_{g,\xi}(\varpi_E) = t_{\langle \varpi \rangle}(A\mathfrak{A}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{A}_{\xi,[g]}).$$

The product of $\zeta_{g,\xi}(\varpi_E)$ is trivial, or we can write

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}|_{\mu_E} = t_{\mu}^1(A\mathfrak{V}_{\xi})t_{\mu}^1(A^*\mathfrak{V}_{\xi})$$

and

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}(\varpi_E) = t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi}).$$

Note that in the second product, the sign without t-factors is equal to (5.6), which is just 1.

When m is odd, we have

$$A\mathfrak{V}_{\xi, \text{sym-unram}} \oplus A^*\mathfrak{V}_{\xi, \text{sym-unram}} = \mathfrak{U}_{\text{sym-unram}}.$$

We then assign the ζ -data to be

$$\zeta_{g,\xi}|_{\mu_E} = t_{\mu}^1(A\mathfrak{V}_{\xi,[g]})t_{\mu}^1(A^*\mathfrak{V}_{\xi,[g]}) = t_{\mu}^1(\mathfrak{U}_{[g]})$$

and

$$\zeta_{g,\xi}(\varpi_E) = -t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi,[g]}) = -t_{\langle \varpi \rangle}(\mathfrak{V}_{[g]}).$$

for all symmetric unramified $[g]$ except the one which stabilizes ϖ_E , in which we assign

$$\zeta_{g,\xi}|_{\mu_E} = t_{\mu}^1(A\mathfrak{V}_{\xi,[g]})t_{\mu}^1(A^*\mathfrak{V}_{\xi,[g]}) = t_{\mu}^1(\mathfrak{U}_{[g]})$$

and

$$\zeta_{g,\xi}(\varpi_E) = t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi,[g]}) = t_{\langle \varpi \rangle}(\mathfrak{V}_{[g]}).$$

In Section 5.2, we checked that the above characters give rise to ζ -data. The product of ζ is hence

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}|_{\mu_E} = t_{\mu}^1(A\mathfrak{V}_{\xi})t_{\mu}^1(A^*\mathfrak{V}_{\xi})$$

and

$$\prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}(\varpi_E) = (-1)^{e+f_{\varpi}+1}t_{\langle \varpi \rangle}(A\mathfrak{V}_{\xi})t_{\langle \varpi \rangle}(A^*\mathfrak{V}_{\xi}),$$

by Proposition 2.6. Again in the second product, the sign without t-factors is equal to (5.6).

5.4 THE MAIN THEOREM

To summarize, we verified the following theorem.

THEOREM 5.5. *Let ξ be an admissible character of E^{\times} over F .*

(i) *Let $A\mathfrak{V}_{\xi}$ (resp. $A^*\mathfrak{V}_{\xi}$) be the finite symplectic module defined by ξ when $G(F) = \text{GL}_m(D)$ (resp. when $G^*(F) = \text{GL}_n(F)$). The following conditions define a collection of ζ -data $\{\zeta_{g,\xi}\}_{g \in \mathcal{D}_{\text{asym}}/\pm \sqcup \mathcal{D}_{\text{sym}}}$.*

(a) *All $\zeta_{g,\xi}$ are tamely ramified.*

(b) If $g \in \mathcal{D}_{\text{asym}/\pm}$, then

$$\zeta_{g,\xi}|_{\mu_{E_g}} = t_{\mu_g}^1(A\mathfrak{V}_{\xi,[g]})t_{\mu_g}^1(A^*\mathfrak{V}_{\xi,[g]})$$

and

$$\begin{aligned} \zeta_{g,\xi}(\varpi_E) &\text{ can be any value satisfying} \\ \zeta_{g,\xi}(\varpi_E)\zeta_{g^{-1},\xi}(\varpi_E) &= t_{\langle\varpi\rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle\varpi\rangle}(A^*\mathfrak{V}_{\xi,[g]}). \end{aligned}$$

(c) If $g \in \mathcal{D}_{\text{sym}}$, then

$$\zeta_{g,\xi}|_{\mu_{E_g}} = t_{\mu_g}^1(A\mathfrak{V}_{\xi,[g]})t_{\mu_g}^1(A^*\mathfrak{V}_{\xi,[g]})$$

and

$$\zeta_{g,\xi}(\varpi_E) = \epsilon_g t_{\langle\varpi\rangle}(A\mathfrak{V}_{\xi,[g]})t_{\langle\varpi\rangle}(A^*\mathfrak{V}_{\xi,[g]}),$$

where ϵ_g is equal to 1 if $g \in (\mathcal{D}_{\text{sym-ram}} - \{\sigma^{e/2}\}) \cup \mathcal{W}_{F[\varpi_E]}$ and is equal to $(-1)^m$ if $g \in (\mathcal{D}_{\text{sym-unram}} - \mathcal{W}_{F[\varpi_E]}) \cup \{\sigma^{e/2}\}$.

(ii) Let ${}_D\nu_\xi$ be the rectifier of ξ and $\{\zeta_{g,\xi}\}_{g \in \mathcal{D}_\pm}$ be the ζ -data in (i), then

$${}_D\nu_\xi = \prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_F / \mathcal{W}_E)'} \zeta_{g,\xi}|_{E^\times}.$$

□

REMARK 5.6. As long as the F -dimension of the division algebra D is fixed, the rectifier ${}_D\nu_\xi$ is independent of the Hasse-invariant $h = h(D)$ of D , as stated in [BH11, Theorem C]. This is because the modules $A\mathfrak{V}_\xi$, where $A = \text{Mat}_n(D)$ and D ranges over all division algebra with same F -dimension, are all isomorphic to each other. Similarly, the ζ -data $\{\zeta_{g,\xi}\}$ are independent of $h(D)$. □

5.5 FUNCTORIAL PROPERTY

Let K be an intermediate subfield in E/F , and write

$$n_K = n(E/K) = f_K e_K = f(E/K)e(E/K) \text{ and } m_K = \gcd(m, n_K).$$

Similar to Section 3.2, we have the Jacquet-Langlands correspondence

$$JL_K : \mathcal{A}_{n_K}^{\text{et}}(K) \rightarrow \mathcal{A}_{m_K}^{\text{et}}(D_K),$$

between essentially tame supercuspidal representations of $G(F)_K = \text{GL}_{m_K}(D_K)$ and its split inner form $G^*(F)_K = \text{GL}_{n_K}(K)$. We can parametrize both collections by the admissible pairs in $P_{n_K}(K)$, and obtain the rectifier map

$${}_{D_K}\nu : P_{n_K}(K) \xrightarrow{\kappa\Pi} \mathcal{A}_{n_K}^{\text{et}}(K) \xrightarrow{JL_K} \mathcal{A}_{m_K}^{\text{et}}(D_K) \xrightarrow{{}_{D_K}\Pi^{-1}} P_{n_K}(K),$$

such that

$${}_{D_K}\nu(E/K, \xi) = (E/F, \xi \cdot {}_{D_K}\nu\xi).$$

for a tamely ramified character ${}_{D_K}\nu\xi$ of E^\times for each pair $(E/K, \xi) \in P_{n_K}(K)$. With the embedding condition for (E_0K, \mathfrak{A}_K) as discussed in Section 2.5, we define the subgroups (see also [BH11, 3.2 Proposition])

$$H_K^1 = H^1(\Xi, \mathfrak{A}) \cap G(F)_K \text{ and } J_K^1 = J^1(\Xi, \mathfrak{A}) \cap G(F)_K$$

Each subgroup above admits a similar factorization as in (3.10). We then obtain

$${}_{A_K}\mathfrak{V}_\xi = J_K^1/H_K^1 = {}_A\mathfrak{V}_\xi \cap \mathfrak{U}_{E/K}$$

and similarly for ${}_{A_K^*}\mathfrak{V}_\xi$.

Denote $\Psi_{E/K} = E^\times/K^\times U_E^1$, and view ${}_{A_K}\mathfrak{V}_\xi$ and ${}_{A_K^*}\mathfrak{V}_\xi$ as $\mathbf{k}_K\Psi_{E/K}$ -submodules of $\mathfrak{U}_{E/K}$. Denote the subgroups of $\Psi_{E/K}$ by

$$\begin{aligned} \boldsymbol{\mu}_{E/K} &= \boldsymbol{\mu}_E/\boldsymbol{\mu}_K \text{ and} \\ \langle \varpi \rangle_{E/K} &= \text{the subgroup generated by the image of } \varpi_E. \end{aligned}$$

Using the results in Section 3.8, with the base field changed from F to K , the values of ${}_{D_K}\nu\xi$ is given by

$$\begin{aligned} {}_{D_K}\nu\xi|_{\boldsymbol{\mu}_E} &= t_{\boldsymbol{\mu}_{E/K}}^1({}_{A_K}\mathfrak{V}_\xi)t_{\boldsymbol{\mu}_{E/K}}^1({}_{A_K^*}\mathfrak{V}_\xi) \\ \text{and } {}_{D_K}\nu\xi(\langle \varpi \rangle_E) &= (-1)^{n_K - m_K + f_{\varpi, K} - m_{\varpi, K}} t_{\langle \varpi \rangle_{E/K}}({}_{A_K}\mathfrak{V}_\xi)t_{\langle \varpi \rangle_{E/K}}({}_{A_K^*}\mathfrak{V}_\xi) \end{aligned}$$

for a prime element $\varpi_E \in E_0K$ (see the beginning of Section 3.8). Here

$$\begin{aligned} f_{\varpi, K} &= f(E/K[\langle \varpi \rangle_E]) \text{ and} \\ m_{\varpi, K} &= \gcd(m_K, f_{\varpi, K}) = \gcd(m, n_K, f(E/K[\langle \varpi \rangle_E])). \end{aligned}$$

Now suppose that $(E/F, \xi) \in P_n(F)$. By the definition of admissibility, we can regard ξ as an admissible character over K and form the pair $(E/K, \xi) \in P_{n_K}(K)$.

PROPOSITION 5.7. *In this situation, we have*

$${}_{D_K}\nu\xi = \prod_{[g] \in (\mathcal{W}_E \backslash \mathcal{W}_K / \mathcal{W}_E)'} \zeta_{g, \xi}|_{E^\times}.$$

Proof. Notice that if \mathfrak{V} is a $\mathbf{k}_F\Psi_{E/F}$ -submodule of $\mathfrak{U}_{E/K}$, we have

$$t_{\boldsymbol{\mu}_{E/K}}^1(\mathfrak{V}) = t_{\boldsymbol{\mu}_{E/F}}^1(\mathfrak{V}) \text{ and } t_{\langle \varpi \rangle_{E/K}}(\mathfrak{V}) = t_{\langle \varpi \rangle_{E/F}}(\mathfrak{V}),$$

where $\boldsymbol{\mu}_{E/F}$ and $\langle \varpi \rangle_{E/F}$ are just $\boldsymbol{\mu}$ and $\langle \varpi \rangle$ respectively considered in (3.20). Hence we have

$${}_{D_K}\nu\xi|_{\boldsymbol{\mu}_E} = \prod_{[g] \in (\mathcal{W}_E \backslash \mathcal{W}_K / \mathcal{W}_E)'} \zeta_{g, \xi}|_{\boldsymbol{\mu}_E}.$$

It remains to consider the values of both characters at ϖ_E . Notice that the left side has value

$$(-1)^{n_K - m_K + f_{\varpi, K} - m_{\varpi, K}} t_{\langle \varpi \rangle_{E/K}}(A_K \mathfrak{Y}_\xi) t_{\langle \varpi \rangle_{E/K}}(A_K^* \mathfrak{Y}_\xi),$$

while the right side has value

$$(\text{a sign}) \cdot \prod_{[g] \in (\mathcal{W}_E \setminus \mathcal{W}_K / \mathcal{W}_E)'} t_{\langle \varpi \rangle}(A_K \mathfrak{Y}_{\xi, [g]}) t_{\langle \varpi \rangle}(A_K^* \mathfrak{Y}_{\xi, [g]}).$$

The t-factors on both sides are clearly equal. We will recall, by Theorem 5.5, the values of the sign on the right side in different cases and show that, in each case, this sign is equal to the one on the left side.

We first consider when f_K is odd, which can be reduced to the case when E/K is totally ramified. We further separate into cases.

- When e is odd, or when e is even and e_K is odd, then m_K is also odd. The sign on the left is $(-1)^{e_K - m_K} = 1$, while that on the right is also 1 since $\sigma^{e/2} \notin \mathcal{W}_K$.
- When e_K is even, the sign on the left is $(-1)^{e_K - m_K} = (-1)^m$ since $m_K \equiv m \pmod 2$, while that on the right is $(-1)^m$ since $\sigma^{e/2} \in \mathcal{W}_K$.

We then consider then f_K is even. Let L be the maximal unramified extension of E/K . We recall, after disregarding the symmetric ramified component (as we did at the beginning of Sub-section 5.3.2), the sign on the left is equal to (see (5.6))

$$(-1)^{e_K - m_L} (-1)^{n_K - m_K + f_{\varpi, K} - m_{\varpi, K}} = \begin{cases} 1 & \text{if } m_K \text{ is even,} \\ (-1)^{e_K + f_{\varpi, K} + 1} & \text{if } m_K \text{ is odd.} \end{cases}$$

Recall from Proposition 2.6 that the number $e_K + f_{\varpi, K} + 1$ is just the cardinality of

$$(\Gamma_E \setminus \Gamma_K / \Gamma_E)_{\text{sym-unram}} - \Gamma_{K[\varpi_E]}.$$

Hence by Theorem 5.5, the sign on the right side is

$$(-1)^{m(e_K + f_{\varpi, K} + 1)}.$$

By knowing that $m_K \equiv m \pmod 2$, the sign above is equal to the one on the left side. □

ACKNOWLEDGEMENTS

This work was started when the author was receiving funding from the European Research Council under the European Community’s Seventh Framework Programme (FP7/2007-2013) / ERC Grant agreement no.290766 (AAMOT) during his visit in IMJ, Paris. It was then finished when he became a post-doc in McMaster University. He would like to thank Vincent Sécherre for a few discussions in Paris, and Shaun Stevens for reading the manuscript and his comments. He would also like to thank the referee for the careful reading.

REFERENCES

- [Bad02] Alexandru Ioan Badulescu, *Correspondance de Jacquet-Langlands pour les corps locaux de caractéristique non nulle*, Ann. Sci. École Norm. Sup. (4) 35 (2002), no. 5, 695–747. MR 1951441 (2004i:11048)
- [BF83] Colin J. Bushnell and Albrecht Fröhlich, *Gauss sums and p -adic division algebras*, Lecture Notes in Mathematics, vol. 987, Springer-Verlag, Berlin-New York, 1983. MR 701540 (84m:12017)
- [BF85] C. J. Bushnell and A. Fröhlich, *Nonabelian congruence Gauss sums and p -adic simple algebras*, Proc. London Math. Soc. (3) 50 (1985), no. 2, 207–264. MR 772712 (86g:11071)
- [BH96] Colin J. Bushnell and Guy Henniart, *Local tame lifting for $GL(N)$. I. Simple characters*, Inst. Hautes Études Sci. Publ. Math. (1996), no. 83, 105–233. MR 1423022 (98m:11129)
- [BH05a] ———, *The essentially tame local Langlands correspondence. I*, J. Amer. Math. Soc. 18 (2005), no. 3, 685–710 (electronic). MR 2138141 (2006a:22014)
- [BH05b] ———, *The essentially tame local Langlands correspondence. II. Totally ramified representations*, Compos. Math. 141 (2005), no. 4, 979–1011. MR 2148193 (2006c:22017)
- [BH10] ———, *The essentially tame local Langlands correspondence, III: the general case*, Proc. Lond. Math. Soc. (3) 101 (2010), no. 2, 497–553. MR 2679700
- [BH11] ———, *The essentially tame Jacquet-Langlands correspondence for inner forms of $GL(n)$* , Pure Appl. Math. Q. 7 (2011), no. 3, Special Issue: In honor of Jacques Tits, 469–538. MR 2848585 (2012f:22027)
- [BHL10] Colin J. Bushnell, Guy Henniart, and Bertrand Lemaire, *Caractère et degré formel pour les formes intérieures de $GL(n)$ sur un corps local de caractéristique non nulle*, Manuscripta Math. 131 (2010), no. 1-2, 11–24. MR 2574988 (2010m:22013)
- [BK93] C.J. Bushnell and P.C. Kutzko, *The admissible dual of $GL(N)$ via compact open subgroups*, Annals of mathematics studies, Princeton University Press, 1993.
- [Bro96] Paul Broussous, *Extension du formalisme de Bushnell et Kutzko au cas d'une algèbre à division*, C. R. Acad. Sci. Paris Sér. I Math. 323 (1996), no. 3, 225–228. MR 1404764 (97f:22028)
- [BSS12] P. Broussous, V. Sécherre, and S. Stevens, *Smooth representations of $GL_m(D)$ V: Endo-classes*, Doc. Math. 17 (2012), 23–77. MR 2889743

- [DKV84] P. Deligne, D. Kazhdan, and M.-F. Vignéras, *Représentations des algèbres centrales simples p -adiques*, Representations of reductive groups over a local field, Travaux en Cours, Hermann, Paris, 1984, pp. 33–117. MR 771672 (86h:11044)
- [Gra07] Martin Grabitz, *Simple characters for principal orders in $M_m(D)$* , J. Number Theory 126 (2007), no. 1, 1–51. MR 2347425 (2008k:11125)
- [Gre55] J. A. Green, *The characters of the finite general linear groups*, Trans. Amer. Math. Soc. 80 (1955), 402–447. MR 0072878 (17,345e)
- [How77] Roger E. Howe, *Tamely ramified supercuspidal representations of GL_n* , Pacific J. Math. 73 (1977), no. 2, 437–460. MR 0492087 (58 #11241)
- [JL70] H. Jacquet and R. P. Langlands, *Automorphic forms on $GL(2)$* , Lecture Notes in Mathematics, Vol. 114, Springer-Verlag, Berlin, 1970. MR 0401654 (53 #5481)
- [KS99] Robert E. Kottwitz and Diana Shelstad, *Foundations of twisted endoscopy*, Astérisque (1999), no. 255, vi+190. MR 1687096 (2000k:22024)
- [Lan94] Serge Lang, *Algebraic number theory*, second ed., Graduate Texts in Mathematics, vol. 110, Springer-Verlag, New York, 1994. MR 1282723 (95f:11085)
- [LS87] R. P. Langlands and D. Shelstad, *On the definition of transfer factors*, Math. Ann. 278 (1987), no. 1-4, 219–271. MR 909227 (89c:11172)
- [Moy86] Allen Moy, *Local constants and the tame Langlands correspondence*, Amer. J. Math. 108 (1986), no. 4, 863–930. MR 853218 (88b:11081)
- [NM43] Tadasi Nakayama and Yozô Matsushima, *Über die multiplikative Gruppe einer p -adischen Divisionsalgebra*, Proc. Imp. Acad. Tokyo 19 (1943), 622–628. MR 0014081 (7,238a)
- [Rei03] I. Reiner, *Maximal orders*, London Mathematical Society Monographs. New Series, vol. 28, The Clarendon Press Oxford University Press, Oxford, 2003, Corrected reprint of the 1975 original, With a foreword by M. J. Taylor. MR 1972204 (2004c:16026)
- [Rog83] Jonathan D. Rogawski, *Representations of $GL(n)$ and division algebras over a p -adic field*, Duke Math. J. 50 (1983), no. 1, 161–196. MR 700135 (84j:12018)
- [Séc04] Vincent Sécherre, *Représentations lisses de $GL(m, D)$. I. Caractères simples*, Bull. Soc. Math. France 132 (2004), no. 3, 327–396. MR 2081220 (2005f:22027)

- [Séc05a] ———, *Représentations lisses de $GL(m, D)$. II. β -extensions*, Compos. Math. 141 (2005), no. 6, 1531–1550. MR 2188448 (2006j:22018)
- [Séc05b] ———, *Représentations lisses de $GL_m(D)$. III. Types simples*, Ann. Sci. École Norm. Sup. (4) 38 (2005), no. 6, 951–977. MR 2216835 (2007a:22010)
- [SS08] V. Sécherre and S. Stevens, *Représentations lisses de $GL(m, D)$. IV. Représentations supercuspidales*, J. Inst. Math. Jussieu 7 (2008), no. 3, 527–574. MR 2427423 (2009d:22023)
- [Tam] Kam-Fai Tam, *Admissible embeddings of L -tori and the essentially tame local Langlands correspondence, to appear in International Mathematics Research Notices*.
- [Tat79] J. Tate, *Number theoretic background*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 3–26. MR 546607 (80m:12009)
- [Zin92] Ernst-Wilhelm Zink, *Representation theory of local division algebras*, J. Reine Angew. Math. 428 (1992), 1–44. MR 1166506 (93i:11141)
- [Zin99] ———, *More on embeddings of local fields in simple algebras*, J. Number Theory 77 (1999), no. 1, 51–61. MR 1695699 (2001d:11116)

K.-F. Tam
Dept. of Math. & Stat.
McMaster University
1280 Main Street West
Hamilton, Ontario, Canada
L8S 4K1
geotam@math.mcmaster.ca

