

COMMUTATIVE ALGEBRAIC GROUPS UP TO ISOGENY

MICHEL BRION

Received: February 10, 2016

Revised: June 28, 2016

Communicated by Henning Krause

ABSTRACT. Consider the abelian category \mathcal{C}_k of commutative group schemes of finite type over a field k . By results of Serre and Oort, \mathcal{C}_k has homological dimension 1 (resp. 2) if k is algebraically closed of characteristic 0 (resp. positive). In this article, we explore the abelian category of commutative algebraic groups up to isogeny, defined as the quotient of \mathcal{C}_k by the full subcategory \mathcal{F}_k of finite k -group schemes. We show that $\mathcal{C}_k/\mathcal{F}_k$ has homological dimension 1, and we determine its projective or injective objects. We also obtain structure results for $\mathcal{C}_k/\mathcal{F}_k$, which take a simpler form in positive characteristics.

2010 Mathematics Subject Classification: 14K02, 14L15, 18E35, 20G07.

Keywords and Phrases: commutative algebraic groups, isogeny category, homological dimension.

1 INTRODUCTION

There has been much recent progress on the structure of algebraic groups over an arbitrary field; in particular, on the classification of pseudo-reductive groups (see [CGP15, CP15]). Yet commutative algebraic groups over an imperfect field remain somewhat mysterious, e.g., extensions with unipotent quotients are largely unknown; see [To13] for interesting results, examples, and questions.

In this article, we explore the category of commutative algebraic groups up to isogeny, in which the problems raised by imperfect fields become tractable; this yields rather simple and uniform structure results.

More specifically, denote by \mathcal{C}_k the category with objects the group schemes of finite type over the ground field k , and with morphisms, the homomorphisms of k -group schemes (all group schemes under consideration will be assumed commutative). By a result of Grothendieck (see [SGA3, VIA, Thm. 5.4.2]),

\mathcal{C}_k is an abelian category. We define the category of ‘algebraic groups up to isogeny’ as the quotient category of \mathcal{C}_k by the Serre subcategory of finite group schemes; then $\mathcal{C}_k/\mathcal{F}_k$ is obtained from \mathcal{C}_k by inverting all isogenies, i.e., all morphisms with finite kernel and cokernel.

It will be easier to deal with the full subcategory $\underline{\mathcal{C}}_k$ of $\mathcal{C}_k/\mathcal{F}_k$ with objects the smooth connected algebraic groups, since these categories turn out to be equivalent, and morphisms in $\underline{\mathcal{C}}_k$ admit a simpler description.

As a motivation for considering the ‘isogeny category’ $\underline{\mathcal{C}}_k$, note that some natural constructions involving algebraic groups are only exact up to isogeny; for example, the formations of the maximal torus or of the largest abelian variety quotient, both of which are not exact in \mathcal{C}_k . Also, some structure theorems for algebraic groups take on a simpler form when reformulated up to isogeny. A classical example is the Poincaré complete reducibility theorem, which is equivalent to the semi-simplicity of the isogeny category of abelian varieties, i.e., the full subcategory $\underline{\mathcal{A}}_k$ of $\underline{\mathcal{C}}_k$ with objects abelian varieties. Likewise, the isogeny category of tori, $\underline{\mathcal{T}}_k$, is semi-simple.

We gather our main results in the following:

THEOREM. (i) *The category $\underline{\mathcal{C}}_k$ is artinian and noetherian. Its simple objects are the additive group $\mathbb{G}_{a,k}$, the simple tori, and the simple abelian varieties.*

(ii) *The product functor $\underline{\mathcal{T}}_k \times \underline{\mathcal{U}}_k \rightarrow \underline{\mathcal{L}}_k$ yields an equivalence of categories, where $\underline{\mathcal{U}}_k$ (resp. $\underline{\mathcal{L}}_k$) denotes the isogeny category of unipotent (resp. linear) algebraic groups.*

(iii) *If $\text{char}(k) > 0$, then the product functor $\underline{\mathcal{S}}_k \times \underline{\mathcal{U}}_k \rightarrow \underline{\mathcal{C}}_k$ yields an equivalence of categories, where $\underline{\mathcal{S}}_k$ denotes the isogeny category of semi-abelian varieties. If in addition k is locally finite, then the product functor $\underline{\mathcal{T}}_k \times \underline{\mathcal{A}}_k \rightarrow \underline{\mathcal{S}}_k$ yields an equivalence of categories as well.*

(iv) *The base change under any purely inseparable field extension k' of k yields an equivalence of categories $\underline{\mathcal{C}}_k \rightarrow \underline{\mathcal{C}}_{k'}$.*

(v) *The homological dimension of $\underline{\mathcal{C}}_k$ is 1.*

We also describe the projective objects of the category $\underline{\mathcal{C}}_k$ (Theorem 5.14) and its injective objects (Theorem 5.16). Moreover, in characteristic 0, we obtain a structure result for that category (Proposition 5.13), which turns out to be more technical than in positive characteristics.

Let us now compare the above statements with known results on \mathcal{C}_k and its full subcategories \mathcal{A}_k (resp. \mathcal{T}_k , \mathcal{U}_k , \mathcal{L}_k , \mathcal{S}_k) of abelian varieties (resp. tori, unipotent groups, linear groups, semi-abelian varieties).

About (i) (an easy result, mentioned by Serre in [Se60]): \mathcal{C}_k is artinian and not noetherian. Also, every algebraic group is an iterated extension of ‘elementary’ groups; these are the simple objects of $\underline{\mathcal{C}}_k$ and the simple finite group schemes.

About (ii): denoting by \mathcal{M}_k the full subcategory of \mathcal{C}_k with objects the algebraic groups of multiplicative type, the product functor $\mathcal{M}_k \times \mathcal{U}_k \rightarrow \mathcal{L}_k$ yields an equivalence of categories if k is perfect. But over an imperfect field, there exist non-zero extensions of unipotent groups by tori, and these are only partially understood (see [To13, §9] again; our study of $\underline{\mathcal{C}}_k$ brings no new insight in these issues).

About (iii): the first assertion follows from recent structure results for algebraic groups (see [Br17, §5]), together with a lifting property for extensions of such groups with finite quotients (see [Br15] and [LA15]). The second assertion is a direct consequence of the Weil-Barsotti isomorphism (see e.g. [Oo66, §III.18]).

About (iv): this is a weak version of a result of Chow on abelian varieties, which asserts (in categorical language) that base change yields a fully faithful functor $\mathcal{A}_k \rightarrow \mathcal{A}_{k'}$ for any primary field extension k' of k (see [Ch55], and [Co06, §3] for a modern proof).

About (v), the main result of this article: recall that the homological dimension of an abelian category \mathcal{D} is the smallest integer, $\text{hd}(\mathcal{D})$, such that $\text{Ext}_{\mathcal{D}}^n(A, B) = 0$ for all objects A, B of \mathcal{D} and all $n > \text{hd}(\mathcal{D})$; these Ext groups are defined as equivalence classes of Yoneda extensions. In particular, $\text{hd}(\mathcal{D}) = 0$ if and only if \mathcal{D} is semi-simple.

It follows from work of Serre (see [Se60, 10.1 Thm. 1] together with [Oo64, Thm. 3.5]) that $\text{hd}(\mathcal{C}_k) = 1$ if k is algebraically closed of characteristic 0. Also, by a result of Oort (see [Oo66, Thm. 14.1]), $\text{hd}(\mathcal{C}_k) = 2$ if k is algebraically closed of positive characteristic. Building on these results, Milne determined $\text{hd}(\mathcal{C}_k)$ when k is perfect (see [Mi70, Thm. 1]); then the homological dimension can be arbitrarily large. In the approach of Serre and Oort, the desired vanishing of higher extension groups is obtained by constructing projective resolutions of elementary groups, in the category of pro-algebraic groups. The latter category contains \mathcal{C}_k as a full subcategory, and has enough projectives.

In contrast, to show that $\text{hd}(\underline{\mathcal{C}}_k) = 1$ over an arbitrary field k , we do not need to go to a larger category. We rather observe that tori are projective objects in $\underline{\mathcal{C}}_k$, and abelian varieties are injective objects there. This yields the vanishing of all but three extension groups between simple objects of $\underline{\mathcal{C}}_k$; two of the three remaining cases are handled directly, and the third one reduces to the known vanishing of $\text{Ext}_{\mathcal{C}_k}^2(\mathbb{G}_{a,k}, \mathbb{G}_{a,k})$ when k is perfect.

When k has characteristic 0, the fact that $\text{hd}(\mathcal{C}_k) \leq 1$ follows from a similar result for the category of Laumon 1-motives up to isogeny (obtained by Mazzari in [Ma10, Thm. 2.5]). Indeed, \mathcal{C}_k is equivalent to a Serre subcategory of the latter category; moreover, if an abelian category has homological dimension at most 1, then the same holds for any Serre subcategory (as follows e.g. from [Oo64, §3]). Likewise, the fact that the category of Deligne 1-motives up to isogeny has homological dimension at most 1 (due to Orgogozo, see [Or04, Prop. 3.2.4]) implies the corresponding assertion for the isogeny category of semi-abelian varieties over an arbitrary field.

Abelian categories of homological dimension 1 are called hereditary. The most studied hereditary categories consist either of finite-dimensional modules over a

finite-dimensional hereditary algebra, or of coherent sheaves on a weighted projective line (see e.g. [Ha01]). Such categories are k -linear and Hom-finite, i.e., all groups of morphisms are finite-dimensional vector spaces over the ground field k . But this seldom holds for the above isogeny categories. More specifically, $\underline{\mathcal{A}}_k$ and $\underline{\mathcal{T}}_k$ are both \mathbb{Q} -linear and Hom-finite, but not $\underline{\mathcal{C}}_k$ unless k is a number field. When k has characteristic 0, we may view $\underline{\mathcal{C}}_k$ as a mixture of k -linear and \mathbb{Q} -linear categories. This is already displayed by the full subcategory $\underline{\mathcal{V}}_k$ with objects the vector extensions of abelian varieties: as shown in §5.1, $\underline{\mathcal{V}}_k$ has enough projectives, and these are either the unipotent groups (k -linear objects), or the vector extensions of simple abelian varieties (\mathbb{Q} -linear objects).

In positive characteristic, one may also consider the quotient category of \mathcal{C}_k by the Serre subcategory \mathcal{I}_k of infinitesimal group schemes. This yields the abelian category of ‘algebraic groups up to purely inseparable isogeny’, which is equivalent to that introduced by Serre in [Se60]; as a consequence, it has homological dimension 1 if k is algebraically closed. For any arbitrary field k , the category $\mathcal{C}_k/\mathcal{I}_k$ is again invariant under purely inseparable field extensions; its homological properties may be worth investigating.

NOTATION AND CONVENTIONS. We will use the book [DG70] as a general reference, especially for affine algebraic groups, and the expository text [Br17] for some further results.

Throughout this text, we fix the ground field k and an algebraic closure \bar{k} ; the characteristic of k is denoted by $\text{char}(k)$. We denote by k_s the separable closure of k in \bar{k} , and by Γ_k the Galois group of k_s over k . We say that k is *locally finite*, if it is algebraic over \mathbb{F}_p for some prime p .

By an *algebraic k -group*, we mean a commutative group scheme G of finite type over k ; we denote by G^0 the neutral component of G . The group law of G will be denoted additively: $(x, y) \mapsto x + y$.

By a *k -subgroup* of G , we mean a closed k -subgroup scheme. *Morphisms* are understood to be homomorphisms of k -group schemes. The (scheme-theoretic) image of a morphism $f : G \rightarrow H$ will be denoted by $\text{Im}(f)$ or $f(G)$, and the (scheme-theoretic) pull-back of a k -subgroup $H' \subset H$, by $G \times_H H'$ or $f^{-1}(H')$. Recall that an *abelian variety* over k is a smooth connected proper algebraic k -group. Also, recall that a k -group scheme G is an affine algebraic k -group if and only if G is isomorphic to a k -subgroup of the general linear group $\text{GL}_{n,k}$ for some n . We will thus call affine algebraic k -groups *linear*. We say that an algebraic k -group G is *of multiplicative type* if G is isomorphic to a k -subgroup of some k -torus.

To simplify the notation, *we will suppress the mention of the ground field k* whenever this yields no confusion. For example, the category \mathcal{C}_k will be denoted by \mathcal{C} , except when we use base change by a field extension.

Given an algebraic group G and two subgroups G_1, G_2 , we denote by $G_1 + G_2$ the subgroup of G generated by G_1 and G_2 . Thus, $G_1 + G_2$ is the image of the morphism $G_1 \times G_2 \rightarrow G$, $(x_1, x_2) \mapsto x_1 + x_2$.

An *isogeny* is a morphism with finite kernel and cokernel. Two algebraic groups G_1, G_2 are *isogenous* if they can be connected by a chain of isogenies.

We say that two subgroups G_1, G_2 of an algebraic group G are *commensurable* if both quotients $G_1/G_1 \cap G_2$ and $G_2/G_1 \cap G_2$ are finite; then G_1 and G_2 are isogenous.

Given an algebraic group G and a non-zero integer n , the multiplication by n yields a morphism

$$n_G : G \longrightarrow G.$$

We denote its kernel by $G[n]$, and call it the *n-torsion subgroup*. We say that G is *divisible* if n_G is an epimorphism for all $n \neq 0$; then n_G is an isogeny for all such n . When $\text{char}(k) = 0$, the divisible groups are the connected algebraic groups; when $\text{char}(k) = p > 0$, they are just the *semi-abelian varieties*, that is, the extensions of abelian varieties by tori (see e.g. [Br17, Thm. 5.6.3] for the latter result).

Still assuming that $\text{char}(k) = p > 0$, we say that an algebraic group G is a *p-group* if $p_G^n = 0$ for $n \gg 0$. Examples of *p-groups* include the unipotent groups and the connected finite algebraic groups, also called *infinitesimal*.

2 STRUCTURE OF ALGEBRAIC GROUPS

2.1 PRELIMINARY RESULTS

We will use repeatedly the following simple observation:

LEMMA 2.1. *Let G be a smooth connected algebraic group.*

- (i) *If G' is a subgroup of G such that G/G' is finite, then $G' = G$.*
- (ii) *Any isogeny $f : H \rightarrow G$ is an epimorphism.*

Proof. (i) The quotient G/G' is smooth, connected and finite, hence zero.

(ii) This follows from (i) applied to $\text{Im}(f) \subset H$. □

The following lifting result for finite quotients will also be frequently used:

LEMMA 2.2. *Let G be an algebraic group, and H a subgroup such that G/H is finite.*

- (i) *There exists a finite subgroup $F \subset G$ such that $G = H + F$.*
- (ii) *If G/H is infinitesimal (resp. a finite p -group), then F may be chosen infinitesimal (resp. a finite p -group) as well.*

Proof. (i) This is a special case of [Br15, Thm. 1.1].

(ii) Assume G/H infinitesimal. Then the quotient $G/(H + F^0)$ is infinitesimal (as a quotient of G/H) and étale (as a quotient of F/F^0), hence zero. Thus, we may replace F with F^0 , an infinitesimal subgroup.

Next, assume that G/H is a finite p -group. Denote by $F[p^\infty]$ the largest p -subgroup of F . Then the quotient $G/(H + F[p^\infty])$ is a finite p -group and is killed by the order of $F/F[p^\infty]$. Since the latter order is prime to p , we obtain $G/(H + F[p^\infty]) = 0$. Thus, we may replace F with $F[p^\infty]$. \square

Next, we recall a version of a theorem of Chevalley:

THEOREM 2.3. (i) *Every algebraic group G contains a linear subgroup L such that G/L is an abelian variety. Moreover, L is unique up to commensurability in G , and G/L is unique up to isogeny.*

(ii) *If G is connected, then there exists a smallest such subgroup, $L = L(G)$, and this subgroup is connected.*

(iii) *If in addition G is smooth, then every morphism from G to an abelian variety factors uniquely through the quotient map $G \rightarrow G/L(G)$.*

Proof. The assertion (ii) follows from [Ra70, Lem. IX 2.7] (see also [BLR90, 9.2 Thm. 1]).

To prove (i), note that G contains a finite subgroup F such that G/F is connected (as follows from Lemma 2.2). Then we may take for L the pull-back of a linear subgroup of G/F with quotient an abelian variety. If L' is another linear subgroup of G such that G/L' is an abelian variety, then $L + L'$ is linear, as a quotient of $L \times L'$. Moreover, the natural map $q : G/L \rightarrow G/(L + L')$ is the quotient by $(L + L')/L$, a linear subgroup of the abelian variety G/L . It follows that $(L + L')/L$ is finite; thus, q is an isogeny and $L'/L \cap L'$ is finite. Likewise, $q' : G/L' \rightarrow G/(L + L')$ is an isogeny and $L/L \cap L'$ is finite; this completes the proof of (i).

Finally, the assertion (iii) is a consequence of [Br17, Thm. 4.3.4]. \square

The structure of linear algebraic groups may be described as follows (see [DG70, Thm. IV.3.1.1]):

THEOREM 2.4. *Let G be a linear algebraic group. Then G has a largest subgroup of multiplicative type, M ; moreover, G/M is unipotent. If k is perfect, then $G = M \times U$, where U denotes the largest unipotent subgroup of G .*

Also, note the following orthogonality relations:

PROPOSITION 2.5. (i) *Let M be a group of multiplicative type, and U a unipotent group. Then $\mathrm{Hom}_{\mathcal{C}}(M, U) = 0 = \mathrm{Hom}_{\mathcal{C}}(U, M)$.*

(ii) *Let L be a linear algebraic group, and A an abelian variety. Then $\mathrm{Hom}_{\mathcal{C}}(A, L) = 0$, and every morphism $L \rightarrow A$ has finite image. Moreover, $\mathrm{Hom}_{\mathcal{C}}(L, A)$ is n -torsion for some positive integer n .*

Proof. (i) This follows from [DG70, Cor. IV.2.2.4].

(ii) The image of a morphism $A \rightarrow L$ is proper, smooth, connected and affine, hence zero. Likewise, the image of a morphism $L \rightarrow A$ is affine and proper, hence finite.

To show the final assertion, we may replace k with any field extension, and hence assume that k is perfect. Then the reduced neutral component L_{red}^0 is a smooth connected subgroup of L , the quotient L/L_{red}^0 is finite, and $\text{Hom}_{\mathcal{C}}(L_{\text{red}}^0, A) = 0$ by the above argument. As a consequence, we have $\text{Hom}_{\mathcal{C}}(L, A) = \text{Hom}_{\mathcal{C}}(L/L_{\text{red}}^0, A)$ and this group is n -torsion, where n denotes the order of the finite group scheme L/L_{red}^0 (indeed, L/L_{red}^0 is n -torsion in view of [SGA3, VIIA, Prop. 8.5]). \square

Next, we obtain a key preliminary result. To state it, recall that a unipotent group G is said to be *split* if it admits a finite increasing sequence of subgroups $0 = G_0 \subset G_1 \subset \dots \subset G_n = G$ such that $G_i/G_{i-1} \cong \mathbb{G}_a$ for $i = 1, \dots, n$.

PROPOSITION 2.6. *Let G be an algebraic group.*

- (i) *There exists a finite subgroup $F \subset G$ such that G/F is smooth and connected.*
- (ii) *If G is unipotent, then we may choose F such that G/F is split.*

Proof. (i) By Lemma 2.2, we have $G = G^0 + F$ for some finite subgroup $F \subset G$. Thus, $G/F \cong G^0/F \cap G^0$ is connected; this yields the assertion if $\text{char}(k) = 0$. If $\text{char}(k) = p > 0$, then we may assume G connected by the above step. Consider the relative Frobenius morphism $F_{G/k} : G \rightarrow G^{(p)}$ and its iterates $F_{G/k}^n : G \rightarrow G^{(p^n)}$, where $n \geq 1$. Then $\text{Ker}(F_{G/k}^n)$ is finite for all n ; moreover, $G/\text{Ker}(F_{G/k}^n)$ is smooth for $n \gg 0$ (see [SGA3, VIIA, Prop. 8.3]), and still connected.

(ii) We argue by induction on the dimension of G . The statement is obvious if $\dim(G) = 0$. In the case where $\dim(G) = 1$, we may assume that G is smooth and connected in view of Lemma 2.2 again; then G is a k -form of \mathbb{G}_a . By [Ru70, Thm. 2.1], there exists an exact sequence

$$0 \longrightarrow G \longrightarrow \mathbb{G}_a^2 \xrightarrow{f} \mathbb{G}_a \longrightarrow 0,$$

where $f \in \mathcal{O}(\mathbb{G}_a^2) \cong k[x, y]$ satisfies $f(x, y) = y^{p^n} - a_0 x - a_1 x^p - \dots - a_m x^{p^m}$ for some integers $m, n \geq 0$ and some $a_0, \dots, a_m \in k$ with $a_0 \neq 0$. Thus, the projection

$$p_1 : G \longrightarrow \mathbb{G}_a, \quad (x, y) \longmapsto x$$

lies in an exact sequence

$$0 \longrightarrow \alpha_{p^n} \longrightarrow G \xrightarrow{p_1} \mathbb{G}_a \longrightarrow 0,$$

where α_{p^n} denotes the kernel of the endomorphism $x \mapsto x^{p^n}$ of \mathbb{G}_a . This yields the assertion in this case.

If $\dim(G) \geq 2$, then we may choose a subgroup $G_1 \subset G$ such that $0 < \dim(G_1) < \dim(G)$ (as follows from [DG70, Prop. IV.2.2.5]). By the induction assumption for G/G_1 , there exists a subgroup $G_2 \subset G$ such that $G_1 \subset G_2$,

G_2/G_1 is finite, and G/G_2 is split. Next, the induction assumption for G_2 yields a finite subgroup $F \subset G_2$ such that G_2/F is split. Then G/F is split as well. \square

REMARK 2.7. By Proposition 2.6, every algebraic group G admits an isogeny $u : G \rightarrow H$, where H is smooth and connected. If k is perfect, then there also exists an isogeny $v : K \rightarrow G$, where K is smooth and connected: just take v to be the inclusion of the reduced neutral component G_{red}^0 . But this fails over any imperfect field k . Indeed, if such an isogeny v exists, then its image must be G_{red}^0 . On the other hand, by [SGA3, VIA, Ex. 1.3.2], there exists a connected algebraic group G such that G_{red} is not a subgroup.

By combining Lemma 2.2, Theorems 2.3 and 2.4, and Proposition 2.6, we obtain readily:

PROPOSITION 2.8. *Every algebraic group G admits a finite increasing sequence of subgroups $0 = G_0 \subset G_1 \subset \dots \subset G_n = G$ such that each subquotient G_i/G_{i-1} , $i = 1, \dots, n$, is finite or isomorphic to \mathbb{G}_a , a simple torus, or a simple abelian variety. Moreover, G is linear if and only if no abelian variety occurs.*

2.2 CHARACTERISTIC ZERO

In this subsection, we assume that $\text{char}(k) = 0$. Recall that every unipotent group is isomorphic to the additive group of its Lie algebra via the exponential map; this yields an equivalence between the category \mathcal{U} of unipotent groups and the category of finite-dimensional k -vector spaces (see [DG70, Prop. IV.2.4.2]). In particular, every unipotent group is connected.

Next, consider a connected algebraic group G . By Theorem 2.3, there is a unique exact sequence $0 \rightarrow L \rightarrow G \rightarrow A \rightarrow 0$, where A is an abelian variety, and L is connected and linear. Moreover, in view of Theorem 2.4, we have $L = T \times U$, where T is a torus and U is unipotent.

We now extend the latter structure results to possibly non-connected groups:

THEOREM 2.9. (i) *Every algebraic group G lies in an exact sequence*

$$0 \longrightarrow M \times U \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where M is of multiplicative type, U is unipotent, and A is an abelian variety. Moreover, U is the largest unipotent subgroup of G : the unipotent radical, $R_u(G)$. Also, M is unique up to commensurability in G , and A is unique up to isogeny.

(ii) *The formation of the unipotent radical commutes with base change under field extensions, and yields an exact functor*

$$R_u : \mathcal{C} \longrightarrow \mathcal{U},$$

right adjoint to the inclusion $\mathcal{U} \rightarrow \mathcal{C}$.

(iii) *The projective objects of \mathcal{C} are the unipotent groups.*

Proof. (i) Theorem 2.3 yields an exact sequence

$$0 \longrightarrow L \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where L is linear (possibly non-connected), and A is an abelian variety. By Theorem 2.4, we have $L = M \times U$, where M is of multiplicative type and U is unipotent.

Since M and A have no non-trivial unipotent subgroups, we have $U = R_u(G)$. Given another exact sequence

$$0 \longrightarrow M' \times U \longrightarrow G \longrightarrow A' \longrightarrow 0$$

satisfying the same assumptions, the image of M' in $A \cong G/(M \times U)$ is finite by Proposition 2.5. In other words, the quotient $M'/M' \cap (M \times U)$ is finite. Likewise, $M/M \cap (M' \times U)$ is finite as well. Since we have $(M \times U) \cap M' = M \cap M' = M \cap (M' \times U)$, we see that M, M' are commensurable in G . Then $A = G/(M \times U)$ and $A' = G/(M' \times U)$ are both quotients of $G/(M \cap M') \times U$ by finite subgroups, and hence are isogenous.

(ii) In view of (i), $G/R_u(G)$ is an extension of an abelian variety by a group of multiplicative type. Since these two classes of algebraic groups are stable under base change by any field extension k' of k , it follows that $(G/R_u(G))_{k'}$ has zero unipotent radical. Thus, $R_u(G)_{k'} = R_u(G_{k'})$.

Next, note that every morphism $f : G \rightarrow H$ sends $R_u(G)$ to $R_u(H)$. Consider an exact sequence

$$0 \longrightarrow G_1 \xrightarrow{f} G_2 \xrightarrow{g} G_3 \longrightarrow 0$$

and the induced complex

$$0 \longrightarrow R_u(G_1) \xrightarrow{R_u(f)} R_u(G_2) \xrightarrow{R_u(g)} R_u(G_3) \longrightarrow 0.$$

Clearly, $R_u(f)$ is a monomorphism. Also, we have

$$\text{Ker}(R_u(g)) = R_u(G_2) \cap \text{Ker}(g) = R_u(G_2) \cap \text{Im}(f) = \text{Im}(R_u(f)).$$

We now show that $R_u(g)$ is an epimorphism. For this, we may replace G_2 with $g^{-1}(R_u(G_3))$, and hence assume that G_3 is unipotent. Next, we may replace G_2 with $G_2/R_u(G_2)$, and hence assume (in view of (i) again) that G_2 is an extension of an abelian variety by a group of multiplicative type. Then $\text{Hom}(G_2, G_3) = 0$ by Proposition 2.5; this completes the proof of the exactness assertion.

The assertion about adjointness follows from the fact that every morphism $U \rightarrow H$, where U is unipotent and H arbitrary, factors through a unique morphism $U \rightarrow R_u(H)$.

(iii) Consider an epimorphism $\varphi : G \rightarrow H$, a unipotent group U , and a morphism $\psi : U \rightarrow H$. Then ψ factors through $R_u(H)$. Also, by (ii), φ restricts to

an epimorphism $R_u(G) \rightarrow R_u(H)$, which admits a section as unipotent groups are just vector spaces. Thus, ψ lifts to a morphism $U \rightarrow G$. This shows that U is projective in \mathcal{C} .

Conversely, let G be a projective object in \mathcal{C} . We claim that the (abstract) group $\mathrm{Hom}_{\mathcal{C}}(G, H)$ is divisible for any divisible algebraic group H . Indeed, the exact sequence

$$0 \longrightarrow H[n] \longrightarrow H \xrightarrow{nH} H \longrightarrow 0$$

yields an exact sequence

$$0 \longrightarrow \mathrm{Hom}_{\mathcal{C}}(G, H[n]) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(G, H) \xrightarrow{\times n} \mathrm{Hom}_{\mathcal{C}}(G, H) \longrightarrow 0,$$

for any positive integer n .

Next, the exact sequence $0 \rightarrow L \rightarrow G \rightarrow A \rightarrow 0$ yields an exact sequence

$$0 \longrightarrow \mathrm{End}_{\mathcal{C}}(A) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(G, A) \longrightarrow \mathrm{Hom}_{\mathcal{C}}(L, A),$$

where the abelian group $\mathrm{End}_{\mathcal{C}}(A)$ is free of finite rank (see [Mi86, Thm. 12.5]), and $\mathrm{Hom}_{\mathcal{C}}(L, A)$ is killed by some positive integer (Proposition 2.5). On the other hand, $\mathrm{Hom}_{\mathcal{C}}(G, A)$ is divisible by the above claim. It follows that $\mathrm{End}_{\mathcal{C}}(A)$ is zero, and hence so is A . Thus, G is linear, and hence $G = M \times U$ as above. Since U is projective, so is M . Choose a torus T containing M ; then again, the group $\mathrm{Hom}_{\mathcal{C}}(M, T)$ is finitely generated and divisible, hence zero. Thus, $T = 0$ and $G = U$. \square

REMARK 2.10. With the notation of the above theorem, we have a natural map

$$G \longrightarrow G/M \times_A G/U,$$

which is a morphism of $M \times U$ -torsors over A , and hence an isomorphism. Moreover, G/M is an extension of an abelian variety by a unipotent group; such ‘vector extensions’ will be studied in detail in §5.1. Also, G/U is an extension of an abelian variety by a group of multiplicative type, and hence of a semi-abelian variety by a finite group. The semi-abelian varieties will be considered in §5.2.

2.3 POSITIVE CHARACTERISTICS

In this subsection, we assume that $\mathrm{char}(k) = p > 0$. Then the assertions of Theorem 2.9 are no longer valid. For example, the formation of the unipotent radical (the largest smooth connected unipotent subgroup) is not exact, and does not commute with arbitrary field extensions either (see Remark 2.12 (i) for details). Also, \mathcal{C} has no non-zero projective objects, as will be shown in Corollary 5.15. Yet Theorem 2.9 has a useful analogue, in which the unipotent radical is replaced by the largest unipotent quotient:

THEOREM 2.11. *Let G be an algebraic group.*

- (i) G has a smallest subgroup H such that $U := G/H$ is unipotent. Moreover, H is an extension of an abelian variety A by a group of multiplicative type M . Also, M is unique up to commensurability in G , and A is unique up to isogeny.
- (ii) Every morphism $H \rightarrow U$ is zero; every morphism $U \rightarrow H$ has finite image.
- (iii) The formation of U commutes with base change under field extensions, and yields a functor

$$U : \mathcal{C} \longrightarrow \mathcal{U},$$

which is left adjoint to the inclusion of \mathcal{U} in \mathcal{C} . Moreover, every exact sequence in \mathcal{C}

$$0 \longrightarrow G_1 \xrightarrow{f} G_2 \xrightarrow{g} G_3 \longrightarrow 0$$

yields a right exact sequence

$$0 \longrightarrow U(G_1) \xrightarrow{U(f)} U(G_2) \xrightarrow{U(g)} U(G_3) \longrightarrow 0,$$

where $\text{Ker}(U(f))$ is a finite p -group.

- (iv) There exists a subgroup $V \subset G$ such that $G = H + V$ and $H \cap V$ is a finite p -group.

Proof. (i) Since the underlying topological space of G is noetherian, we may choose a subgroup $H \subset G$ such that G/H is unipotent, and H is minimal for this property. Let $H' \subset G$ be another subgroup such that G/H' is unipotent. Then so is $G/H \cap H'$ in view of the exact sequence

$$0 \longrightarrow G/H \cap H' \longrightarrow G/H \times G/H'.$$

By minimality of H , it follows that $H \cap H' = H$, i.e., $H \subset H'$. Thus, H is the smallest subgroup with unipotent quotient.

Since the class of unipotent groups is stable under extensions, every unipotent quotient of H is zero. Also, by the affinization theorem (see [Br17, Thm. 1, Prop. 5.5.1]), H is an extension of a linear algebraic group L by a semi-abelian variety S . Then every unipotent quotient of L is zero, and hence L must be of multiplicative type in view of Theorem 2.4. By [DG70, Cor. IV.1.3.9], the reduced neutral component L_{red}^0 is its maximal torus, T ; the quotient L/T is a finite group of multiplicative type. Denote by S' the preimage of T in H ; then S' is a semi-abelian variety (extension of T by S) and we have an exact sequence

$$0 \longrightarrow S' \longrightarrow H \longrightarrow L/T \longrightarrow 0.$$

By Lemma 2.2, there exists a finite subgroup $F \subset H$ such that $H = S' + F$; equivalently, the quotient morphism $H \rightarrow L/T$ restricts to an epimorphism

$F \rightarrow L/T$. Also, by Theorem 2.4 again, F has a largest subgroup of multiplicative type, M_F , and the quotient F/M_F is unipotent. Since L/T is of multiplicative type, it follows that the composition $M_F \rightarrow F \rightarrow L/T$ is an epimorphism as well. Thus, we may replace F with M_F , and assume that F is of multiplicative type. Let T' be the maximal torus of the semi-abelian variety S' , and $M := T' + F$. Then M is of multiplicative type; moreover, H/M is a quotient of S'/T' , and hence is an abelian variety. The uniqueness assertions may be checked as in the proof of Theorem 2.9.

(ii) This follows readily from Proposition 2.5.

(iii) The assertion on base change under field extensions follows from the stability of the classes of unipotent groups, abelian varieties, and groups of multiplicative type, under such base changes. The adjointness assertion may be checked as in the proof of Theorem 2.9 (ii).

Next, consider an exact sequence as in the statement. Clearly, $U(g)$ is an epimorphism. Also, $\text{Ker}(U(g))/\text{Im}(U(f)) \cong g^{-1}(H_3)/(u(G_1) + H_2)$, where H_i denotes the kernel of the quotient map $G_i \rightarrow U(G_i)$. Thus, $\text{Ker}(U(g))/\text{Im}(U(f))$ is a quotient of $g^{-1}(H_3)/u(G_1) \cong H_3$. As $\text{Ker}(U(g))/\text{Im}(U(f))$ is unipotent, it is trivial by (ii). Finally, $\text{Ker}(U(f)) \cong (G_1 \cap u^{-1}(H_2))/H_1$ is isomorphic to a subgroup of $H_2/u(H_1)$. Moreover, $H_2/u(H_1)$ is an extension of an abelian variety by a group of multiplicative type. Since $\text{Ker}(U(f))$ is unipotent, it is finite by (ii) again.

(iv) If G is linear, then $H = M$ is the largest subgroup scheme of multiplicative type of G . By [Br17, Thm. 5.3.1], there exists a subgroup scheme $W \subset G$ such that $G = H + W$ and $H \cap W$ is finite. Equivalently, the composition $W \rightarrow G \rightarrow U$ is an epimorphism and its kernel is finite. As a consequence, W is n -torsion for some $n \geq 1$, and hence $W = V + F$ where V is a p -group and F is finite of order prime to p . Thus, $G/(H + V)$ is a quotient of U and a quotient of F , hence trivial. So $G = H + V$, where $H \cap V$ is a finite p -group. For an arbitrary algebraic group G , Theorem 2.3 yields an exact sequence

$$0 \longrightarrow L \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where L is linear and A is an abelian variety. In view of (iii) and the vanishing of $U(A)$, we obtain an exact sequence

$$0 \longrightarrow F \longrightarrow U(L) \longrightarrow U(G) \longrightarrow 0,$$

where F is a finite p -group. By the preceding step, there exists a subgroup $V \subset L$ such that the composition $V \rightarrow L \rightarrow U(L)$ is an epimorphism with kernel a finite p -group. Thus, so is the composition $V \rightarrow G \rightarrow U(G)$. \square

REMARKS 2.12. (i) The formation of the unipotent radical does not commute with purely inseparable field extensions, in view of [SGA3, XVII.C.5]. This formation is not exact either, as seen e.g. from the exact sequence

$$0 \longrightarrow \alpha_p \longrightarrow \mathbb{G}_a \xrightarrow{F} \mathbb{G}_a \longrightarrow 0,$$

where F denotes the relative Frobenius endomorphism. When k is perfect, one may show that every exact sequence

$$0 \longrightarrow G_1 \longrightarrow G_2 \longrightarrow G_3 \longrightarrow 0$$

in \mathcal{C} yields a complex

$$0 \longrightarrow R_u(G_1) \longrightarrow R_u(G_2) \longrightarrow R_u(G_3) \longrightarrow 0$$

with finite homology groups. But this fails when k is imperfect; more specifically, choose a finite purely inseparable field extension K of k of degree p , and consider $G := R_{K/k}(\mathbb{G}_{m,K})$, where $R_{K/k}$ denotes the Weil restriction. By [CGP15, Prop. A.5.11], G is smooth, connected, and lies in an exact sequence

$$0 \longrightarrow \mathbb{G}_m \longrightarrow G \longrightarrow U \longrightarrow 0,$$

where U is unipotent of dimension $p - 1$. Moreover, every morphism from a smooth connected unipotent group to G is constant, as follows from the adjointness property of the Weil restriction (see [CGP15, (A.5.1)]). In other terms, $R_u(G) = 0$.

(ii) The functor U of Theorem 2.11 is not left exact, as seen from the exact sequence

$$0 \longrightarrow \alpha_p \longrightarrow E \xrightarrow{F} E^{(p)} \longrightarrow 0,$$

where E denotes a supersingular elliptic curve, and F its relative Frobenius morphism. Also, note that the torsion subgroups $E[p^n]$, where $n \geq 1$, form a strictly increasing sequence of infinitesimal unipotent groups; in particular, E has no largest connected unipotent subgroup.

COROLLARY 2.13. *Let G be an algebraic group.*

- (i) *There exists a finite subgroup $F \subset G$ such that $G/F \cong S \times U$, where S is a semi-abelian variety, and U a split unipotent group. Moreover, S and U are unique up to isogeny.*
- (ii) *If k is locally finite, then we may choose F so that $S \cong T \times A$, where T is a torus, and A an abelian variety. Moreover, T and A are unique up to isogeny.*

Proof. (i) With the notation of Theorem 2.11, we have isomorphisms

$$G/H \cap V \cong G/V \times G/H \cong (H/H \cap V) \times U.$$

Also, $H/H \cap V$ is an extension of an abelian variety, $H/(H \cap V) + M$, by a group of multiplicative type, $M/M \cap V$. Moreover, U is an extension of a split unipotent group by a finite group (Proposition 2.6). Thus, we may assume that $G = H$. Then G_{red}^0 is a semi-abelian variety, as follows from [Br17, Lem. 5.6.1]. Since G/G_{red}^0 is finite, applying Lemma 2.2 yields that G is an extension of a semi-abelian variety by a finite group.

(ii) By [Br17, Cor. 5.5.5], there exists an abelian subvariety $A \subset S$ such that $S = T + A$, where $T \subset S$ denotes the maximal torus. Then $T \cap A$ is finite, and $S/T \cap A \cong T/T \cap A \times A/T \cap A$.

This completes the proof of the existence assertions in (i) and (ii). The uniqueness up to isogeny follows from Proposition 2.5. \square

3 THE ISOGENY CATEGORY OF ALGEBRAIC GROUPS

3.1 DEFINITION AND FIRST PROPERTIES

Recall that \mathcal{C} denotes the category of commutative algebraic groups, and \mathcal{F} the full subcategory of finite groups. Since \mathcal{F} is stable under taking subobjects, quotients and extensions, we may form the quotient category \mathcal{C}/\mathcal{F} ; it has the same objects as \mathcal{C} , and its morphisms are defined by

$$\mathrm{Hom}_{\mathcal{C}/\mathcal{F}}(G, H) = \lim_{\rightarrow} \mathrm{Hom}_{\mathcal{C}}(G', H/H'),$$

where the direct limit is taken over all subgroups $G' \subset G$ such that G/G' is finite, and all finite subgroups $H' \subset H$. The category \mathcal{C}/\mathcal{F} is abelian, and comes with an exact functor

$$Q : \mathcal{C} \longrightarrow \mathcal{C}/\mathcal{F},$$

which is the identity on objects and the natural map

$$\mathrm{Hom}_{\mathcal{C}}(G, H) \longrightarrow \lim_{\rightarrow} \mathrm{Hom}_{\mathcal{C}}(G', H/H'), \quad f \longmapsto \underline{f}$$

on morphisms. The quotient functor Q satisfies the following universal property: given an exact functor $R : \mathcal{C} \rightarrow \mathcal{D}$, where \mathcal{D} is an abelian category, such that $R(F) = 0$ for any finite group F , there exists a unique exact functor $S : \mathcal{C}/\mathcal{F} \rightarrow \mathcal{D}$ such that $R = S \circ Q$ (see [Ga62, Cor. III.1.2, Cor. III.1.3] for these results).

Alternatively, \mathcal{C}/\mathcal{F} may be viewed as the localization of \mathcal{C} at the multiplicative system of isogenies (see [GZ67, §I.2] or [SP16, §4.26] for localization of categories); this is easily checked by arguing as in the proof of [SP16, Lem. 12.9.6]. We now show that \mathcal{C}/\mathcal{F} is equivalent to a category with somewhat simpler objects and morphisms:

LEMMA 3.1. *Let $\underline{\mathcal{C}}$ be the full subcategory of \mathcal{C}/\mathcal{F} with objects the smooth connected algebraic groups.*

- (i) *The inclusion of $\underline{\mathcal{C}}$ in \mathcal{C}/\mathcal{F} is an equivalence of categories.*
- (ii) *$\mathrm{Hom}_{\underline{\mathcal{C}}}(G, H) = \lim \mathrm{Hom}_{\mathcal{C}}(G, H/H')$, where the direct limit is taken over all finite subgroups $H' \subset H$.*

- (iii) Let $\underline{f} \in \text{Hom}_{\underline{\mathcal{C}}}(G, H)$ be represented by a morphism $f : G \rightarrow H/H'$ in \mathcal{C} . Then \underline{f} is zero (resp. a monomorphism, an epimorphism, an isomorphism) if and only if f is zero (resp. has a finite kernel, is an epimorphism, is an isogeny).

Proof. (i) This follows from Proposition 2.6.

(ii) This follows from Lemma 2.1.

(iii) By [Ga62, Lem. III.1.2], \underline{f} is zero (resp. a monomorphism, an epimorphism) if and only if $\text{Im}(f)$ (resp. $\overline{\text{Ker}}(f)$, $\text{Coker}(f)$) is finite. By Lemma 2.1 again, the finiteness of $\text{Im}(f)$ is equivalent to $f = 0$, and the finiteness of $\text{Coker}(f)$ is equivalent to f being an epimorphism. As a consequence, \underline{f} is an isomorphism if and only if f is an isogeny. \square

The abelian category $\underline{\mathcal{C}}$ will be called the *isogeny category of (commutative) algebraic groups*. Every exact functor $R : \mathcal{C} \rightarrow \mathcal{D}$, where \mathcal{D} is an abelian category and $R(f)$ is an isomorphism for any isogeny f , factors uniquely through $\underline{\mathcal{C}}$ (indeed, R must send any finite group to zero).

We may now prove the assertion (i) of the main theorem:

PROPOSITION 3.2. (i) *The category $\underline{\mathcal{C}}$ is noetherian and artinian.*

- (ii) *The simple objects of $\underline{\mathcal{C}}$ are \mathbb{G}_a , the simple tori, and the simple abelian varieties.*

Proof. (i) Let G be a smooth connected algebraic group, and $(G_n)_{n \geq 0}$ an increasing sequence of subobjects of G in $\underline{\mathcal{C}}$, i.e., each G_n is smooth, connected, and equipped with a \mathcal{C} -morphism

$$\varphi_n : G_n \longrightarrow G/G'_n,$$

where $\text{Ker}(\varphi_n)$ and G'_n are finite; moreover, we have \mathcal{C} -morphisms

$$\psi_n : G_n \longrightarrow G_{n+1}/G''_{n+1},$$

where $\text{Ker}(\psi_n)$ and G''_{n+1} are finite. Thus, $\dim(G_n) \leq \dim(G_{n+1}) \leq \dim(G)$. It follows that $\dim(G_n) = \dim(G_{n+1})$ for $n \gg 0$, and hence ψ_n is an isogeny. So $G_n \cong G_{n+1}$ in $\underline{\mathcal{C}}$ for $n \gg 0$. This shows that $\underline{\mathcal{C}}$ is noetherian. One may check likewise that $\underline{\mathcal{C}}$ is artinian.

- (ii) This follows from Proposition 2.8. \square

Next, we relate the short exact sequences in \mathcal{C} with those in $\underline{\mathcal{C}}$:

LEMMA 3.3. *Consider a short exact sequence in \mathcal{C} ,*

$$\xi : 0 \longrightarrow G_1 \xrightarrow{u} G_2 \xrightarrow{v} G_3 \longrightarrow 0,$$

where G_1, G_2, G_3 are smooth and connected. Then ξ splits in $\underline{\mathcal{C}}$ if and only if the push-out f_ξ splits in \mathcal{C} for some epimorphism with finite kernel $f : G_1 \rightarrow H$.*

Proof. The exact sequence ξ splits in $\underline{\mathcal{C}}$ if and only if there exists a $\underline{\mathcal{C}}$ -morphism $\underline{g} : G_2 \rightarrow G_1$ such that $\underline{g} \circ u = \text{id}$ in $\underline{\mathcal{C}}$. Equivalently, there exists a finite subgroup $G'_1 \subset G_1$ and a $\underline{\mathcal{C}}$ -morphism $g : G_2 \rightarrow G_1/G'_1$ such that $g \circ u$ is the quotient map $f_1 : G_1 \rightarrow G_1/G'_1$.

If such a pair (G'_1, g) exists, then g factors through a \mathcal{C} -morphism $G_2/u(G'_1) \rightarrow G_1/G'_1$, which splits the bottom exact sequence in the push-out diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u} & G_2 & \xrightarrow{v} & G_3 & \longrightarrow & 0 \\ & & f_1 \downarrow & & f_2 \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & G_1/G'_1 & \xrightarrow{u'} & G_2/u(G'_1) & \xrightarrow{v'} & G_3 & \longrightarrow & 0. \end{array}$$

Replacing G'_1 by a larger finite subgroup, we may assume that G_1/G'_1 is smooth and connected (Lemma 2.2).

Conversely, a splitting of the bottom exact sequence in the above diagram is given by a \mathcal{C} -morphism $g' : G_2/u(G'_1) \rightarrow G_1/G'_1$ such that $g' \circ u' = \text{id}$ in \mathcal{C} . Let $g : G_2 \rightarrow G_1/G'_1$ denote the composition $G_2 \xrightarrow{f_2} G_2/u(G'_1) \xrightarrow{g'} G_1/G'_1$. Then $g \circ u = g' \circ f_2 \circ u = g' \circ u' \circ f_1 = f_1$ as desired. \square

We may now construct non-split exact sequences in $\underline{\mathcal{C}}$, thereby showing that $\text{hd}(\underline{\mathcal{C}}) \geq 1$:

EXAMPLES 3.4. (i) Consider an exact sequence

$$\xi : 0 \longrightarrow \mathbb{G}_a \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where A is an abelian variety. Then ξ , viewed as an extension of A by \mathbb{G}_a in \mathcal{C} , is classified by an element $\eta \in H^1(A, \mathcal{O}_A)$ (see [Ro58] or [MM74, §1.9]).

If $\text{char}(k) = 0$, then every epimorphism with finite kernel $f : \mathbb{G}_a \rightarrow H$ may be identified with the multiplication by some $t \in k^*$, viewed as an endomorphism of \mathbb{G}_a ; then the push-out $f_*\xi$ is classified by $t\eta$. In view of Lemma 3.3, it follows that ξ is non-split in $\underline{\mathcal{C}}$ whenever $\eta \neq 0$.

In contrast, if $\text{char}(k) = p > 0$, then ξ splits in $\underline{\mathcal{C}}$. Indeed, the multiplication map p_A yields an isomorphism in $\underline{\mathcal{C}}$, and $p_{\mathbb{G}_a} = 0$ whereas $(p_{\mathbb{G}_a})_*\xi = p_A^*\xi$.

(ii) Assume that $\text{char}(k) = p > 0$ and consider the algebraic group W_2 of Witt vectors of length 2. This group comes with an exact sequence

$$\xi : 0 \longrightarrow \mathbb{G}_a \longrightarrow W_2 \longrightarrow \mathbb{G}_a \longrightarrow 0,$$

see e.g. [DG70, §V.1.1.6]. Every epimorphism with finite kernel $f : \mathbb{G}_a \rightarrow H$ may be identified with a non-zero endomorphism of \mathbb{G}_a . In view of [DG70, Cor. V.1.5.2], it follows that the push-forward $f_*\xi$ is non-split. Thus, ξ does not split in $\underline{\mathcal{C}}$.

PROPOSITION 3.5. Consider an exact sequence

$$0 \longrightarrow G_1 \xrightarrow{u_1} G_2 \xrightarrow{u_2} \dots \xrightarrow{u_{n-1}} G_n \longrightarrow 0$$

in $\underline{\mathcal{C}}$. Then there exists an exact sequence

$$0 \longrightarrow H_1 \xrightarrow{v_1} H_2 \xrightarrow{v_2} \dots \xrightarrow{v_{n-1}} H_n \longrightarrow 0$$

in \mathcal{C} , and epimorphisms with finite kernels $f_i : G_i \rightarrow H_i$ ($i = 1, 2, \dots, n$), such that the diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u_1} & G_2 & \xrightarrow{u_2} & \dots & \xrightarrow{u_{n-1}} & G_n & \longrightarrow & 0 \\ & & \downarrow f_1 & & \downarrow f_2 & & & & \downarrow f_n & & \\ 0 & \longrightarrow & H_1 & \xrightarrow{v_1} & H_2 & \xrightarrow{v_2} & \dots & \xrightarrow{v_{n-1}} & H_n & \longrightarrow & 0 \end{array}$$

commutes in $\underline{\mathcal{C}}$.

Proof. We argue by induction on the length n . If $n = 2$, then we just have an isomorphism $\underline{u} : G_1 \rightarrow G_2$ in $\underline{\mathcal{C}}$. Then \underline{u} is represented by an isomorphism $u : G_1/G'_1 \rightarrow G_2/G'_2$ in \mathcal{C} , for some finite groups $G'_1 \subset G_1$ and $G'_2 \subset G_2$. If $n = 3$, then the $\underline{\mathcal{C}}$ -morphism $\underline{u}_2 : G_2 \rightarrow G_3$ is represented by an epimorphism $u_2 : G_2 \rightarrow G_3/G'_3$ in \mathcal{C} , where G'_3 is a finite subgroup of G_3 . We may thus replace G_3 with G_3/G'_3 , and assume that u_2 is an epimorphism in \mathcal{C} . Next, $\underline{u}_1 : G_1 \rightarrow G_2$ is represented by a morphism $u_1 : G_1 \rightarrow G_2/G'_2$ with finite kernel, where G'_2 is a finite subgroup of G_2 . We may thus replace G_1 (resp. G_2, G_3) with $G_1/\text{Ker}(u_1)$ (resp. $G_2/G'_2, G_3/u_2(G'_2)$) and assume that u_1 is a monomorphism in \mathcal{C} . Then $u_2 \circ u_1$ has finite image, and hence is zero since G_1 is smooth and connected. We now have a complex in \mathcal{C}

$$0 \longrightarrow G_1 \xrightarrow{u_1} G_2 \xrightarrow{u_2} G_3 \longrightarrow 0,$$

where u_1 is a monomorphism, u_2 an epimorphism, and $\text{Ker}(u_2)/\text{Im}(u_1)$ is finite. By Lemma 2.2, we may choose a finite subgroup $F \subset \text{Ker}(u_2)$ such that $\text{Ker}(u_2) = \text{Im}(u_1) + F$. This yields a commutative diagram in \mathcal{C}

$$\begin{array}{ccccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u_1} & G_2 & \xrightarrow{u_2} & G_3 & \longrightarrow & 0 \\ & & \downarrow q_1 & & \downarrow q_2 & & \downarrow \text{id} & & \\ 0 & \longrightarrow & G_1/u_1^{-1}(F) & \xrightarrow{v_1} & G_2/F & \xrightarrow{v_2} & G_3 & \longrightarrow & 0, \end{array}$$

where q_1, q_2 denote the quotient maps. Clearly, v_1 is a monomorphism, and v_2 an epimorphism. Also, $v_2 \circ v_1 = 0$, since q_1 is an epimorphism. Finally, q_2 restricts to an epimorphism $\text{Ker}(u_2) \rightarrow \text{Ker}(v_2)$, and hence $\text{Ker}(v_2) = \text{Im}(v_1)$. This completes the proof in the case where $n = 3$.

For an arbitrary length $n \geq 4$, we cut the given exact sequence into two exact sequences in $\underline{\mathcal{C}}$

$$\begin{array}{l} 0 \longrightarrow G_1 \xrightarrow{u_1} G_2 \longrightarrow K \longrightarrow 0, \\ 0 \longrightarrow K \longrightarrow G_3 \xrightarrow{u_3} \dots \xrightarrow{u_{n-1}} G_n \longrightarrow 0. \end{array}$$

By the induction assumption, there exists a commutative diagram in $\underline{\mathcal{C}}$

$$\begin{array}{ccccccc} 0 & \longrightarrow & K & \longrightarrow & G_3 & \xrightarrow{u_3} & \cdots \xrightarrow{u_{n-1}} & G_n & \longrightarrow & 0 \\ & & \underline{f} \downarrow & & \underline{f}_3 \downarrow & & & \underline{f}_n \downarrow & & \\ 0 & \longrightarrow & L & \xrightarrow{v} & H_3 & \xrightarrow{v_3} & \cdots \xrightarrow{v_{n-1}} & H_n & \longrightarrow & 0, \end{array}$$

where f, f_3, \dots, f_n are epimorphisms with finite kernels, and the bottom sequence comes from an exact sequence in \mathcal{C} . Since \underline{f} is an isomorphism in $\underline{\mathcal{C}}$, we have an exact sequence

$$0 \longrightarrow G_1 \xrightarrow{u_1} G_2 \longrightarrow L \longrightarrow 0$$

in $\underline{\mathcal{C}}$, and hence another commutative diagram in $\underline{\mathcal{C}}$

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u_1} & G_2 & \longrightarrow & L & \longrightarrow & 0 \\ & & \underline{f}_1 \downarrow & & \underline{f}_2 \downarrow & & \underline{g} \downarrow & & \\ 0 & \longrightarrow & H_1 & \xrightarrow{v_1} & H_2 & \longrightarrow & M & \longrightarrow & 0, \end{array}$$

where again f_1, f_2, g are epimorphisms with finite kernels, and the bottom sequence comes from an exact sequence in \mathcal{C} . Let $H'_3 := H_3/v(F)$, and denote by F the kernel of $g : L \rightarrow M$; then we have a commutative diagram in $\underline{\mathcal{C}}$

$$\begin{array}{ccccccc} 0 & \longrightarrow & L & \longrightarrow & G_3 & \xrightarrow{u_3} & G_4 & \xrightarrow{u_4} & \cdots \xrightarrow{u_{n-1}} & G_n & \longrightarrow & 0 \\ & & \underline{g} \downarrow & & \underline{h} \downarrow & & \underline{f}_4 \downarrow & & & \underline{f}_n \downarrow & & \\ 0 & \longrightarrow & M & \longrightarrow & H'_3 & \xrightarrow{w} & H_4 & \xrightarrow{v_4} & \cdots \xrightarrow{v_{n-1}} & H_n & \longrightarrow & 0, \end{array}$$

satisfying similar properties. By concatenating the two latter diagrams, we obtain the desired commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u_1} & G_2 & \xrightarrow{u_2} & G_3 & \xrightarrow{u_3} & \cdots \xrightarrow{u_{n-1}} & G_n & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & & \downarrow & & \\ 0 & \longrightarrow & H_1 & \xrightarrow{v_1} & H_2 & \longrightarrow & H'_3 & \xrightarrow{w} & \cdots \xrightarrow{v_{n-1}} & H_n & \longrightarrow & 0. \end{array}$$

□

3.2 DIVISIBLE GROUPS

Given a divisible algebraic group G and a non-zero integer n , the morphism $n_G : G \rightarrow G$ factors through an isomorphism $G/G[n] \xrightarrow{\cong} G$. We denote the inverse isomorphism by

$$u_n : G \xrightarrow{\cong} G/G[n].$$

By construction, we have a commutative triangle

$$\begin{array}{ccc}
 G & & \\
 n_G \downarrow & \searrow q & \\
 G & \xrightarrow{u_n} & G/G[n],
 \end{array}$$

where q denotes the quotient morphism. Since q yields the identity morphism in $\underline{\mathcal{C}}$, we see that u_n yields the inverse of the $\underline{\mathcal{C}}$ -automorphism n_G of G . As a consequence, $\text{End}_{\underline{\mathcal{C}}}(G)$ is a \mathbb{Q} -algebra.

More generally, we have the following:

PROPOSITION 3.6. *Let G, H be smooth connected algebraic groups, and assume that H is divisible.*

- (i) *Every extension group $\text{Ext}_{\underline{\mathcal{C}}}^n(G, H)$ is a \mathbb{Q} -vector space.*
- (ii) *The natural map $Q : \text{Hom}_{\mathcal{C}}(G, H) \rightarrow \text{Hom}_{\underline{\mathcal{C}}}(G, H)$ is injective and induces an isomorphism*

$$\gamma : \mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{C}}(G, H) \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(G, H), \quad \frac{1}{n} \otimes f \longmapsto Q(u_n \otimes f).$$

- (iii) *If G is divisible as well, then the natural map*

$$Q^1 : \text{Ext}_{\underline{\mathcal{C}}}^1(G, H) \rightarrow \text{Ext}_{\underline{\mathcal{C}}}^1(G, H)$$

induces an isomorphism

$$\gamma^1 : \mathbb{Q} \otimes_{\mathbb{Z}} \text{Ext}_{\underline{\mathcal{C}}}^1(G, H) \longrightarrow \text{Ext}_{\underline{\mathcal{C}}}^1(G, H).$$

Proof. (i) Just note that $\text{Ext}_{\underline{\mathcal{C}}}^n(G, H)$ is a module over the \mathbb{Q} -algebra $\text{End}_{\underline{\mathcal{C}}}(H)$.

(ii) Let $f \in \text{Hom}_{\mathcal{C}}(G, H)$, and n a positive integer. If $\gamma(\frac{1}{n} \otimes f) = 0$, then $\gamma(f) = 0$, i.e., $\underline{f} = 0$. Thus, $f = 0$ in view of Lemma 3.1; this shows the injectivity of γ .

For the surjectivity, consider a $\underline{\mathcal{C}}$ -morphism $\underline{f} : G \rightarrow H$ represented by a \mathcal{C} -morphism $f : G \rightarrow H/H'$, where H' is a finite subgroup of H . Then $H' \subset H[n]$ for some positive integer n , which we may take to be the order of H' . Thus, we may assume that $H' = H[n]$. Then the \mathcal{C} -morphism $\varphi := u_n^{-1} \circ f : G \rightarrow H$ satisfies $\underline{\varphi} = n\underline{f}$, i.e., $\underline{f} = \gamma(\frac{1}{n} \otimes \varphi)$.

(iii) Consider $\eta \in \text{Ext}_{\underline{\mathcal{C}}}^1(G, H)$ such that $\gamma^1(\frac{1}{n} \otimes \eta) = 0$ for some positive integer n . Then of course $\gamma^1(\eta) = 0$, i.e., η is represented by an exact sequence in \mathcal{C}

$$0 \longrightarrow H \xrightarrow{u} E \xrightarrow{v} G \longrightarrow 0,$$

which splits in $\underline{\mathcal{C}}$. By Lemma 3.3 and the divisibility of H , it follows that the push-out by m_H of the above extension splits in \mathcal{C} for some $m > 0$. But

$(m_H)_*\eta = m\eta$ (see e.g. [Oo66, Lem. I.3.1]), and hence $m\eta = 0$. This shows the injectivity of γ^1 .

For the surjectivity, we adapt the proof of Proposition 3.5. Let $\eta \in \text{Ext}_{\mathcal{C}}^1(G, H)$ be represented by an exact sequence in \mathcal{C}

$$0 \longrightarrow H \xrightarrow{u} E \xrightarrow{v} G \longrightarrow 0.$$

Since G is divisible, v is represented by a \mathcal{C} -morphism $v : E \rightarrow G/G[m]$ for some positive integer m . Replacing η with its pull-back $u_m^*\eta = (m_G^*)^{-1}\eta = \gamma^1(\frac{1}{m}\otimes\eta)$, we may thus assume that v is represented by a \mathcal{C} -epimorphism $v : E \rightarrow G$.

Likewise, as H is divisible, u is represented by a \mathcal{C} -morphism $u : H \rightarrow E/E[n]$. Then η is represented by the exact sequence in \mathcal{C}

$$0 \longrightarrow H \xrightarrow{u} E/E[n] \xrightarrow{v_n} G \longrightarrow 0,$$

where $v_n : E/E[n] \rightarrow G/G[n]$ is the \mathcal{C} -epimorphism induced by v . So we may further assume that u is represented by a \mathcal{C} -morphism $u : H \rightarrow E$. By Lemma 3.1, we then have $v \circ u = 0$; moreover, $\text{Ker}(u)$ and $\text{Ker}(v)/\text{Im}(u)$ are finite. In view of Lemma 2.2, we have $\text{Ker}(v) = \text{Im}(u) + E'$ for some finite subgroup $E' \subset E$. This yields a commutative diagram in \mathcal{C}

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H & \xrightarrow{u} & E & \xrightarrow{v} & G & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & H/u^{-1}(E') & \xrightarrow{u_1} & E/E' & \xrightarrow{v_1} & G & \longrightarrow & 0, \end{array}$$

where the bottom sequence is exact, and $u^{-1}(E')$ is finite.

We may thus choose a positive integer r such that $u^{-1}(E') \subset H[r]$. Taking the push-out by the quotient map $H/u^{-1}(E') \rightarrow H/H[r]$ yields a commutative diagram in \mathcal{C}

$$\begin{array}{ccccccccc} 0 & \longrightarrow & H & \xrightarrow{u} & E & \xrightarrow{v} & G & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & H/H[r] & \xrightarrow{u_2} & E/E' + u(H[r]) & \xrightarrow{v_2} & G & \longrightarrow & 0, \end{array}$$

where the bottom sequence is exact again. Thus, $r\eta = (r_H)_*\eta$ is represented by an exact sequence in \mathcal{C} . □

REMARKS 3.7. (i) Given two divisible groups G, H , the map

$$Q^1 : \text{Ext}_{\mathcal{C}}^1(G, H) \longrightarrow \text{Ext}_{\mathcal{C}}^1(G, H)$$

is not necessarily injective. Indeed, the group $\text{Ext}_{\mathcal{C}}^1(A, \mathbb{G}_m)$ has non-zero torsion for any non-zero abelian variety A over (say) a separably closed field, as follows from the Weil-Barsotti isomorphism.

(ii) We may also consider the natural maps

$$\gamma^n : \mathbb{Q} \otimes_{\mathbb{Z}} \text{Ext}_{\mathcal{C}}^n(G, H) \longrightarrow \text{Ext}_{\underline{\mathcal{C}}}^n(G, H)$$

for $n \geq 2$. But these maps turn out to be zero for any algebraic groups G, H , since $\text{Ext}_{\underline{\mathcal{C}}}^n(G, H) = 0$ (Lemma 4.11) and $\text{Ext}_{\mathcal{C}}^n(G, H)$ is torsion (Remark 4.12).

As a first application of Proposition 3.6, we obtain:

PROPOSITION 3.8. *Assume that $\text{char}(k) = 0$.*

(i) *The composition of the inclusion $\mathcal{U} \rightarrow \mathcal{C}$ with the quotient $Q : \mathcal{C} \rightarrow \underline{\mathcal{C}}$ identifies \mathcal{U} with a full subcategory of $\underline{\mathcal{C}}$.*

(ii) *The unipotent radical functor yields an exact functor*

$$\underline{R}_u : \underline{\mathcal{C}} \longrightarrow \mathcal{U},$$

which is right adjoint to the inclusion. Moreover, \underline{R}_u commutes with base change under field extensions.

(iii) *Every unipotent group is a projective object in $\underline{\mathcal{C}}$.*

Proof. (i) Recall that a morphism of unipotent groups is just a linear map of the associated k -vector spaces. In view of Proposition 3.6, it follows that the natural map $\text{Hom}_{\mathcal{C}}(U, V) \rightarrow \text{Hom}_{\underline{\mathcal{C}}}(U, V)$ is an isomorphism for any unipotent groups U, V .

(ii) The functor $R_u : \mathcal{C} \rightarrow \mathcal{U}$ is exact by Theorem 2.9, and sends every finite group to 0. By the universal property of Q , there exists a unique exact functor $S : \mathcal{C}/\mathcal{F} \rightarrow \mathcal{U}$ such that $R_u = S \circ Q$. Since R_u commutes with base change under field extensions (Theorem 2.9 again), so does S by uniqueness. Thus, composing S with the inclusion $\underline{\mathcal{C}} \rightarrow \mathcal{C}/\mathcal{F}$ yields the desired exact functor.

For any unipotent group U and any algebraic group G , the natural map

$$\text{Hom}_{\mathcal{U}}(U, R_u(G)) \longrightarrow \text{Hom}_{\mathcal{C}}(U, G)$$

is an isomorphism. By Proposition 3.6 again, the natural map

$$Q : \text{Hom}_{\mathcal{C}}(U, G) \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(U, G)$$

is an isomorphism as well. It follows that \underline{R}_u is right adjoint to the inclusion.

(iii) Let U be a unipotent group. Then the functor on \mathcal{C} defined by

$$G \longmapsto \text{Hom}_{\mathcal{U}}(U, R_u(G))$$

is exact, since the unipotent radical functor is exact and the category \mathcal{U} is semi-simple. Thus, $G \mapsto \text{Hom}_{\underline{\mathcal{C}}}(U, G)$ is exact as well; this yields the assertion. \square

3.3 FIELD EXTENSIONS

Let k' be a field extension of k . Then the assignment $G \mapsto G_{k'} := G \otimes_k k'$ yields the *base change functor*

$$\otimes_k k' : \mathcal{C}_k \longrightarrow \mathcal{C}_{k'}.$$

Clearly, this functor is exact and faithful; also, note that an algebraic group G is connected (resp. smooth, finite, linear, unipotent, a torus, an abelian variety, a semi-abelian variety) if and only if so is $G_{k'}$.

LEMMA 3.9. *With the above notation, the functor $\otimes_k k'$ yields an exact functor*

$$\underline{\otimes}_k k' : \underline{\mathcal{C}}_k \longrightarrow \underline{\mathcal{C}}_{k'}.$$

Proof. As the composite functor $Q_{k'} \circ \otimes_k k' : \mathcal{C}_k \rightarrow \underline{\mathcal{C}}_{k'}$ is exact and sends every finite k -group to 0, it factors through a unique exact functor $\mathcal{C}_k/\mathcal{F}_k \rightarrow \underline{\mathcal{C}}_{k'}$. This yields the existence and exactness of $\underline{\otimes}_k k'$. \square

LEMMA 3.10. *Let k' be a purely inseparable field extension of k , and G' a k' -group.*

- (i) *There exists a smooth k -group G and an epimorphism $f : G' \rightarrow G_{k'}$ such that $\text{Ker}(f)$ is infinitesimal.*
- (ii) *If $G' \subset H_{k'}$ for some k -group H , then there exists a k -subgroup $G \subset H$ such that $G' \subset G_{k'}$ and $G_{k'}/G'$ is infinitesimal.*

Proof. (i) Let n be a positive integer and consider the n th relative Frobenius morphism

$$F_{G'/k'}^n : G' \longrightarrow G'^{(p^n)}.$$

Recall that the quotient $G'/\text{Ker}(F_{G'/k'}^n)$ is smooth for $n \gg 0$. Since $\text{Ker}(F_{G'/k'}^n)$ is infinitesimal, we may assume that G' is smooth. Then $F_{G'/k'}^n$ is an epimorphism in view of [SGA3, VIIA, Cor. 8.3.1].

Next, note that G' is defined over some finite subextension k'' of k' , i.e., there exists a k'' -subgroup G'' such that $G' = G'' \otimes_{k''} k'$. By transitivity of base change, we may thus assume that k' is finite over k . Let $q := [k' : k]$, then $q = p^n$, where $p = \text{char}(k)$ and n is a positive integer; also, $k'^q \subset k$. Consider again the morphism $F_{G'/k'}^n$; then by construction, $G'^{(p^n)} \cong G' \otimes_{k'} k'$, where k' is sent to itself via the q th power map. Thus, $G'^{(p^n)} \cong G_{k'}$, where G denotes the k -group $G' \otimes_{k'} k$; here k' is sent to k via the q th power map again. So the induced map $G' \rightarrow G_{k'}$ is the desired morphism.

(ii) As above, we may reduce to the case where k' is finite over k . Then the statement follows by similar arguments, see [Br17, Lem. 4.3.5] for details. \square

We now are ready to prove Theorem 1 (iv):

THEOREM 3.11. *Let k' be a purely inseparable field extension of k . Then the base change functor $\underline{\otimes}_k k' : \underline{\mathcal{C}}_k \rightarrow \underline{\mathcal{C}}_{k'}$ is an equivalence of categories.*

Proof. By Lemma 3.10, every k' -group G' is isogenous to $G_{k'}$ for some smooth k -group G . It follows that $\underline{\otimes}_k k'$ is essentially surjective.

Next, let G, H be smooth connected k -groups, and $\underline{f} : G \rightarrow H$ a $\underline{\mathcal{C}}_k$ -morphism, represented by a \mathcal{C}_k -morphism $f : G \rightarrow H/H'$ for some finite k -subgroup H' of H . If $\underline{f}_{k'} : G_{k'} \rightarrow H_{k'}$ is zero in $\underline{\mathcal{C}}_{k'}$, then the image of $f_{k'} : G_{k'} \rightarrow H_{k'}/H'_{k'}$ is finite. By Lemma 2.1, it follows that $f_{k'} = 0$. This shows that $\underline{\otimes}_k k'$ is faithful. We now check that $\underline{\otimes}_k k'$ is full. Let again G, H be smooth connected k -groups, and let $\underline{f} \in \text{Hom}_{\underline{\mathcal{C}}_{k'}}(G_{k'}, H_{k'})$. We show that there exists a finite k -subgroup $H' \subset H$ and a k -morphism $\varphi : G \rightarrow H/H'$ such that $\varphi_{k'}$ represents \underline{f} . For this, we may replace H with its quotient by any finite k -subgroup.

Choose a representative $f : G_{k'} \rightarrow H_{k'}/H''$ of \underline{f} , where $H'' \subset H_{k'}$ is a finite k' -subgroup. By Lemma 3.10, there exists a \bar{k} -subgroup $I \subset H$ such that $H'' \subset I_{k'}$ and $I_{k'}/H''$ is finite; then $I_{k'}$ is finite as well, and hence so is I . We may thus replace H by H/I , and f by its composition with the quotient morphism $H_{k'}/H'' \rightarrow H_{k'}/I_{k'} = (H/I)_{k'}$. Then \underline{f} is represented by a morphism $f : G_{k'} \rightarrow H_{k'}$.

Consider the graph $\Gamma(f) \subset G_{k'} \times_{k'} H_{k'}$. By Lemma 3.10 again, there exists a k -subgroup $\Delta \subset G \times H$ such that $\Gamma(f) \subset \Delta_{k'}$ and $\Delta_{k'}/\Gamma(f)$ is finite. Then the intersection $\Delta_{k'} \cap (0 \times H_{k'})$ is finite, since $\Gamma(f) \cap (0 \times H_{k'})$ is zero. Thus, $\Delta \cap (0 \times H)$ is finite as well; equivalently, the k -group $H' := H \cap (0 \times \text{id})^{-1}(\Delta)$ is finite. Denoting by Γ the image of Δ in $G \times H/H'$, we have a cartesian square

$$\begin{array}{ccc} \Delta & \longrightarrow & G \times H \\ \downarrow & & \downarrow \\ \Gamma & \longrightarrow & G \times H/H', \end{array}$$

where the horizontal arrows are closed immersions, and the left (resp. right) vertical arrow is the quotient by $\Delta \cap (0 \times H)$ (resp. by H' acting on H via addition). So Γ is a k -subgroup of $G \times H/H'$, and $\Gamma \cap (0 \times H/H')$ is zero; in other words, the projection $\pi : \Gamma \rightarrow G$ is a closed immersion. Since G is smooth and connected, and $\dim(\Gamma) = \dim(\Delta) = \dim \Gamma(f) = \dim(G)$, it follows that π is an isomorphism. In other words, Γ is the graph of a k -morphism $\varphi : G \rightarrow H/H'$. Since the above cartesian square lies in a push-out diagram,

$$\begin{array}{ccccccc} 0 & \longrightarrow & \Delta & \longrightarrow & G \times H & \xrightarrow{\varphi - q} & H/H' \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow \\ 0 & \longrightarrow & \Gamma & \longrightarrow & G \times H/H' & \xrightarrow{\varphi - \text{id}} & H/H' \longrightarrow 0, \end{array}$$

where $q : H \rightarrow H/H'$ denotes the quotient morphism, it follows that $\Delta = \ker(\varphi - q)$. As $\Gamma(f) \subset \Delta_{k'}$, we see that $\varphi_{k'} = q_{k'} \circ f$. Thus, \underline{f} is represented by $\varphi_{k'}$; this completes the proof of the fullness assertion. \square

REMARKS 3.12. (i) Likewise, the base change functor induces equivalences of categories $\underline{\mathcal{U}}_k \rightarrow \underline{\mathcal{U}}_{k'}$, $\underline{\mathcal{T}}_k \rightarrow \underline{\mathcal{T}}_{k'}$, $\underline{\mathcal{L}}_k \rightarrow \underline{\mathcal{L}}_{k'}$, and $\underline{\mathcal{A}}_k \rightarrow \underline{\mathcal{A}}_{k'}$. For tori, this follows much more directly from the anti-equivalence of $\underline{\mathcal{T}}_k$ with the category of rational representations of the absolute Galois group, see Proposition 4.1.

(ii) In particular, the category $\underline{\mathcal{U}}_k$ is equivalent to $\underline{\mathcal{U}}_{k_i}$, where k_i denotes the perfect closure of k in \bar{k} . Recall from [DG70, Thm. V.1.4.3, Cor. V.1.4.4] that the category \mathcal{U}_{k_i} is anti-equivalent to the category of finitely generated modules over the Dieudonné ring $\mathbb{D} = \mathbb{D}_{k_i}$ which are killed by some power of the Verschiebung map V . Moreover, the category $\underline{\mathcal{U}}_{k_i}$ is anti-equivalent to the category of finitely generated modules over the localization $\mathbb{D}_{(V)}$ which are killed by some power of V ; see [DG70, §V.3.6.7].

By work of Schoeller, Kraft, and Takeuchi (see [Sc72, Kr75, Tak75]), the category \mathcal{U}_k is anti-equivalent to a category of finitely generated modules over a certain k -algebra, which generalizes the Dieudonné ring but seems much less tractable.

4 TORI, ABELIAN VARIETIES, AND HOMOLOGICAL DIMENSION

4.1 TORI

Denote by \mathcal{T} (resp. \mathcal{M} , \mathcal{FM}) the full subcategory of \mathcal{C} with objects the tori (resp. the groups of multiplicative type, the finite groups of multiplicative type). Then \mathcal{T} is stable under taking quotients and extensions, but not subobjects; in particular, \mathcal{T} is an additive subcategory of \mathcal{C} , but not an abelian subcategory. Also, \mathcal{F} and \mathcal{FM} are stable under taking subobjects, quotients and extensions. Thus, we may form the quotient abelian category \mathcal{M}/\mathcal{FM} , as in §3.1. One may readily check that \mathcal{M}/\mathcal{FM} is a full subcategory of \mathcal{C}/\mathcal{F} .

Let $\underline{\mathcal{T}}$ be the full subcategory of \mathcal{M}/\mathcal{FM} with objects the tori. Since these are the smooth connected objects of \mathcal{M} , one may check as in Lemma 3.1 that the inclusion of $\underline{\mathcal{T}}$ in \mathcal{M}/\mathcal{FM} is an equivalence of categories. The remaining statements of Lemma 3.1 also adapt to this setting; note that we may replace the direct limits over all finite subgroups with those over all n -torsion subgroups, since tori are divisible. Also, Proposition 3.6 yields natural isomorphisms

$$\mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(T_1, T_2) \xrightarrow{\cong} \mathrm{Hom}_{\underline{\mathcal{T}}}(T_1, T_2)$$

for any tori T_1, T_2 .

By assigning with each group of multiplicative type G its character group,

$$X(G) := \mathrm{Hom}_{k_s}(G_{k_s}, \mathbb{G}_{m, k_s}),$$

one obtains an anti-equivalence between \mathcal{M} (resp. \mathcal{FM}) and the category of finitely generated (resp. finite) abstract commutative groups equipped with the discrete topology and a continuous action of the Galois group Γ ; see [DG70, Thm. IV.1.3.6]. Thus, the assignment

$$X_{\mathbb{Q}} : G \longmapsto \mathbb{Q} \otimes_{\mathbb{Z}} X(G) =: X(G)_{\mathbb{Q}}$$

yields a contravariant exact functor from \mathcal{M} to the category $\text{Rep}_{\mathbb{Q}}(\Gamma)$ of finite-dimensional \mathbb{Q} -vector spaces equipped with a continuous representation of Γ as above; moreover, every finite group of multiplicative type is sent to 0. This yields in turn a contravariant exact functor

$$\underline{X}_{\mathbb{Q}} : \underline{\mathcal{T}} \longrightarrow \text{Rep}_{\mathbb{Q}}(\Gamma).$$

PROPOSITION 4.1. *The functor $\underline{X}_{\mathbb{Q}}$ is an anti-equivalence of categories. In particular, the category $\underline{\mathcal{T}}$ is semi-simple, and $\text{Hom}_{\underline{\mathcal{T}}}(T_1, T_2)$ is a finite-dimensional \mathbb{Q} -vector space for any tori T_1, T_2 .*

Proof. Given a finite-dimensional \mathbb{Q} -vector space V equipped with a continuous action of Γ , there exists a finitely generated Γ -stable subgroup $M \subset V$ which spans V ; thus, $V \cong X(T)_{\mathbb{Q}}$, where T denotes the torus with character group M . So $\underline{X}_{\mathbb{Q}}$ is essentially surjective.

Given two tori T_1, T_2 , the natural isomorphism

$$\text{Hom}_{\mathcal{M}}(T_1, T_2) \cong \text{Hom}^{\Gamma}(X(T_2), X(T_1))$$

yields an isomorphism

$$\text{Hom}_{\underline{\mathcal{T}}}(T_1, T_2) \cong \text{Hom}^{\Gamma}(X(T_2)_{\mathbb{Q}}, X(T_1)_{\mathbb{Q}})$$

in view of Proposition 3.6. It follows that $\underline{X}_{\mathbb{Q}}$ is fully faithful. □

- LEMMA 4.2. (i) *Every algebraic group G has a unique maximal torus, $T(G)$.*
 (ii) *Every morphism of algebraic groups $u : G \rightarrow H$ sends $T(G)$ to $T(H)$.*
 (iii) *The formation of $T(G)$ commutes with base change under field extensions.*

Proof. (i) This follows from the fact that $T_1 + T_2$ is a torus for any subtori $T_1, T_2 \subset G$.

(ii) Just note that the image of a torus under any morphism is still a torus.

(iii) Consider an algebraic group G , its maximal torus T , and a field extension k' of k . If $\text{char}(k) = 0$, then Theorem 2.9 implies that G/T is an extension of an abelian variety by a product $M \times U$, where M is finite and U unipotent. As a consequence, a similar assertion holds for $G_{k'}/T_{k'}$; it follows that $G_{k'}/T_{k'}$ contains no non-zero torus, and hence $T_{k'}$ is the maximal torus of $G_{k'}$. On the other hand, if $\text{char}(k) > 0$, then G/T is a 3-step extension of a unipotent group by an abelian variety by a finite group, in view of Theorem 2.11. It follows similarly that $T_{k'}$ is the maximal torus of $G_{k'}$. □

By Lemma 4.2, the assignment $G \mapsto T(G)$ yields a functor

$$T : \mathcal{C} \longrightarrow \mathcal{T},$$

the *functor of maximal tori*. This functor is not exact, as seen from the exact sequence

$$0 \longrightarrow G[n] \longrightarrow G \xrightarrow{n\mathcal{C}} G \longrightarrow 0,$$

where G is a non-zero torus and n a non-zero integer. But T is exact up to finite groups, as shown by the following;

LEMMA 4.3. *Every exact sequence in \mathcal{C}*

$$0 \longrightarrow G_1 \xrightarrow{u} G_2 \xrightarrow{v} G_3 \longrightarrow 0$$

yields a complex in \mathcal{C}

$$0 \longrightarrow T(G_1) \xrightarrow{T(u)} T(G_2) \xrightarrow{T(v)} T(G_3) \longrightarrow 0,$$

where $T(u)$ is a monomorphism, $T(v)$ an epimorphism, and $\text{Ker } T(v)/\text{Im } T(u)$ is finite.

Proof. We argue as in the proof of Theorem 2.11 (iii). Clearly, $T(u)$ is a monomorphism. Also, the group

$$\text{Ker } T(v)/\text{Im } T(u) = T(G_2) \cap u(G_1)/u(T(G_1))$$

is the quotient of a group of multiplicative type by its maximal torus, and hence is finite.

To show that $T(v)$ is an epimorphism, we may replace G_2 with $v^{-1}(T(G_3))$, and hence assume that G_3 is a torus. Next, we may replace G_2 with $G_2/T(G_2)$, and hence assume that $T(G_2)$ is zero. We then have to check that G_3 is zero. If $\text{char}(k) = 0$, then there is an exact sequence

$$0 \longrightarrow M \times U \longrightarrow G_2 \longrightarrow A \longrightarrow 0$$

as in Theorem 2.9, where M is finite. Thus, every morphism $G_2 \rightarrow G_3$ has finite image. Since $v : G_2 \rightarrow G_3$ is an epimorphism, it follows that $G_3 = 0$. On the other hand, if $\text{char}(k) > 0$, then there are exact sequences

$$0 \longrightarrow H \longrightarrow G_2 \longrightarrow U \longrightarrow 0, \quad 0 \longrightarrow M \longrightarrow H \longrightarrow A \longrightarrow 0$$

as in Theorem 2.11, where M is finite. This implies again that every morphism $G_2 \rightarrow G_3$ has finite image, and hence that $G_3 = 0$. \square

PROPOSITION 4.4. (i) *The functor of maximal tori yields an exact functor*

$$\underline{T} : \underline{\mathcal{C}} \longrightarrow \underline{\mathcal{T}},$$

right adjoint to the inclusion $\underline{\mathcal{T}} \rightarrow \underline{\mathcal{C}}$. Moreover, \underline{T} commutes with base change under field extensions.

(ii) *Every torus is a projective object in $\underline{\mathcal{C}}$.*

Proof. (i) Composing T with the functor $\mathcal{T} \rightarrow \underline{\mathcal{T}}$ induced by the quotient functor Q , we obtain an exact functor $\mathcal{C} \rightarrow \underline{\mathcal{T}}$ (Lemma 4.3), which sends every

finite group to 0. This yields an exact functor $\underline{T} : \underline{\mathcal{C}} \rightarrow \underline{\mathcal{T}}$. The adjointness assertion follows from the natural isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(T, G) \cong \mathrm{Hom}_{\mathcal{T}}(T, T(G))$$

for any torus T and any algebraic group G , which yields a natural isomorphism

$$\mathrm{Hom}_{\underline{\mathcal{C}}}(T, G) \cong \mathrm{Hom}_{\underline{\mathcal{T}}}(T, T(G))$$

by using Lemmas 3.1 and 4.3. Finally, the assertion on field extensions is a direct consequence of Lemma 4.2.

(ii) This follows by arguing as in the proof of Proposition 3.8 (iii). □

4.2 ABELIAN VARIETIES

Denote by \mathcal{A} (resp. \mathcal{P}) the full subcategory of \mathcal{C} with objects the abelian varieties (resp. the proper groups, i.e., those algebraic groups G such that the structure map $G \rightarrow \mathrm{Spec}(k)$ is proper). Like the category of tori, \mathcal{A} is stable under taking quotients and extensions, but not subobjects; so \mathcal{A} is an additive subcategory of \mathcal{C} , but not an abelian subcategory. Also, \mathcal{P} is stable under taking subobjects, quotients and extensions; it also contains the category \mathcal{F} of finite groups. We may thus form the quotient abelian category \mathcal{P}/\mathcal{F} , which is a full subcategory of \mathcal{C}/\mathcal{F} .

Next, let $\underline{\mathcal{A}}$ be the full subcategory of \mathcal{P}/\mathcal{F} with objects the abelian varieties. As in §4.1, the inclusion of $\underline{\mathcal{A}}$ in \mathcal{P}/\mathcal{F} is an equivalence of categories, and the remaining statements of Lemma 3.1 adapt to this setting. Also, Proposition 3.6 yields natural isomorphisms

$$\mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{C}}(A_1, A_2) \xrightarrow{\cong} \mathrm{Hom}_{\underline{\mathcal{A}}}(A_1, A_2)$$

for any abelian varieties A_1, A_2 . Since the abelian group $\mathrm{Hom}_{\mathcal{C}}(A_1, A_2)$ is free of finite rank (see [Mi86, Thm. 12.5]), $\mathrm{Hom}_{\underline{\mathcal{A}}}(A_1, A_2)$ is a finite-dimensional \mathbb{Q} -vector space. Moreover, the category $\underline{\mathcal{A}}$ is semi-simple, in view of the Poincaré complete reducibility theorem (which holds over an arbitrary field, see [Co06, Cor. 3.20] or [Br17, Cor. 4.2.6]).

LEMMA 4.5. (i) *Every smooth connected algebraic group G has a largest abelian variety quotient,*

$$\alpha = \alpha_G : G \longrightarrow A(G).$$

Moreover, $\mathrm{Ker}(\alpha)$ is linear and connected.

(ii) *Every morphism $u : G \rightarrow H$, where H is smooth and connected, induces a unique morphism $A(u) : A(G) \rightarrow A(H)$ such that the square*

$$\begin{array}{ccc} G & \xrightarrow{u} & H \\ \alpha_G \downarrow & & \alpha_H \downarrow \\ A(G) & \xrightarrow{A(u)} & A(H) \end{array}$$

commutes.

- (iii) For any field extension k' of k , the natural morphism $A(G_{k'}) \rightarrow A(G)_{k'}$ is an isomorphism if $\text{char}(k) = 0$, and an isogeny if $\text{char}(k) > 0$.

Proof. (i) and (ii) The assertions are direct consequences of Theorem 2.3 (ii). (iii) By (i), we have an exact sequence

$$0 \longrightarrow L(G) \longrightarrow G \xrightarrow{\alpha} A(G) \longrightarrow 0,$$

where $L(G)$ is linear and connected. This yields an exact sequence

$$0 \longrightarrow L(G)_{k'} \longrightarrow G_{k'} \longrightarrow A(G)_{k'} \longrightarrow 0.$$

Thus, $L(G_{k'}) \subset L(G)_{k'}$ and we obtain an exact sequence

$$0 \longrightarrow L(G)_{k'}/L(G_{k'}) \longrightarrow A(G_{k'}) \longrightarrow A(G)_{k'} \longrightarrow 0.$$

Since $L(G)_{k'}$ is linear, the quotient $L(G)_{k'}/L(G_{k'})$ must be finite; this yields the assertion when $\text{char}(k) > 0$.

When $\text{char}(k) = 0$, we may characterize $L(G)$ as the largest connected linear subgroup of G . It follows that $L(G)_{k'} \subset L(G_{k'})$; hence equality holds, and $A(G_{k'}) \xrightarrow{\cong} A(G)_{k'}$. \square

REMARKS 4.6. (i) An arbitrary algebraic group G may admit no largest abelian variety quotient, as shown by the following variant of [Br17, Ex. 4.3.8]: let A be a non-zero abelian variety, and choose an integer $n \geq 2$. Let

$$G := (A \times A[n^2])/\text{diag}(A[n]),$$

where $A[n]$ is viewed as a subgroup of $A[n^2]$. Consider the subgroups of G

$$H_1 := (A[n] \times A[n^2])/\text{diag}(A[n]), \quad H_2 := \text{diag}(A[n^2])/\text{diag}(A[n]).$$

Then G/H_1 , G/H_2 are both isomorphic to A . Also, $H_1 \cap H_2 = 0$, and G is not an abelian variety.

(ii) The assignment $G \mapsto A(G)$ does not preserve exactness of sequences of smooth connected algebraic groups. For example, consider an elliptic curve E equipped with a k -rational point of prime order $\ell \geq 2$. Assume that k contains a nontrivial ℓ th root of unity; this identifies μ_ℓ with the constant group scheme $\mathbb{Z}/\ell\mathbb{Z}$. Consider the quotient

$$G := (E \times \mathbb{G}_m)/\text{diag}(\mu_\ell),$$

with an obvious notation. Then G is a smooth connected algebraic group, which lies in an exact sequence

$$0 \longrightarrow E \longrightarrow G \longrightarrow \mathbb{G}_m \longrightarrow 0.$$

Moreover, the induced map $A(E) \rightarrow A(G)$ is just the quotient map $E \rightarrow E/\mu_\ell$.

We now show that the assignment $G \mapsto A(G)$ is exact up to finite groups:

LEMMA 4.7. *Consider an exact sequence in \mathcal{C}*

$$0 \longrightarrow G_1 \xrightarrow{u} G_2 \xrightarrow{v} G_3 \longrightarrow 0,$$

where G_1, G_2, G_3 are smooth and connected. Then we have a commutative diagram in \mathcal{C}

$$\begin{array}{ccccccccc} 0 & \longrightarrow & G_1 & \xrightarrow{u} & G_2 & \xrightarrow{v} & G_3 & \longrightarrow & 0 \\ & & \alpha_1 \downarrow & & \alpha_2 \downarrow & & \alpha_3 \downarrow & & \\ 0 & \longrightarrow & A(G_1) & \xrightarrow{A(u)} & A(G_2) & \xrightarrow{A(v)} & A(G_3) & \longrightarrow & 0, \end{array}$$

where $A(v)$ is an epimorphism, and $\text{Ker } A(u), \text{Ker } A(v)/\text{Im } A(u)$ are finite.

Proof. Clearly, $A(v)$ is an epimorphism. Let $L_i := \text{Ker}(\alpha_i)$ for $i = 1, 2, 3$; then each L_i is connected and linear by Theorem 2.3. We have isomorphisms $\text{Ker } A(u) \cong u^{-1}(L_2)/L_1$, $\text{Im } A(u) \cong u(G_1)/u(G_1) \cap L_2$ and $\text{Ker } A(v) \cong v^{-1}(L_3)/L_2$. Since $u^{-1}(L_2)$ is linear, $\text{Ker } A(u)$ is linear as well; it is also proper, and hence finite. Also, $\text{Ker } A(v)/\text{Im } A(u) \cong v^{-1}(L_3)/L_2 + u(G_1)$ is a quotient of $v^{-1}(L_3)/u(G_1) \cong L_3$. It follows similarly that $\text{Ker } A(v)/\text{Im } A(u)$ is finite. \square

Next, we obtain a dual version of Proposition 4.4:

PROPOSITION 4.8. (i) *The Albanese functor yields an exact functor*

$$\underline{A} : \underline{\mathcal{C}} \longrightarrow \underline{\mathcal{A}},$$

which is left adjoint to the inclusion $\underline{\mathcal{A}} \rightarrow \underline{\mathcal{C}}$. Moreover, \underline{A} commutes with base change under field extensions.

(ii) *Every abelian variety is an injective object in $\underline{\mathcal{C}}$.*

Proof. (i) Consider a $\underline{\mathcal{C}}$ -morphism $\underline{f} : G \rightarrow H$ and choose a representative by a \mathcal{C} -morphism $f : G \rightarrow H/H'$, where H' is a finite subgroup of H . This yields an \mathcal{A} -morphism

$$A(f) : A(G) \longrightarrow A(H/H') = A(H)/\alpha_H(H'),$$

and hence an $\underline{\mathcal{A}}$ -morphism $\underline{A}(f) : A(G) \rightarrow A(H)$. One may readily check that $\underline{A}(f)$ depends only on \underline{f} , in a covariant way. By Proposition 3.5 and Lemma 4.5, the resulting functor \underline{A} is exact and commutes with base change under field extensions.

To show the adjointness assertion, consider a smooth connected algebraic group G and an abelian variety A . Then the map

$$\text{Hom}_{\mathcal{A}}(A(G), A) \longrightarrow \text{Hom}_{\mathcal{C}}(G, A), \quad f \longmapsto f \circ \alpha_G$$

is an isomorphism by Lemma 4.5 again. In view of Proposition 3.6, it follows that the analogous map

$$\mathrm{Hom}_{\underline{\mathcal{A}}}(A(G), A) \longrightarrow \mathrm{Hom}_{\underline{\mathcal{C}}}(G, A)$$

is an isomorphism as well.

(ii) This follows formally from (i). \square

REMARKS 4.9. (i) Denote by \widehat{A} the dual of an abelian variety A . Then the assignment $A \mapsto \widehat{A}$ yields a contravariant endofunctor of \mathcal{A} , which is involutive and preserves isogenies and finite products. As an easy consequence, we obtain a contravariant endofunctor of $\underline{\mathcal{A}}$, which is involutive and exact. Note that each abelian variety is (non-canonically) $\underline{\mathcal{A}}$ -isomorphic to its dual, via the choice of a polarization.

(ii) Let k' be a field extension of k . Then the assignment

$$A \mapsto \mathbb{Q} \otimes_{\mathbb{Z}} A(k') =: A(k')_{\mathbb{Q}}$$

yields a functor from \mathcal{A} to the category of \mathbb{Q} -vector spaces (possibly of infinite dimension), which preserves finite products. Moreover, each isogeny

$$f : A \longrightarrow B$$

yields an isomorphism

$$A(k')_{\mathbb{Q}} \xrightarrow{\cong} B(k')_{\mathbb{Q}},$$

since this holds for the multiplication maps n_A . Thus, the above assignment yields an exact functor from $\underline{\mathcal{A}}$ to the category of \mathbb{Q} -vector spaces.

4.3 VANISHING OF EXTENSION GROUPS

In this subsection, we prove the assertion (v) of Theorem 1: $\mathrm{hd}(\underline{\mathcal{C}}) = 1$. We first collect general vanishing results for extension groups in $\underline{\mathcal{C}}$:

LEMMA 4.10. *Let G be a smooth connected algebraic group, U a smooth connected unipotent group, A an abelian variety, and T a torus.*

- (i) $\mathrm{Ext}_{\underline{\mathcal{C}}}^n(T, G) = 0 = \mathrm{Ext}_{\underline{\mathcal{C}}}^n(G, A) = 0$ for all $n \geq 1$.
- (ii) If $\mathrm{char}(k) = 0$, then $\mathrm{Ext}_{\underline{\mathcal{C}}}^n(U, G) = 0$ for all $n \geq 1$.
- (iii) If $\mathrm{char}(k) = p > 0$ and G is divisible, then $\mathrm{Ext}_{\underline{\mathcal{C}}}^n(U, G) = 0 = \mathrm{Ext}_{\underline{\mathcal{C}}}^n(G, U)$ for all $n \geq 0$.
- (iv) If G is linear, then $\mathrm{Ext}_{\underline{\mathcal{C}}}^n(G, T) = 0$ for all $n \geq 1$. If moreover $\mathrm{char}(k) = 0$, then $\mathrm{Ext}_{\underline{\mathcal{C}}}^n(G, U) = 0$ for all $n \geq 1$ as well.

Proof. (i) Just recall that T is projective in $\underline{\mathcal{C}}$ (Proposition 4.4), and A is injective in $\underline{\mathcal{C}}$ (Proposition 4.8).

(ii) Likewise, U is projective in $\underline{\mathcal{C}}$ by Proposition 3.8.

(iii) Since U is unipotent, there exists a positive integer m such that $p_G^m = 0$. It follows that both groups $\text{Ext}_{\underline{\mathcal{C}}}^m(U, G)$ and $\text{Ext}_{\underline{\mathcal{C}}}^m(G, U)$ are p^m -torsion (see e.g. [Oo66, Lem. I.3.1]). But these groups are also modules over $\text{End}_{\underline{\mathcal{C}}}(G)$, and hence \mathbb{Q} -vector spaces by Proposition 3.6. This yields the assertion.

(iv) By Proposition 2.8 and the long exact sequence for Ext groups, we may assume that G is unipotent or a torus. In the latter case, both assertions follows from (i); in the former case, the first assertion follows from (ii) and (iii), and the second assertion, from the fact that unipotent groups are just vector spaces. \square

Next, recall that $\text{hd}(\underline{\mathcal{C}}) \geq 1$ in view of Examples 3.4. So, to conclude that $\text{hd}(\underline{\mathcal{C}}) = 1$, it suffices to check the following:

LEMMA 4.11. *For any smooth connected algebraic groups G, H and any integer $n \geq 2$, we have $\text{Ext}_{\underline{\mathcal{C}}}^n(G, H) = 0$.*

Proof. Let $\eta \in \text{Ext}_{\underline{\mathcal{C}}}^n(G, H)$, where $n \geq 3$. Then η is represented by an exact sequence in $\underline{\mathcal{C}}$

$$0 \longrightarrow H \longrightarrow G_1 \longrightarrow \cdots \longrightarrow G_n \longrightarrow G \longrightarrow 0,$$

which we may cut into two exact sequences in $\underline{\mathcal{C}}$

$$0 \longrightarrow H \longrightarrow G_1 \longrightarrow G_2 \longrightarrow K \longrightarrow 0,$$

$$0 \longrightarrow K \longrightarrow G_3 \longrightarrow \cdots \longrightarrow G_n \longrightarrow G \longrightarrow 0.$$

Thus, η can be written as a Yoneda product $\eta_1 \cup \eta_2$, where $\eta_1 \in \text{Ext}_{\underline{\mathcal{C}}}^2(G, K)$ and $\eta_2 \in \text{Ext}_{\underline{\mathcal{C}}}^{n-2}(K, H)$. So it suffices to show the assertion when $n = 2$.

Using the long exact sequences for Ext groups, we may further reduce to the case where G, H are simple objects in $\underline{\mathcal{C}}$, i.e., \mathbb{G}_a , simple tori T , or simple abelian varieties A (Proposition 3.2). In view of Lemma 4.10, it suffices in turn to check that

$$(i) \quad \text{Ext}_{\underline{\mathcal{C}}}^2(A, T) = 0,$$

$$(ii) \quad \text{Ext}_{\underline{\mathcal{C}}}^2(A, \mathbb{G}_a) = 0 \text{ when } \text{char}(k) = 0,$$

$$(iii) \quad \text{Ext}_{\underline{\mathcal{C}}}^2(\mathbb{G}_a, \mathbb{G}_a) = 0 \text{ when } \text{char}(k) > 0.$$

For (i), we adapt the argument of [Oo66, Prop. II.12.3]. Let $\eta \in \text{Ext}_{\underline{\mathcal{C}}}^2(A, T)$ be represented by an exact sequence in $\underline{\mathcal{C}}$

$$0 \longrightarrow T \xrightarrow{u} G_1 \xrightarrow{v} G_2 \xrightarrow{w} A \longrightarrow 0.$$

As above, $\eta = \eta_1 \cup \eta_2$, where η_1 denotes the class of the extension

$$0 \longrightarrow T \xrightarrow{u} G_1 \longrightarrow K \longrightarrow 0,$$

and η_2 that of the extension

$$0 \longrightarrow K \longrightarrow G_2 \xrightarrow{w} A \longrightarrow 0.$$

Also, note that $\underline{A}(w) : \underline{A}(G_2) \rightarrow A$ is an epimorphism in $\underline{\mathcal{A}}$, and hence has a section, say s . Denoting by H_2 the pull-back of $s(A)$ under the Albanese morphism $G_2 \rightarrow \underline{A}(G_2)$, we have a monomorphism $\iota : H_2 \rightarrow G_2$ in $\underline{\mathcal{C}}$, such that $\underline{A}(w \circ \iota) : \underline{A}(H_2) \rightarrow A$ is an isomorphism. This yields a commutative diagram of exact sequences in $\underline{\mathcal{C}}$

$$\begin{array}{ccccccccc} 0 & \longrightarrow & T & \longrightarrow & H_1 & \longrightarrow & H_2 & \longrightarrow & A & \longrightarrow & 0 \\ & & \text{id} \downarrow & & \downarrow & & \iota \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & T & \xrightarrow{u} & G_1 & \xrightarrow{v} & G_2 & \xrightarrow{w} & A & \longrightarrow & 0. \end{array}$$

Thus, η is represented by the top exact sequence, and hence we may assume that $\underline{A}(w)$ is an isomorphism. Then K is linear in view of Theorem 2.3. Thus, $\kappa_1 = 0$ by Lemma 4.10 (iv). So $\eta = 0$; this completes the proof of (i).

For (ii), we replace T with \mathbb{G}_a in the above argument, and use the vanishing of $\text{Ext}_{\underline{\mathcal{C}}}^1(L, \mathbb{G}_a)$ for L linear (Lemma 4.10 (iv) again).

Finally, for (iii), it suffices to show that $\text{Ext}_{\underline{\mathcal{U}}}^2(\mathbb{G}_a, \mathbb{G}_a) = 0$. Also, we may assume that k is perfect, in view of Theorem 3.11. Let $\eta \in \text{Ext}_{\underline{\mathcal{U}}}^2(\mathbb{G}_a, \mathbb{G}_a)$ be represented by an exact sequence in $\underline{\mathcal{U}}$,

$$0 \longrightarrow \mathbb{G}_a \longrightarrow G_1 \longrightarrow G_2 \longrightarrow \mathbb{G}_a \longrightarrow 0.$$

Then Proposition 3.5 yields an exact sequence in $\underline{\mathcal{U}}$,

$$0 \longrightarrow \mathbb{G}_a \longrightarrow H_1 \longrightarrow H_2 \longrightarrow \mathbb{G}_a \longrightarrow 0,$$

and a commutative diagram in $\underline{\mathcal{U}}$,

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathbb{G}_a & \longrightarrow & G_1 & \longrightarrow & G_2 & \longrightarrow & \mathbb{G}_a & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & \mathbb{G}_a & \longrightarrow & H_1 & \longrightarrow & H_2 & \longrightarrow & \mathbb{G}_a & \longrightarrow & 0. \end{array}$$

where the vertical arrows are isomorphisms in $\underline{\mathcal{U}}$ (here we use the fact that the quotient of \mathbb{G}_a by a finite subgroup is isomorphic to \mathbb{G}_a). As $\text{Ext}_{\underline{\mathcal{U}}}^2(\mathbb{G}_a, \mathbb{G}_a) = 0$ (see [DG70, V.1.5.1, V.1.5.2]), the bottom exact sequence is equivalent to the trivial exact sequence in $\underline{\mathcal{U}}$, and hence in $\underline{\mathcal{U}}$. Thus $\eta = 0$ as desired. \square

REMARK 4.12. When k is perfect, the groups $\text{Ext}_{\underline{\mathcal{C}}}^n(G, H)$ are torsion for all $n \geq 2$ and all algebraic groups G, H , in view of [Mi70, Cor., p. 439]. In fact,

this assertion extends to an arbitrary field k : indeed, it clearly holds when G or H is finite, or more generally m -torsion for some positive integer m . Using Proposition 2.8, one may thus reduce to the case when G, H are simple objects of $\underline{\mathcal{C}}$. Then the assertion is obtained by combining Proposition 3.6, Lemma 4.10, and the proof of Lemma 4.11.

5 STRUCTURE OF ISOGENY CATEGORIES

5.1 VECTOR EXTENSIONS OF ABELIAN VARIETIES

In this subsection, we assume that $\text{char}(k) = 0$. Recall that a *vector extension* of an abelian variety A is an algebraic group G that lies in an extension

$$\xi : 0 \longrightarrow U \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where U is unipotent. Then $U = R_u(G)$ and $A = A(G)$ are uniquely determined by G ; also, the extension ξ has no non-trivial automorphisms, since $\text{Hom}_{\mathcal{C}}(A, U) = 0$. Thus, the data of the algebraic group G and the extension ξ are equivalent.

We denote by \mathcal{V} the full subcategory of \mathcal{C} with objects the vector extensions (of all abelian varieties). By Theorems 2.3 and 2.4, the objects of \mathcal{V} are those smooth connected algebraic groups that admit no non-zero subtorus. In view of Lemmas 4.2 and 4.3, this readily implies:

- LEMMA 5.1. (i) *Let $0 \rightarrow G_1 \rightarrow G_2 \rightarrow G_3 \rightarrow 0$ be an exact sequence in \mathcal{C} , where G_2 is connected. Then G_2 is an object of \mathcal{V} if and only if so are G_3 and G_1^0 .*
- (ii) *Let $f : G \rightarrow H$ be an isogeny of connected algebraic groups. Then G is an object of \mathcal{V} if and only if so is H .*
- (iii) *Let k' be a field extension of k , and G an algebraic k -group. Then G is an object of \mathcal{V}_k if and only if $G_{k'}$ is an object of $\mathcal{V}_{k'}$.*

In particular, \mathcal{V} is stable under taking quotients and extensions, but not subobjects; like \mathcal{T} and \mathcal{A} , it is an additive subcategory of \mathcal{C} , but not an abelian subcategory.

Next, recall from [Ro58] or [MM74, §1.9] that every abelian variety A has a universal vector extension,

$$\xi(A) : 0 \longrightarrow U(A) \longrightarrow E(A) \longrightarrow A \longrightarrow 0,$$

where $U(A)$ is the additive group of the vector space $H^1(A, \mathcal{O}_A)^*$; moreover, $\dim U(A) = \dim A$. Also, $E(A)$ is anti-affine, i.e., every morphism from $E(A)$ to a linear algebraic group is zero (see e.g. [Br17, Prop. 5.5.8]).

PROPOSITION 5.2. (i) *The assignments $A \mapsto E(A)$, $A \mapsto U(A)$ yield additive functors*

$$E : \mathcal{A} \longrightarrow \mathcal{V}, \quad U : \mathcal{A} \longrightarrow \mathcal{U},$$

which commute with base change under field extensions.

(ii) For any morphism $f : A \rightarrow B$ of abelian varieties, the induced morphism $U(f) : U(A) \rightarrow U(B)$ is the dual of the pull-back morphism $f^* : H^1(B, \mathcal{O}_B) \rightarrow H^1(A, \mathcal{O}_A)$. Moreover, $U(f)$ is zero (resp. an isomorphism) if and only if f is zero (resp. an isogeny).

(iii) E is left adjoint to the Albanese functor $A : \mathcal{V} \rightarrow \mathcal{A}$.

Proof. We prove (i) and (ii) simultaneously. Let $f : A \rightarrow B$ be a morphism of abelian varieties. Consider the pull-back diagram of exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U(B) & \longrightarrow & F & \longrightarrow & A & \longrightarrow & 0 \\ & & \text{id} \downarrow & & \downarrow & & f \downarrow & & \\ 0 & \longrightarrow & U(B) & \longrightarrow & E(B) & \longrightarrow & B & \longrightarrow & 0. \end{array}$$

The universal property of $\xi(A)$ yields a commutative diagram of exact sequences

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U(A) & \longrightarrow & E(A) & \longrightarrow & A & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & U(B) & \longrightarrow & F & \longrightarrow & A & \longrightarrow & 0, \end{array}$$

and hence another such diagram,

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U(A) & \longrightarrow & E(A) & \longrightarrow & A & \longrightarrow & 0 \\ & & U(f) \downarrow & & E(f) \downarrow & & f \downarrow & & \\ 0 & \longrightarrow & U(B) & \longrightarrow & E(B) & \longrightarrow & B & \longrightarrow & 0, \end{array}$$

which defines morphisms $E(f)$ and $U(f)$.

Next, let $\eta \in H^1(B, \mathcal{O}_B)$, so that we have a push-out diagram of extensions

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U(B) & \longrightarrow & E(B) & \longrightarrow & B & \longrightarrow & 0 \\ & & \eta \downarrow & & \downarrow & & \text{id} \downarrow & & \\ 0 & \longrightarrow & \mathbb{G}_a & \longrightarrow & E_\eta & \longrightarrow & B & \longrightarrow & 0. \end{array}$$

By construction, the pull-back of E_η by f is the push-out of $\xi(A)$ by the map $\eta \circ U(f) : U(A) \rightarrow \mathbb{G}_a$. Hence $U(f)$ is the dual of $f^* : H^1(B, \mathcal{O}_B) \rightarrow H^1(A, \mathcal{O}_A)$. As a consequence, U is a covariant functor, and hence so is E .

Since the formation of the universal vector extension commutes with base change under field extensions, the functors E and U commute with such base change as well. Clearly, they are additive; this completes the proof of (i).

To complete the proof of (ii), recall the canonical isomorphism

$$H^1(A, \mathcal{O}_A) \cong \text{Lie}(\widehat{A}),$$

where the right-hand side denotes the Lie algebra of the dual abelian variety (see [Mi86, Rem. 9.4]). This isomorphism identifies f^* with $\text{Lie}(\hat{f})$, where $\hat{f} : \hat{B} \rightarrow \hat{A}$ denotes the dual morphism of f . As a consequence,

$$U(f) = 0 \Leftrightarrow f^* = 0 \Leftrightarrow \hat{f} = 0 \Leftrightarrow f = 0,$$

where the second equivalence holds since $\text{char}(k) = 0$, and the third one follows from biduality of abelian varieties. Likewise, $U(f)$ is an isomorphism if and only if f^* is an isomorphism; equivalently, \hat{f} is an isogeny, i.e., f is an isogeny. (iii) Given a vector extension $0 \rightarrow U \rightarrow G \rightarrow A(G) \rightarrow 0$, we check that the map

$$\alpha : \text{Hom}_{\mathcal{V}}(E(A), G) \rightarrow \text{Hom}_{\mathcal{A}}(A, A(G)), \quad u \mapsto A(u)$$

is an isomorphism.

Consider a morphism $u : E(A) \rightarrow G$ such that $A(u) = 0$. Then u factors through a morphism $E(A) \rightarrow R_u(G)$, and hence $u = 0$ as $E(A)$ is anti-affine. Next, consider a morphism $v : A \rightarrow A(G)$. By (i), we have a commutative square

$$\begin{array}{ccc} E(A) & \xrightarrow{E(v)} & E(A(G)) \\ \downarrow & & \downarrow \\ A & \xrightarrow{v} & A(G). \end{array}$$

Also, the universal property of $\xi(A)$ yields a commutative square

$$\begin{array}{ccc} E(A(G)) & \xrightarrow{\delta} & G \\ \downarrow & & \alpha_G \downarrow \\ A(G) & \xrightarrow{\text{id}} & A(G). \end{array}$$

Thus, $w := E(v) \circ \delta \in \text{Hom}_{\mathcal{V}}(E(A), G)$ satisfies $\alpha(w) = u$. □

Denote by $\underline{\mathcal{V}}$ the isogeny category of vector extensions, that is, the full subcategory of $\underline{\mathcal{C}}$ with the same objects as \mathcal{V} . Then $\underline{\mathcal{V}}$ is an abelian category in view of Lemma 5.1. Also, Proposition 3.6 yields natural isomorphisms

$$\mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{C}}(G_1, G_2) \xrightarrow{\cong} \text{Hom}_{\underline{\mathcal{V}}}(G_1, G_2), \quad \mathbb{Q} \otimes_{\mathbb{Z}} \text{Ext}_{\mathcal{C}}^1(G_1, G_2) \xrightarrow{\cong} \text{Ext}_{\underline{\mathcal{V}}}^1(G_1, G_2)$$

for any objects G_1, G_2 of $\underline{\mathcal{V}}$.

COROLLARY 5.3. (i) *The functors $E : \mathcal{A} \rightarrow \mathcal{V}$, $U : \mathcal{A} \rightarrow \mathcal{U}$ yield exact functors*

$$\underline{E} : \underline{\mathcal{A}} \rightarrow \underline{\mathcal{V}}, \quad \underline{U} : \underline{\mathcal{A}} \rightarrow \underline{\mathcal{U}},$$

which commute with base change under field extensions. Moreover, \underline{E} is left adjoint to the Albanese functor $\underline{A} : \underline{\mathcal{V}} \rightarrow \underline{\mathcal{A}}$.

- (ii) *The universal vector extension of any abelian variety is a projective object of $\underline{\mathcal{V}}$.*

Proof. (i) This follows from Propositions 3.5 and 5.2.

(ii) We have canonical isomorphisms for any object G of $\underline{\mathcal{V}}$:

$$\mathrm{Hom}_{\underline{\mathcal{V}}}(E(A), G) \cong \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{V}}(E(A), G) \cong \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{A}}(A, A(G)),$$

where the first isomorphism follows from Proposition 3.6, and the second one from Proposition 5.2. Using Proposition 3.6 again, this yields a canonical isomorphism

$$\mathrm{Hom}_{\underline{\mathcal{V}}}(E(A), G) \cong \mathrm{Hom}_{\underline{\mathcal{A}}}(A, \underline{A}(G)).$$

Since the Albanese functor \underline{A} is exact (Proposition 4.8), it follows that the functor $G \mapsto \mathrm{Hom}_{\underline{\mathcal{V}}}(E(A), G)$ is exact as well. \square

Next, let G be an object of \mathcal{V} . Form and label the commutative diagram of exact sequences in \mathcal{C}

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U(A) & \xrightarrow{\iota} & E(A) & \longrightarrow & A & \longrightarrow & 0 \\ & & \gamma \downarrow & & \delta \downarrow & & \mathrm{id} \downarrow & & \\ 0 & \longrightarrow & U & \longrightarrow & G & \longrightarrow & A & \longrightarrow & 0, \end{array}$$

where $U = U(G)$, $A = A(G)$, and $\gamma = \gamma_G$ classifies the bottom extension. This yields an exact sequence in \mathcal{C}

$$\xi : 0 \longrightarrow U(A) \xrightarrow{\gamma - \iota} U \times E(A) \longrightarrow G \longrightarrow 0.$$

PROPOSITION 5.4. *Keep the above notation.*

- (i) ξ yields a projective resolution of G in $\underline{\mathcal{V}}$.
- (ii) For any object H of \mathcal{V} , we have an exact sequence

$$0 \longrightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(G, H) \xrightarrow{\varphi} \mathrm{Hom}_{\mathcal{U}}(U(G), U(H)) \times \mathrm{Hom}_{\underline{\mathcal{A}}}(A(G), A(H)) \\ \xrightarrow{\psi} \mathrm{Hom}_{\mathcal{U}}(U(A(G)), U(H)) \longrightarrow \mathrm{Ext}_{\underline{\mathcal{V}}}^1(G, H) \longrightarrow 0,$$

where $\varphi(\underline{f}) := (U(\underline{f}), \underline{A}(\underline{f}))$, and $\psi(u, v) := u \circ \gamma_G - \gamma_H \circ U(v)$.

Proof. (i) This holds as $U, U(A)$ are projective in \mathcal{C} (Theorem 2.9), and $E(A)$ is projective in $\underline{\mathcal{C}}$ (Proposition 5.2).

(ii) In view of (i), this follows readily from the long exact sequence of extension groups

$$0 \rightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(G, H) \rightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(U \times E(A), H) \rightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(U, H) \rightarrow \mathrm{Ext}_{\underline{\mathcal{V}}}^1(G, H) \rightarrow 0$$

associated with the short exact sequence ξ . \square

As a direct consequence of Proposition 5.4, we obtain:

COROLLARY 5.5. *The following conditions are equivalent for an object G of \mathcal{V} :*

- (i) $G \cong E(A)$ in $\underline{\mathcal{V}}$ for some abelian variety A .
- (ii) $\mathrm{Hom}_{\underline{\mathcal{V}}}(G, \mathbb{G}_a) = \mathrm{Ext}_{\underline{\mathcal{V}}}^1(G, \mathbb{G}_a) = 0$.

As a further consequence, we describe the projective or injective objects of $\underline{\mathcal{V}}$:

COROLLARY 5.6. (i) *The projective objects of $\underline{\mathcal{V}}$ are the products $V \times E(A)$, where V is unipotent, and A is an abelian variety.*

(ii) *The injective objects of $\underline{\mathcal{V}}$ are the abelian varieties.*

Proof. Let G be an extension of an abelian variety A by a unipotent group U .

(i) If G is projective in $\underline{\mathcal{V}}$, then $\mathrm{Ext}_{\underline{\mathcal{V}}}^1(G, \mathbb{G}_a) = 0$. In view of Proposition 5.4, it follows that the map $\mathrm{Hom}_{\mathcal{U}}(U, \mathbb{G}_a) \rightarrow \mathrm{Hom}_{\mathcal{U}}(U(A), \mathbb{G}_a)$, $u \mapsto u \circ \gamma$ is surjective. Equivalently, γ is injective; hence so is $\delta : E(A) \rightarrow G$. Identifying $E(A)$ with a subgroup of G , it follows that $G = U + E(A)$, and $U(A) \subset U$. We may choose a complement $V \subset U$ to the subspace $U(A) \subset U$; then $G \cong V \times E(A)$. Conversely, every such product is projective by Proposition 5.2. This yields the assertion.

(ii) If G is injective in $\underline{\mathcal{V}}$, then $\mathrm{Ext}_{\underline{\mathcal{V}}}^1(B, G) = 0$ for any abelian variety B . Thus, we have an exact sequence

$$0 \longrightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(B, U) \longrightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(B, G) \longrightarrow \mathrm{Hom}_{\underline{\mathcal{V}}}(B, A) \xrightarrow{\partial} \mathrm{Ext}_{\underline{\mathcal{V}}}^1(B, U) \longrightarrow 0.$$

Moreover, $\mathrm{Ext}_{\underline{\mathcal{V}}}^1(B, U) \cong \mathrm{Hom}_{\mathcal{U}}(U(B), U)$, as follows e.g. from Proposition 5.4. Since $\mathrm{Hom}_{\underline{\mathcal{V}}}(B, A) = \mathrm{Hom}_{\underline{\mathcal{A}}}(B, A)$ is a finite-dimensional \mathbb{Q} -vector space, so is $\mathrm{Hom}_{\mathcal{U}}(U(B), U)$.

When k is not a number field, i.e., k is an infinite-dimensional \mathbb{Q} -vector space, this forces $U = 0$, since $U(B) \neq 0$ for any non-zero abelian variety B . Thus, G is an abelian variety.

On the other hand, when k is a number field, there are only finitely many isomorphism classes of abelian varieties that are isogenous to any prescribed abelian variety (see [MW93] for a quantitative version of this finiteness result). As a consequence, we may choose a simple abelian variety B , not isogenous to any simple factor of A . Then $\mathrm{Hom}_{\underline{\mathcal{A}}}(B, A) = 0$; as above, this yields $U = 0$, i.e., G is an abelian variety.

Conversely, every abelian variety is injective in $\underline{\mathcal{V}}$ by Proposition 4.8. □

We now describe the structure of \mathcal{V} and $\underline{\mathcal{V}}$ in terms of linear algebra. Let \mathcal{D} be the category with objects the triples (A, U, γ) , where A is an abelian variety, U a unipotent group, and $\gamma : U(A) \rightarrow U$ a morphism; the \mathcal{D} -morphisms

from (A_1, U_1, γ_1) to (A_2, U_2, γ_2) are those pairs of \mathcal{C} -morphisms $u : U_1 \rightarrow U_2$, $v : A_1 \rightarrow A_2$ such that the square

$$\begin{array}{ccc} U(A_1) & \xrightarrow{U(v)} & U(A_2) \\ \gamma_1 \downarrow & & \gamma_2 \downarrow \\ U_1 & \xrightarrow{u} & U_2 \end{array}$$

commutes. We also introduce the ‘isogeny category’ $\underline{\mathcal{D}}$, by allowing v to be a $\underline{\mathcal{C}}$ -morphism in the above definition (this makes sense in view of Corollary 5.3). Next, define a functor

$$D : \mathcal{V} \longrightarrow \mathcal{D}$$

by assigning to each object G the triple $(A(G), R_u(G), \gamma)$, where $\gamma : U(A(G)) \rightarrow R_u(G)$ denotes the classifying map, and to each morphism $f : G_1 \rightarrow G_2$, the pair $(A(f), U(f))$. By Corollary 5.3 again, we may define similarly a functor

$$\underline{D} : \underline{\mathcal{V}} \longrightarrow \underline{\mathcal{D}}.$$

PROPOSITION 5.7. *With the above notation, the functors D and \underline{D} yield equivalences of categories.*

We omit the easy proof.

5.2 SEMI-ABELIAN VARIETIES

Recall that a semi-abelian variety is an algebraic group G that lies in an extension

$$\xi : 0 \longrightarrow T \longrightarrow G \longrightarrow A \longrightarrow 0,$$

where T is a torus, and A an abelian variety. We now adapt part of the results of §5.1 to this setting, leaving the (easy) verifications to the motivated reader. The algebraic groups $T = T(G)$ and $A = A(G)$ are uniquely determined by G , and the extension ξ has no non-trivial automorphisms. Thus, the data of G and of the extension ξ are equivalent. Moreover, recall the natural isomorphism

$$c : \mathrm{Ext}_{\mathcal{C}}^1(A, T) \xrightarrow{\cong} \mathrm{Hom}^\Gamma(X(T), \widehat{A}(k_s)),$$

which arises from the Weil-Barsotti isomorphism

$$\mathrm{Ext}_{\mathcal{C}_{k_s}}^1(A_{k_s}, \mathbb{G}_{m, k_s}) \xrightarrow{\cong} \widehat{A}(k_s)$$

together with the pairing

$$\mathrm{Ext}_{\mathcal{C}}^1(A, T) \times X(T) \longrightarrow \mathrm{Ext}_{\mathcal{C}_{k_s}}^1(A_{k_s}, \mathbb{G}_{m, k_s})$$

given by push-out of extensions via characters of T .

Denote by \mathcal{S} the full subcategory of \mathcal{C} with objects the semi-abelian varieties. Then the analogue of Lemma 5.1 holds in view e.g. of [Br17, §5.4] (but there

is no analogue of the universal vector extension in this setting). Thus, the isogeny category of semi-abelian varieties, $\underline{\mathcal{S}}$, is an abelian category. As for vector extensions of abelian varieties, we have natural isomorphisms

$$\mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{C}}(G_1, G_2) \xrightarrow{\cong} \text{Hom}_{\underline{\mathcal{S}}}(G_1, G_2), \quad \mathbb{Q} \otimes_{\mathbb{Z}} \text{Ext}_{\mathcal{C}}^1(G_1, G_2) \xrightarrow{\cong} \text{Ext}_{\underline{\mathcal{S}}}^1(G_1, G_2)$$

for any objects G_1, G_2 of $\underline{\mathcal{S}}$. This yields a natural isomorphism

$$\text{Ext}_{\underline{\mathcal{C}}}^1(A, T) \xrightarrow{\cong} \text{Hom}^{\Gamma}(X(T)_{\mathbb{Q}}, \widehat{A}(k_s)_{\mathbb{Q}}).$$

Note that the assignment

$$A \longmapsto \widehat{A}(k_s)_{\mathbb{Q}}$$

yields an exact functor from $\underline{\mathcal{A}}$ to the category of \mathbb{Q} -vector spaces equipped with the discrete topology and a continuous representation of Γ , as follows e.g. from Remarks 4.9.

Next, we obtain a description of $\underline{\mathcal{S}}$ in terms of linear algebra. Let $\underline{\mathcal{E}}$ be the category with objects the triples (A, M, c) , where A is an abelian variety, M a finite-dimensional \mathbb{Q} -vector space equipped with a continuous action of Γ , and $c : M \rightarrow \widehat{A}(k_s)_{\mathbb{Q}}$ a Γ -equivariant linear map; the $\underline{\mathcal{E}}$ -morphisms from (A_1, M_1, c_1) to (A_2, M_2, c_2) are those pairs (\underline{u}, v) , where $\underline{u} : A_1 \rightarrow A_2$ is a $\underline{\mathcal{A}}$ -morphism and $v : M_2 \rightarrow M_1$ a Γ -equivariant linear map, such that the square

$$\begin{array}{ccc} M_2 & \xrightarrow{c_2} & \widehat{A}_2(k_s)_{\mathbb{Q}} \\ v \downarrow & & \underline{u} \downarrow \\ M_1 & \xrightarrow{c_1} & \widehat{A}_1(k_s)_{\mathbb{Q}} \end{array}$$

commutes. Then one may check that the assignment

$$G \longmapsto (A(G), X(T(G))_{\mathbb{Q}}, c(G)_{\mathbb{Q}})$$

yields an equivalence of categories $\underline{\mathcal{S}} \rightarrow \underline{\mathcal{E}}$. Moreover, the sequence

$$\begin{aligned} 0 &\longrightarrow \text{Hom}_{\underline{\mathcal{S}}}(G_1, G_2) \xrightarrow{\varphi} \text{Hom}^{\Gamma}(M_2, M_1) \times \text{Hom}_{\underline{\mathcal{A}}}(A_1, A_2) \\ &\xrightarrow{\psi} \text{Hom}^{\Gamma}(M_2, \widehat{A}_1(k_s)_{\mathbb{Q}}) \longrightarrow \text{Ext}_{\underline{\mathcal{S}}}^1(G_1, G_2) \longrightarrow 0 \end{aligned}$$

is exact for any semi-abelian varieties G_1, G_2 , where $\underline{\mathcal{E}}(G_i) := (A_i, M_i, c_i)$ for $i = 1, 2$, $\varphi(\underline{f}) := (X_{\mathbb{Q}} \circ T)(\underline{f}), \underline{\mathcal{A}}(\underline{f})$, and $\psi(u, v) := c_1 \circ u - \hat{v} \circ c_2$.

Yet there are important differences between the isogeny categories of vector extensions and semi-abelian varieties. For example, the latter does not have enough projectives in general:

PROPOSITION 5.8. (i) *If k is not locally finite, then the projective objects of $\underline{\mathcal{S}}$ are the tori.*

(ii) *If k is locally finite, then the product functor $\underline{\mathcal{T}} \times \underline{\mathcal{A}} \rightarrow \underline{\mathcal{S}}$ yields an equivalence of categories.*

Proof. (i) Let G be a semi-abelian variety, extension of an abelian variety A by a torus T . Denote by $c : X(T) \rightarrow \widehat{A}(k_s)$ the classifying map, and by

$$c_{\mathbb{Q}} : X(T)_{\mathbb{Q}} \longrightarrow \widehat{A}(k_s)_{\mathbb{Q}}$$

the corresponding \mathbb{Q} -linear map; recall that c and $c_{\mathbb{Q}}$ are Γ -equivariant. If G is projective in $\underline{\mathcal{C}}$, then $\text{Ext}_{\underline{\mathcal{C}}}^1(G, T') = 0$ for any torus T' . Thus, we have an exact sequence

$$0 \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(A, T') \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(G, T') \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(T, T') \xrightarrow{\partial} \text{Ext}_{\underline{\mathcal{C}}}^1(A, T') \longrightarrow 0.$$

Moreover, the boundary map ∂ may be identified with the map

$$\text{Hom}^{\Gamma}(X(T')_{\mathbb{Q}}, X(T)_{\mathbb{Q}}) \longrightarrow \text{Hom}^{\Gamma}(X(T')_{\mathbb{Q}}, \widehat{A}(k_s)_{\mathbb{Q}}). \quad f \longmapsto c_{\mathbb{Q}} \circ f.$$

Since ∂ is surjective, and $X(T')_{\mathbb{Q}}$ may be chosen arbitrarily among finite-dimensional \mathbb{Q} -vector spaces equipped with a continuous representation of Γ , the map $c_{\mathbb{Q}}$ is surjective as well. In particular, the abelian group $\widehat{A}(k_s)$ has finite rank. In view of [FJ74, Thm. 9.1], this forces A to be zero, i.e., G is a torus.

(ii) This follows readily from Proposition 2.5 and Corollary 2.13. □

5.3 PRODUCT DECOMPOSITIONS

In this subsection, we first prove the remaining assertions (ii) and (iii) of Theorem 1. Then we describe the isogeny category $\underline{\mathcal{C}}$ in characteristic 0, and its projective or injective objects in arbitrary characteristics.

PROPOSITION 5.9. (i) *If k is perfect, then the product functor $\mathcal{M} \times \mathcal{U} \rightarrow \mathcal{L}$ yields an equivalence of categories.*

(ii) *For any field k , the product functor $\underline{\mathcal{T}} \times \underline{\mathcal{U}} \rightarrow \underline{\mathcal{L}}$ yields an equivalence of categories.*

Proof. (i) This follows readily from Theorem 2.4 and Proposition 2.5. (ii) This is a consequence of (i) in view of Theorem 3.11. □

PROPOSITION 5.10. *If $\text{char}(k) > 0$, then the product functor $\underline{\mathcal{S}} \times \underline{\mathcal{U}} \rightarrow \underline{\mathcal{C}}$ yields an equivalence of categories.*

Proof. Let G be a smooth connected algebraic group. By Corollary 2.13, there exists a finite subgroup $F \subset G$ such that $G/F \cong S \times U$, where S is a semi-abelian variety, and U is unipotent. Thus, the product functor Π is essentially surjective.

Next, let S_1, S_2 be semi-abelian varieties, and U_1, U_2 smooth connected unipotent groups. We check that Π induces an isomorphism

$$\text{Hom}_{\underline{\mathcal{C}}}(S_1, S_2) \times \text{Hom}_{\underline{\mathcal{C}}}(U_1, U_2) \longrightarrow \text{Hom}_{\underline{\mathcal{C}}}(S_1 \times U_1, S_2 \times U_2), \quad (\underline{\varphi}, \underline{\psi}) \longmapsto \underline{\varphi} \times \underline{\psi}.$$

Assume that $\varphi \times \psi = 0$. Choose representatives $\varphi : S_1 \rightarrow S_2/S'_2$ and $\psi : U_1 \rightarrow U_2/U'_2$, where S'_2, U'_2 are finite. Then $\varphi \times \psi : S_1 \times U_1 \rightarrow (S_2 \times U_2)/(S'_2 \times U'_2)$ has finite image, and hence is zero by Lemma 2.1. So $\varphi = \psi = 0$.

Let $\gamma \in \text{Hom}_{\mathcal{C}}(S_1 \times U_1, S_2 \times U_2)$ be represented by $\gamma : \overline{S}_1 \times \overline{U}_1 \rightarrow (S_2 \times U_2)/F$, where F is finite. Then $F \subset S'_2 \times U'_2$ for some finite subgroups $S'_2 \subset S_2, U'_2 \subset U_2$. Thus, we may assume that $F = S'_2 \times U'_2$. Then the composite morphisms

$$\begin{aligned} S_1 &\longrightarrow S_1 \times U_1 \xrightarrow{\gamma} S_2/S'_2 \times U_2/U'_2 \longrightarrow U_2/U'_2, \\ U_1 &\longrightarrow S_1 \times U_1 \xrightarrow{\gamma} S_2/S'_2 \times U_2/U'_2 \longrightarrow S_2/S'_2 \end{aligned}$$

are zero by Lemma 2.1 and Proposition 2.5. Thus, $\gamma = \varphi \times \psi$ for some morphisms $\varphi : S_1 \rightarrow S_2/S'_2, \psi : U_1 \rightarrow U_2/U'_2$. □

Combining Propositions 5.8 (i) and 5.10, we obtain readily:

COROLLARY 5.11. *If k is locally finite, then the product functor*

$$\mathcal{I} \times \underline{\mathcal{A}} \times \underline{\mathcal{U}} \longrightarrow \underline{\mathcal{C}}$$

yields an equivalence of categories.

REMARKS 5.12. (i) With the notation of the above corollary, each of the categories $\mathcal{I}, \underline{\mathcal{A}}, \underline{\mathcal{U}}$ admits a description of its own. By Proposition 4.1, \mathcal{I} is equivalent to the category of \mathbb{Q} -vector spaces equipped with an automorphism of finite order. Also, the isomorphism classes of abelian varieties over a finite field are classified by the Honda-Tate theorem (see [Ho68, Tat66]); their endomorphism rings are investigated in [Wa69]. Finally, the structure of $\underline{\mathcal{U}}$ (obtained in [DG70, §V.3.6.7]) has been described in Remark 3.12.

(ii) Combining Lemma 2.2, Theorem 2.4 and Lemma 3.10, one may show that the product functor $\mathcal{M}/\mathcal{IM} \times \mathcal{U}/\mathcal{IU} \rightarrow \mathcal{L}/\mathcal{I}$ yields an equivalence of categories. Here \mathcal{I} denotes the category of infinitesimal algebraic groups, and \mathcal{IM} (resp. \mathcal{IU}) the full subcategory of infinitesimal groups of multiplicative type (resp. unipotent).

Next, assume that $\text{char}(k) = 0$. Then every algebraic group is isogenous to a fibered product $E \times_A S$, where E is a vector extension of the abelian variety A , and S is semi-abelian with Albanese variety isomorphic to A (see e.g. Remark 2.10). This motivates the consideration of the fibered product $\mathcal{V} \times_{\underline{\mathcal{A}}} \underline{\mathcal{S}}$: this is the category with objects the triples (E, S, f) , where E is a vector extension of an abelian variety, S a semi-abelian variety, and $f : A(E) \rightarrow A(S)$ an $\underline{\mathcal{A}}$ -isomorphism. The morphisms from (E_1, S_1, f_1) to (E_2, S_2, f_2) are those pairs of $\underline{\mathcal{C}}$ -morphisms $\underline{u} : E_1 \rightarrow E_2, \underline{v} : S_1 \rightarrow S_2$ such that the square

$$\begin{array}{ccc} A(E_1) & \xrightarrow{A(\underline{u})} & A(E_2) \\ \underline{f}_1 \downarrow & & \underline{f}_2 \downarrow \\ A(S_1) & \xrightarrow{A(\underline{v})} & A(S_2) \end{array}$$

commutes in $\underline{\mathcal{A}}$.

PROPOSITION 5.13. *If $\text{char}(k) = 0$, then $\underline{\mathcal{C}}$ is equivalent to $\underline{\mathcal{V}} \times_{\underline{\mathcal{A}}} \underline{\mathcal{S}}$.*

The proof is similar to that of Proposition 5.10, and will be omitted. Note that the descriptions of $\underline{\mathcal{V}}$ and $\underline{\mathcal{S}}$ in terms of linear algebra, obtained in §5.1 and §5.2, can also be reformulated in terms of fibered products of categories. Returning to an arbitrary field k , we obtain:

THEOREM 5.14. *The projective objects of $\underline{\mathcal{C}}$ are:*

- *the linear algebraic groups, if $\text{char}(k) = 0$.*
- *the semi-abelian varieties, if k is locally finite.*
- *the tori, if $\text{char}(k) > 0$ and k is not locally finite.*

Proof. Let G be a smooth connected algebraic group. As a consequence of Theorem 2.9 and Proposition 5.10, we have an exact sequence in $\underline{\mathcal{C}}$

$$0 \longrightarrow U \longrightarrow G \longrightarrow S \longrightarrow 0,$$

where U is smooth, connected, and unipotent, and S is a semi-abelian variety. If G is projective in $\underline{\mathcal{C}}$, then $\text{Ext}_{\underline{\mathcal{C}}}^1(G, T') = 0$ for any torus T' . Since $\text{Hom}_{\underline{\mathcal{C}}}(U, T') = 0$ (as a consequence of Proposition 2.5) and $\text{Ext}_{\underline{\mathcal{C}}}^1(U, T') = 0$ (by Lemma 4.10), the long exact sequence for Ext groups yields $\text{Ext}_{\underline{\mathcal{C}}}^1(S, T') = 0$ as well. By arguing as in the proof of Proposition 5.8, this forces either $A(S)$ to be zero, or k to be locally finite.

If $A(S) = 0$, then G is linear, and hence $G \cong T \times U$ in $\underline{\mathcal{C}}$ by Proposition 5.9. Moreover, tori are projective in $\underline{\mathcal{C}}$ by Proposition 4.4; thus, we may assume that G is unipotent. If $\text{char}(k) = 0$, then every unipotent group is projective, as follows e.g. from Lemma 4.10. If $\text{char}(k) > 0$ and $G \neq 0$, then there exists an exact sequence

$$0 \longrightarrow H \longrightarrow G \longrightarrow \mathbb{G}_a \longrightarrow 0$$

in \mathcal{C} . Since $\text{Ext}_{\underline{\mathcal{C}}}^1(G, \mathbb{G}_a) = 0 = \text{Ext}_{\underline{\mathcal{C}}}^2(H, \mathbb{G}_a)$, it follows that $\text{Ext}_{\underline{\mathcal{C}}}^1(\mathbb{G}_a, \mathbb{G}_a) = 0$. But this contradicts Example 3.4 (ii), hence $G = 0$.

On the other hand, if k is locally finite, then $G \cong T \times A \times U$ in $\underline{\mathcal{C}}$ (by Corollary 5.11) and it follows as above that U is zero. Conversely, every semi-abelian variety is projective in $\underline{\mathcal{C}}$, by Corollary 5.11 again. \square

COROLLARY 5.15. *If $\text{char}(k) > 0$, then \mathcal{C} has no non-zero projective objects.*

Proof. Let G be a projective object of \mathcal{C} . By the claim in the proof of Theorem 2.9 (iii), the (abstract) group $\text{Hom}_{\mathcal{C}}(G, H)$ is divisible for any divisible group H . By arguing as in that proof, it follows that G is linear. Also, for any torus T , the group $\text{Hom}_{\mathcal{C}}(G, T)$ is finitely generated and divisible, hence zero.

Next, we show that G is connected. Indeed, the quotient G/G^0 is finite and étale, hence contained in a torus T . As $\text{Hom}_{\mathcal{C}}(G, T) = 0$, this yields the assertion.

In view of Theorem 2.4, we obtain an exact sequence

$$0 \longrightarrow M \longrightarrow G \longrightarrow U \longrightarrow 0,$$

where M is of multiplicative type, and U is unipotent and connected. We now show that U is projective in the unipotent category \mathcal{U} . Indeed, given an exact sequence $U_1 \rightarrow U_2 \rightarrow 0$ in \mathcal{U} and a morphism $\varphi : U \rightarrow U_2$, we may lift the composition $\psi : G \rightarrow U \rightarrow U_2$ to a morphism $\gamma : G \rightarrow U_1$. Then $\gamma(M) = 0$ (Proposition 2.5) and hence γ factors through a morphism $\delta : U \rightarrow U_1$, which lifts φ .

Let F be an infinitesimal subgroup of U such that U/F is smooth; then U/F is an object of \mathcal{U} , and one easily checks by using Proposition 3.5 that U/F is projective in $\underline{\mathcal{U}}$. In view of Proposition 5.10 and Theorem 5.14, it follows that U/F is zero, hence U is infinitesimal. If $U \neq 0$ then there exists an epimorphism $U \rightarrow \alpha_p$ and hence a non-zero morphism $U \rightarrow A$ for some (supersingular) abelian variety A . Since $\text{Hom}_{\mathcal{C}}(G, A) = 0$, this yields a contradiction. Thus, $U = 0$, i.e., G is of multiplicative type, hence contained in a torus, hence zero. \square

THEOREM 5.16. *The injective objects of $\underline{\mathcal{C}}$ are the semi-abelian varieties if k is locally finite, and the abelian varieties otherwise.*

Proof. Let G be an injective object of $\underline{\mathcal{C}}$, and

$$0 \longrightarrow G_1 \longrightarrow G \longrightarrow G_2 \longrightarrow 0$$

an exact sequence in $\underline{\mathcal{C}}$. Since $\text{hd}(\underline{\mathcal{C}}) = 1$, the map $\text{Ext}_{\underline{\mathcal{C}}}^1(H, G) \rightarrow \text{Ext}_{\underline{\mathcal{C}}}^1(H, G_2)$ is an epimorphism for any object H of $\underline{\mathcal{C}}$. Thus, every quotient of G is injective. In particular, so is the largest semi-abelian quotient S (in view of Remark 2.10 and Theorem 2.11).

We now adapt the argument of Corollary 5.6. Recall that S lies in a unique extension

$$0 \longrightarrow T \longrightarrow S \longrightarrow A \longrightarrow 0,$$

classified by a Γ -equivariant morphism $c : X(T) \rightarrow \widehat{A}(k_s)$. Since S is injective in $\underline{\mathcal{C}}$, we have $\text{Ext}_{\underline{\mathcal{C}}}^1(B, S) = 0$ for any abelian variety B , and hence the connecting homomorphism $\partial : \text{Hom}_{\underline{\mathcal{C}}}(B, A) \rightarrow \text{Ext}_{\underline{\mathcal{C}}}^1(B, T)$ is surjective. Under the isomorphisms

$$\text{Hom}_{\underline{\mathcal{C}}}(B, A) \cong \mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{C}}(B, A), \quad \text{Ext}_{\underline{\mathcal{C}}}^1(B, T) \cong \mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}^{\Gamma}(X(T), \widehat{B}(k_s))$$

(Proposition 3.6 and §5.2), the map ∂ is identified with the composition of the natural map

$$\mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}_{\mathcal{C}}(B, A) \longrightarrow \mathbb{Q} \otimes_{\mathbb{Z}} \text{Hom}^{\Gamma}(\widehat{A}(k_s), \widehat{B}(k_s))$$

with the map

$$\gamma : \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}^{\Gamma}(\widehat{A}(k_s), \widehat{B}(k_s)) \longrightarrow \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}^{\Gamma}(X(T), \widehat{B}(k_s)), \quad u \longmapsto u \circ c.$$

Next, consider a free abelian group M of finite rank, equipped with a continuous action of Γ . Then for any abelian variety B , the tensor product $B_{k_s} \otimes_{\mathbb{Z}} M$ is a k_s -abelian variety equipped with the diagonal Γ -action, and descends to a unique k -abelian variety $B(M)$. Replacing B with $B(M)$, it follows that the corresponding map

$$\partial(M) : \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{C}_{k_s}}^{\Gamma}(B_{k_s} \otimes_{\mathbb{Z}} M, A_{k_s}) \longrightarrow \mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}^{\Gamma}(X(T) \otimes_{\mathbb{Z}} M, \widehat{B}(k_s))$$

is surjective as well. As $\mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}_{\mathcal{C}_{k_s}}(B_{k_s}, A_{k_s})$ and $\mathbb{Q} \otimes_{\mathbb{Z}} \mathrm{Hom}(X(T), \widehat{B}(k_s))$ are direct sums of continuous, finite-dimensional Γ -modules, and M is arbitrary, it follows that the natural map

$$\mathrm{Hom}_{\mathcal{C}_{k_s}}(B_{k_s}, A_{k_s}) \longrightarrow \mathrm{Hom}(X(T), \widehat{B}(k_s))$$

is surjective over the rationals.

In the case where k is not locally finite, recall from [FJ74, Thm. 9.1] that the group $\widehat{B}(k_s)$ has infinite rank for any non-zero abelian variety B . Since $\mathrm{Hom}_{\mathcal{C}_{k_s}}(B_{k_s}, A_{k_s})$ has finite rank, it follows that $X(T) = 0$, i.e., S is an abelian variety. We now distinguish between two subcases.

If $\mathrm{char}(k) = 0$, then G is a vector extension of A , and hence is injective in the category $\underline{\mathcal{Y}}$. Thus, $G = A$ in view of Corollary 5.6.

If $\mathrm{char}(k) > 0$, then $G \cong A \times U$ in $\underline{\mathcal{C}}$ for some split unipotent group U , which must be injective in $\underline{\mathcal{C}}$. If in addition $U \neq 0$, then it follows that \mathbb{G}_a (a quotient of U) is injective in $\underline{\mathcal{C}}$ as well. But $\mathrm{Ext}_{\underline{\mathcal{C}}}^1(\mathbb{G}_a, \mathbb{G}_a) \neq 0$ in view of Example 3.4, a contradiction. Thus, $G = A$ again.

Finally, in the case where k is locally finite, we have $G \cong T \times A \times U$ in $\underline{\mathcal{C}}$ with an obvious notation. As above, we obtain that $U = 0$; on the other hand, T and A are injective in $\underline{\mathcal{C}}$ by Corollary 5.11. This completes the proof. \square

REMARK 5.17. The category \mathcal{C} has no non-zero injective objects. This result should be well-known, but we could not find it in the literature; also, it does not seem to follow from Theorem 5.16, as the relation between injective objects in \mathcal{C} and $\underline{\mathcal{C}}$ is unclear. So we sketch a direct proof: let G be an injective object of \mathcal{C} . For any positive integer n which is prime to $\mathrm{char}(k)$, the n -torsion subgroup $G[n]$ is finite and étale, hence isomorphic to a subgroup of some torus T_n . The inclusion $G[n] \subset G$ extends to a morphism $T_n \rightarrow G$, and hence $G[n]$ is contained in the maximal torus $T(G)$. Likewise, $G[n]$ is isomorphic to a subgroup of an abelian variety A_n , and hence is contained in the largest abelian subvariety of G . Since this abelian variety intersects $T(G)$ along a finite subgroup, it follows that $G[n] = 0$ for $n \gg 0$.

If $\mathrm{char}(k) = 0$, then there is an exact sequence

$$0 \longrightarrow R_u(G) \longrightarrow G \longrightarrow H \longrightarrow 0,$$

where H^0 is a semi-abelian variety. Since $n_{R_u(G)}$ is an isomorphism, we have $G[n] \cong H[n]$ for all n ; it follows that H is finite. Thus, $G = R_u(G) + F$ for some finite group F , which embeds into some torus T . As above, it follows that $F = 0$, i.e., G is unipotent. If $G \neq 0$, then $\text{Ext}_{\mathcal{C}}^1(A, G) \neq 0$ for any abelian variety A , a contradiction.

If $\text{char}(k) = p > 0$, then there is an exact sequence

$$0 \longrightarrow H \longrightarrow G \longrightarrow U \longrightarrow 0,$$

where H^0 is again a semi-abelian variety, H/H^0 is of multiplicative type, and U is unipotent. Thus, $H[n] \cong G[n]$ for all n prime to p , and hence H is finite. It follows as above that $H = 0$, i.e., G is unipotent. If $G \neq 0$, then G contains a copy of α_p , which embeds into some abelian variety; this yields a contradiction.

ACKNOWLEDGEMENTS. I warmly thank Claire Amiot, Brian Conrad, Hélène Esnault, Stéphane Guillermou, Bruno Kahn, Giancarlo Lucchini Arteche, Gaël Rémond, Jean-Pierre Serre, and Catharina Stroppel for very helpful discussions or e-mail exchanges. Special thanks are due to an anonymous referee for a careful reading and valuable comments.

REFERENCES

- [BLR90] S. Bosch, W. Lütkebohmert, M. Raynaud, *Néron models*, Ergebnisse Math. Grenzg. (3) 21, Springer, 1990.
- [Br15] M. Brion, *On extensions of algebraic groups with finite quotient*, Pacific J. Math. 279 (2015), 135–153.
- [Br17] M. Brion, *Some structure theorems for algebraic groups*, Proc. Symp. Pure Math. 94 (2017), 53–125.
- [Ch55] W.-L. Chow, *Abelian varieties over function fields*, Trans. Amer. Math. Soc. 78 (1955), 253–275.
- [Co06] B. Conrad, *Chow’s K/k -image and K/k -trace, and the Lang-Néron theorem*, Enseign. Math. (2) 52 (2006), no. 1-2, 37–108.
- [CGP15] B. Conrad, O. Gabber, G. Prasad, *Pseudo-reductive groups. Second edition*, New Math. Monogr. 26, Cambridge Univ. Press, Cambridge, 2015.
- [CP15] B. Conrad, G. Prasad, *Classification of pseudo-reductive groups*, Ann. of Math. Stud. 191, Princeton Univ. Press, 2015.
- [DG70] M. Demazure, P. Gabriel, *Groupes algébriques*, Masson, Paris, 1970.
- [FJ74] G. Frey, M. Jarden, *Approximation theory and the rank of abelian varieties over large algebraic fields*, Proc. London Math. Soc. (3) 28 (1974), 112–128.

- [Ga62] P. Gabriel, *Des catégories abéliennes*, Bull. Soc. Math. France 90 (1962), 323–448.
- [GZ67] P. Gabriel, M. Zisman, *Calculus of fractions and homotopy theory*, Ergeb. Math. Grenzgeb. 35, Springer-Verlag, New York, 1967.
- [Ha01] D. Happel, *A characterization of hereditary categories with tilting object*, Invent. Math. 144 (2001), no. 2, 381–398.
- [Ho68] T. Honda, *Isogeny classes of abelian varieties over finite fields*, J. Math. Soc. Japan 20 (1968), 83–95.
- [Kr75] H. Kraft, *Kommutative algebraische Gruppen und Ringe*, Lecture Notes in Math. 455, Springer-Verlag, Berlin-New York, 1975.
- [LA15] G. Lucchini Arteche, *Extensions of algebraic groups with finite quotient*, preprint, arXiv:1503:06582.
- [MM74] B. Mazur, W. Messing, *Universal extensions and one dimensional crystalline cohomology*, Lecture Notes in Math. 370, Springer-Verlag, Berlin-New York, 1974.
- [MW93] D. Masser, G. Wüstholz: *Isogeny estimates for abelian varieties, and finiteness theorems*, Ann. Math. (2) 137 (1993), no. 3, 459–472.
- [Ma10] N. Mazzari, *Cohomological dimension of Laumon 1-motives up to isogenies*, J. Théor. Nombres Bordeaux 22 (2010), no. 3, 719–726.
- [Mi70] J. S. Milne, *The homological dimension of commutative group schemes over a perfect field*, J. Algebra 16 (1970), 436–441.
- [Mi86] J. S. Milne, *Abelian varieties*, in: Arithmetic geometry, 103–150, Springer-Verlag, New York, 1986.
- [Oo64] F. Oort, *Yoneda extensions in abelian categories*, Math. Annalen 153 (1964), no. 3, 227–235.
- [Oo66] F. Oort, *Commutative group schemes*, Lecture Notes in Math. 15, Springer-Verlag, Berlin-New York, 1966.
- [Or04] F. Orgogozo, *Isomotifs de dimension inférieure ou égale à un*, Manuscripta Math. 115 (2004), no. 3, 339–360.
- [Ra70] M. Raynaud, *Faisceaux amples sur les schémas en groupes et les espaces homogènes*, Lecture Notes in Math. 119, Springer-Verlag, Berlin-New York, 1970.
- [Ro56] M. Rosenlicht, *Some basic theorems on algebraic groups*, Amer. J. Math. 78 (1956), 401–443.

- [Ro58] M. Rosenlicht, *Extensions of vector groups by abelian varieties*, Amer. J. Math. 80 (1958), 685–714.
- [Ru70] P. Russell, *Forms of the affine line and its additive group*, Pacific J. Math. 32 (1970), 527–539.
- [SGA3] M. Demazure, A. Grothendieck, *Schémas en groupes I, II, III (SGA 3)*, Springer Lecture Notes in Math. 151, 152, 153 (1970); revised version edited by P. Gille and P. Polo, vols. I and III, Soc. Math. de France, Paris, 2011.
- [Sc72] C. Schoeller, *Groupes affines, commutatifs, unipotents sur un corps non parfait*, Bull. Soc. Math. France 100 (1972), 241–300.
- [Se59] J.-P. Serre, *Groupes algébriques et corps de classes*, Hermann, Paris, 1959.
- [Se60] J.-P. Serre, *Groupes proalgébriques*, Publ. Math. IHÉS 7 (1960).
- [SP16] *Stacks Project*, <http://stacks.math.columbia.edu>, 2016.
- [Tat66] J. Tate, *Endomorphisms of abelian varieties over finite fields*, Invent. Math. 2 (1966), 134–144.
- [Tak75] M. Takeuchi, *On the structure of commutative affine group schemes over a non-perfect field*, Manuscripta Math. 16 (1975), no. 2, 101–136.
- [To13] B. Totaro, *Pseudo-abelian varieties*, Ann. Sci. Éc. Norm. Sup. (4) 46 (2013), no. 5, 693–721.
- [Wa69] W. C. Waterhouse, *Abelian varieties over finite fields*, Ann. Sci. École Norm. Sup. (4) 2 (1969), 521–560.

Michel Brion
Université Grenoble Alpes
Institut Fourier
100 rue des maths
38610 Gières
France
Michel.Brion@ujf-grenoble.fr

